

JUNE EXAMINATION GRADE 12

2023

PHYSICAL SCIENCES (PHYSICS) (PAPER 1)

TIME: 3 hours

MARKS: 150

16 pages and 4 data sheets



INSTRUCTIONS AND INFORMATION


1. This question paper consists of 9 questions. Answer ALL the questions in the ANSWER BOOK.
2. Start the answers to each question on a NEW page in the ANSWER BOOK.
3. Number the answers correctly according to the numbering system used in this question paper.
4. Leave ONE line open between two sub-sections, for example between QUESTION 2.1 and QUESTION 2.2.
5. You may use a non-programmable calculator.
6. You may use appropriate mathematical instruments.
7. You are advised to use the attached DATA SHEETS.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round-off your final numerical answers to a minimum of TWO decimal places.
10. Give brief motivations, discussions, et cetera where required.
11. Write neatly and legibly.



QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A – D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, e.g. 1.11 E.

- 1.1 A box is shown resting on the ground. Newton's Third Law implies that four forces of equal magnitude are involved. These forces are labelled **P**, **Q**, **R** and **S**. Forces **P** and **Q** act on the box. Forces **R** and **S** act on the Earth. For clarity, the forces are shown slightly separated.



Which statement about the forces is correct?





- A Force **P** is equal and opposite to force **Q** and both are forces of contact.
 B Force **Q** is equal and opposite to force **P** and both are gravitational forces.
 C Force **R** is equal and opposite to force **S** and both are forces of contact.
 D Force **S** is equal and opposite to force **Q** and both are gravitational forces. (2)
- 1.2 A child on a sledge slides down a hill with acceleration **a**. The hill makes an angle θ with the horizontal. The total mass of the child and the sledge is **m**. The acceleration of free fall is **g**.



The frictional force **F** is represented by:

- A $m(g \cos \theta - a)$
 B $m(g \cos \theta + a)$
 C $m(g \sin \theta - a)$
 D $m(g \sin \theta + a)$ (2)

- 1.3 A small steel ball falls freely under gravity after being released from rest. Which graph best represents the variation of the height h of the ball with time t when using the ground as reference?



(2)


- 1.4 If air resistance is negligible, the total mechanical energy of a free-falling body ...

- A increases.
- B decreases.
- C becomes zero.
- D remains constant.

(2)



- 1.5 The diagram below shows two identical spheres **X** and **Y**. Initially, **X** moves with speed **v** directly towards **Y**, while **Y** is stationary. The spheres collide elastically.



The following occurs:

	X	Y
A	Stops	Moves with a speed v to the right
B	Moves with a speed v to the left	Remains stationary
C	Moves with a speed $\frac{1}{2}v$ to the left	Moves with a speed $\frac{1}{2}v$ to the right
D	Moves with a speed $\frac{1}{2}v$ to the right	Moves with a speed $\frac{1}{2}v$ to the right


(2)

- 1.6 The direction of an object's momentum is always the same as the direction of the object's ...

- A velocity.
- B weight.
- C inertia.
- D potential energy.

(2)

- 1.7 A satellite orbits Earth at a point **x**. The gravitational force on **x** is a quarter ($\frac{1}{4}$) of the gravitational force it experiences on the surface of the Earth.




If the radius of Earth is **R**, then the height of the satellite ABOVE THE SURFACE of Earth, will be:





- A 4R
- B 2R
- C R
- D $\frac{1}{2}R$

(2)

- 1.8 The warning siren on an ambulance has a frequency of 600 Hz. The speed of sound is $330 \text{ m}\cdot\text{s}^{-1}$ in air. The ambulance is travelling at a constant velocity of $25 \text{ m}\cdot\text{s}^{-1}$ towards an observer.




Which graph illustrates the change in observed frequency?



(2)

- 1.9 Which diagram below best represents the electric field pattern between two point charges of equal magnitude and opposite sign?



(2)

- 1.10 Which of the following represents the correct units for electric field strength?


- A T
 B $\text{N}\cdot\text{C}^{-1}$
 C $\text{J}\cdot\text{C}^{-1}$
 D $\text{N}\cdot\text{m}^2\cdot\text{C}^{-2}$

(2)

[20]


QUESTION 2 (Start on a new page.)

A log of mass m kg is pulled up a slope by a light non-elastic wire of negligible mass, attached to a motor, as shown in the diagram below.



The angle that the slope makes with the horizontal is 12° . The kinetic coefficient of friction between the log and the surface is 0,3. The tension in the wire is 1 570 N while the log moves up the slope with a constant velocity.

- 2.1 State Newton's First Law in words. (2)
- 2.2 Draw a labelled, free-body diagram to show all the forces acting on the log. (4)
- 2.3 Calculate the mass of the log. (5)
- 2.4 The angle of the slope is increased to 14° .




Identify the effect of this increased angle on each of the following:
(Write only INCREASES, DECREASES or REMAINS THE SAME.)

- 2.4.1 Kinetic coefficient of friction between the log and the surface (1)
- 2.4.2 Frictional force (1)
- 2.5 The wire snaps and the log slides down the slope until it stops.
Give a reason why this cannot be seen as free fall or projectile motion. (2)

[15]

QUESTION 3 (Start on a new page.)

A soft ball is thrown vertically down towards the ground whereafter it bounces back up as illustrated in the sketch below.



When the ball reaches point **A**, it has a speed of $8,4 \text{ m}\cdot\text{s}^{-1}$. The height of **A** is $5,0 \text{ m}$ above the ground. The ball hits the ground and bounces up to a maximum height at position **B**. Assume that the air resistance is negligible.

- 3.1 Explain the term *projectile motion*. (2)
- 3.2 Calculate the speed of the ball as it hits the ground. (3)
- 3.3 Show, by calculation that the time for the ball to reach the ground is $0,47 \text{ s}$. (3)
- 3.4 The ball bounces vertically upwards with a speed of $4,2 \text{ m}\cdot\text{s}^{-1}$ as it leaves the ground. The time the ball is in contact with the ground is $0,20 \text{ s}$.
 - 3.4.1 What is the magnitude and direction of the acceleration of the ball at point **B**? (2)
 - 3.4.2 Calculate the maximum height that the ball reaches after the first bounce off the floor. (3)
 - 3.4.3 Sketch a velocity time graph for the motion from the moment that the ball passes **A** until it reaches position **B**.

Clearly indicate the following on your graph:

- The velocity of the ball at **A**
- The velocity of the ball as it hits the ground
- Time taken for the ball to reach the ground **and** leave the ground
- The velocity at the maximum height of point **B** (6)

3.5 The soft ball is replaced by a very hard ball. How would this affect the following:
(Write only INCREASES, DECREASES or REMAINS THE SAME.)


3.5.1 The contact time when the ball bounces on the ground (1)

3.5.2 The height of the ball after it strikes the ground (2)
Explain the answer. [22]



QUESTION 4 (Start on a new page.)

Ball **X**, with a mass of 50 g, and ball **Y**, with a mass twice that of **X**, are supported by long strings from a ceiling, as shown in the diagram below.




The balls are each pulled back and released so that they move towards each other. When the balls collide at the position shown in the diagram above, the strings are vertical. The balls rebound in opposite directions. The velocity of ball **X** after the collision is $2,1 \text{ m}\cdot\text{s}^{-1}$ to the left. Ignore all effects of friction.

- 4.1 State the *principle of conservation of linear momentum*. (2)
- 4.2 Calculate the velocity of ball **Y** immediately after the collision. (5)
- 4.3 Is the collision ELASTIC or INELASTIC?
Explain the answer by means of suitable calculations. (5)
- 4.4 Ball **X** and **Y** are replaced by two identical balls. How would this affect the conservation of the momentum of the system if the initial velocities remain the same?
(Write only INCREASES, DECREASES or REMAINS THE SAME.)
Explain the answer. (3)

[15]

QUESTION 5 (Start on a new page.)

A ball of mass 0,03 kg moves along a curved track, as shown in the diagram below.



The speed of the ball is v when it is at point **A** at a height of 0,31 m. The ball moves down the frictionless track and collides with a vertical wall at point **B**. The kinetic energy of the ball just before the collision with the wall, is 0,12 J. After the collision the ball rebounds up the track. Ignore all frictional forces.

- 5.1 State the *principle of conservation of mechanical energy*. (2)
- 5.2 Calculate the speed v of the ball at point **A**. (4)
- 5.3 The change in momentum of the ball due to the collision with the wall is $0,096 \text{ kg}\cdot\text{m}\cdot\text{s}^{-1}$. The ball is in contact with the wall for a time of 0,02 s.

For the ball hitting the wall, calculate:


- 5.3.1 The speed immediately after the collision (4)
- 5.3.2 The magnitude of the average force on the wall (4)

[14]




QUESTION 6 (Start on a new page.)

The diagram below shows a fairground carriage moving at a constant speed of $9,5 \text{ m}\cdot\text{s}^{-1}$ over a frictionless surface. The total mass of the people and the carriage is 600 kg . The carriage moves up a slope inclined at 30° to the horizontal. The carriage comes to a stop after travelling up the rough slope to a vertical height of $4,1 \text{ m}$ at point **C**.



- 6.1 State the *work-energy theorem*, in words. (2)
- 6.2 Calculate the kinetic energy of the carriage and passengers as they reach point **B**. (3)
- 6.3 Draw a labelled, free-body diagram of all the forces that act on the carriage while moving up the slope. (3)
- 6.4 Use energy principles and calculate the frictional force between point **B** and **C**. (5)
- 6.5 A small tractor is rated at 9 kW and pulls an empty carriage across the ground at a constant speed with a horizontal force of 300 N .




- 6.5.1 Define a *non-conservative force*. (2)
- 6.5.2 Calculate the time that it would take the tractor to tow the empty carriage a distance of 850 m . (4)
- 6.5.3 How would the time change if a tractor with a greater power output was used?
(Write down only INCREASES, DECREASES or REMAINS THE SAME.)

Explain the answer.

(3)
[22]

QUESTION 7 (Start on a new page.)

Two charged spheres, **A** and **B** are placed 15 cm apart in a vacuum. Sphere **A** has a charge of +3,0 pC and sphere **B** has an unknown positive charge. The arrangement is illustrated in the diagram below.



Point **P** lies on the line joining the charged spheres at a distance of 5,0 cm from sphere **A**.

- 7.1 Describe an *electric field*. (2)
- 7.2 The net electric field strength at point **P** is zero. Calculate the charge of sphere **B**. (5)
- 7.3 State *Coulomb's Law* in words. (2)
- 7.4 Charge **B** is replaced by a negative charge of magnitude -12 pC. Calculate the magnitude of the electrostatic force between charged spheres **A** and **B**. (3)
- 7.5 How would the magnitude of the force calculated in QUESTION 7.4 be affected, if the distance between the charges is decreased?
(Write only INCREASES, DECREASES or REMAINS THE SAME.)

Explain the answer.


(3)
[15]

QUESTION 8 (Start on a new page.)

An experiment is conducted to investigate the relationship between the force experienced by a charge if the electric field strength is kept constant. The table of results is given below.

Force in N	0	0,05	0,1	0,15	0,2	0,25	0,28	0,35	X
Charge in ($\times 10^{-9}$ C)	0	7,6	15,2	22,8	30,4	38	42,6	53,2	63,8

The following graph is obtained.



8.1 Define the *electric field at a point*. (2)

8.2 Give the relationship between *force* and *charge*.

Explain the answer. (2)

8.3 Use the information above and calculate the:

8.3.1 Gradient of the graph (3)

8.3.2 Value of **X** (4)

8.4 Sketch the electric field pattern around the charge if it had been a negative charge. (3)

[14]

QUESTION 9 (Start on a new page.)

9.1 One way to tell if a mosquito is about to bite is to listen to the Doppler effect of the sound emitted by the mosquito as it is flying. The buzzing of a mosquito's wings emits sound at a frequency of 1 050 Hz. The speed of sound is $330 \text{ m}\cdot\text{s}^{-1}$ in air.

9.1.1 State the *Doppler effect for sound*, in words. (2)

9.1.2 If you hear a frequency of 1 034 Hz, does this mean that the mosquito is coming in for a landing or that it has just bitten you and is flying away? Explain your answer. (3)

9.1.3 Calculate the speed of the mosquito when you hear the frequency of 1 034 Hz. (5)

9.2 The Doppler effect has many other uses.

9.2.1 One of the uses of the Doppler effect is in the field of Astronomy. While studying the stars, a red shift emission is observed. Is the star moving away from the earth or towards the earth? (1)

9.2.2 Another use of the Doppler effect is in the medical field. Give TWO uses of the Doppler flow meter on humans. (2)

[13]

TOTAL: 150



DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 1 (PHYSICS)

TABLE 1: PHYSICAL CONSTANTS

NAME	SYMBOL	VALUE
Acceleration due to gravity	g	$9,8 \text{ m}\cdot\text{s}^{-2}$
Universal gravitational constant	G	$6,67 \times 10^{-11} \text{ N}\cdot\text{m}^2\cdot\text{kg}^{-2}$
Speed of light in a vacuum	c	$3,0 \times 10^8 \text{ m}\cdot\text{s}^{-1}$
Planck's constant	h	$6,63 \times 10^{-34} \text{ J}\cdot\text{s}$
Coulomb's constant	k	$9,0 \times 10^9 \text{ N}\cdot\text{m}^2\cdot\text{C}^{-2}$
Charge on electron	e	$1,6 \times 10^{-19} \text{ C}$
Electron mass	m_e	$9,11 \times 10^{-31} \text{ kg}$
Mass of earth	M	$5,98 \times 10^{24} \text{ kg}$
Radius of earth	R_E	$6,38 \times 10^6 \text{ m}$



TABLE 2: FORMULAE

MOTION

$v_f = v_i + a\Delta t$	$\Delta x = v_i\Delta t + \frac{1}{2}a\Delta t^2$ or $\Delta y = v_i\Delta t + \frac{1}{2}a\Delta t^2$
$v_f^2 = v_i^2 + 2a\Delta a$ or $v_f^2 = v_i^2 + 2a\Delta a$	$\Delta x = \left(\frac{v_i + v_f}{2}\right)\Delta t$ or $\Delta y = \left(\frac{v_i + v_f}{2}\right)\Delta t$

FORCE

$F_{\text{net}} = ma$	$p = mv$
$f_s^{\text{max}} = \mu_s N$	$f_k = \mu_k N$
$F_{\text{net}}\Delta t = \Delta p$ $\Delta p = mv_f - mv_i$	$w = mg$
$F = G \frac{m_1 m_2}{d^2}$ or $F = G \frac{m_1 m_2}{r^2}$	$g = G \frac{M}{d^2}$ or $g = G \frac{M}{r^2}$

WORK, ENERGY AND POWER

$W = F\Delta x \cos\theta$	$U = mgh$ or $E_p = mgh$
$K = \frac{1}{2}mv^2$ or $E_k = \frac{1}{2}mv^2$	$W_{\text{net}} = \Delta K$ or $W_{\text{net}} = \Delta E_k$
	$\Delta K = K_f - K_i$ or $\Delta E_k = E_{kf} - E_{ki}$
$W_{\text{nc}} = \Delta K + \Delta U$ or $W_{\text{nc}} = \Delta E_k + \Delta E_p$	$P = \frac{W}{\Delta t}$
$P_{\text{ave}} = Fv_{\text{ave}}$	



WAVES, SOUND AND LIGHT

$v = f \lambda$	$T = \frac{1}{f}$
$f_L = \frac{v \pm v_L}{v \pm v_b} f_b$	$E = hf \quad \text{or} \quad E = h \frac{c}{\lambda}$
$E = W_0 + E_{k(\max)}$ or $E = W_0 + K_{\max}$ where $E = hf$ and $W_0 = hf_0$ and $E_{k(\max)} = \frac{1}{2}mv_{\max}^2$ or $K_{\max} = \frac{1}{2}mv_{\max}^2$	

ELECTROSTATICS

$F = \frac{kQ_1Q_2}{r^2}$	$E = \frac{kQ}{r^2}$
$V = \frac{W}{q}$	$E = \frac{F}{q}$
$n = \frac{Q}{e}$ or $n = \frac{Q}{q_e}$	




ELECTRIC CIRCUITS

$R = \frac{V}{I}$	$\text{emf } (\mathcal{E}) = I(R + r)$
$R_s = R_1 + R_2 + \dots$ $\frac{1}{R_p} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$	$q = I\Delta t$
$W = Vq$ $W = VI\Delta t$ $W = I^2R\Delta t$ $W = \frac{V^2\Delta t}{R}$	$P = \frac{W}{\Delta t}$ $P = VI$ $P = I^2R$ $P = \frac{V^2}{R}$

ALTERNATING CURRENT

$I_{\text{rms}} = \frac{I_{\text{max}}}{\sqrt{2}}$ $V_{\text{rms}} = \frac{V_{\text{max}}}{\sqrt{2}}$	$P_{\text{ave}} = V_{\text{rms}} I_{\text{rms}}$ $P_{\text{ave}} = I_{\text{rms}}^2 R$ $P_{\text{ave}} = \frac{V_{\text{rms}}^2}{R}$
--	--





**JUNE EXAMINATION
*JUNIE EKSAMEN***

GRADE/GRAAD 12

2023

**MARKING GUIDELINES/
*NASIENRIGLYNE***

**PHYSICAL SCIENCES: PHYSICS/
*FISIESE WETENSKAPPE: FISIKA***

(PAPER/VRAESTEL 1)

14 pages/*bladsye*



QUESTION/VRAAG 1

- | | | |
|------|------|-----|
| 1.1 | D ✓✓ | (2) |
| 1.2 | C ✓✓ | (2) |
| 1.3 | B ✓✓ | (2) |
| 1.4 | D ✓✓ | (2) |
| 1.5 | A ✓✓ | (2) |
| 1.6 | A ✓✓ | (2) |
| 1.7 | C ✓✓ | (2) |
| 1.8 | C ✓✓ | (2) |
| 1.9 | D ✓✓ | (2) |
| 1.10 | B ✓✓ | (2) |

[20]





QUESTION/VRAAG 2

- 2.1 A body continues at rest or constant velocity unless acted on by a net (external) force. ✓✓

’n Liggaam bly in rus of beweeg teen konstante snelheid tensy ’n nie-nul resultante (netto eksterne) krag daarop inwerk.

(2)

2.2



OR/OF

Accepted labels/Aanvaarde byskrifte	
w ✓	F_g/F_w /weight/mg/gravitational force F_g/F_w /gewig/ gravitasiekrag
f ✓	F_{friction}/F_f /friction/ f_k F_{wrywing}/F_f / wrywing/ f_k
N ✓	F_N/F_{normal} /normal force F_N/F_{normaal} /normaalkrag
T ✓	F_T /tension F_T /spanning/ spankrag
	Deduct 1 mark for any additional force. Marks are given for both arrow and label. If everything is correct, but no arrows, deduct a mark.
	<i>Trek een punt af vir enige addisionele kragte. Punte word gegee vir beide pylpunt en byskrif. Indien alles korrek is, maar geen pyle, trek ’n punt af.</i>

(4)

- 2.3 In x-direction:/In x-rigting:

$$F_{\text{net}} = ma \quad \checkmark$$

$$T - f_k - W_{II} = 0$$

$$1\,570 - 0,3(m)(9,8)\cos 12^\circ \checkmark - m(9,8)\sin 12^\circ \checkmark = 0 \quad \checkmark$$

$$1\,570 = 2,876\,m + 2,039\,m$$

$$1\,570 = 4,914\,m$$

$$m = 319,495\,kg \quad \checkmark$$

(5)

- 2.4 2.4.1 REMAINS THE SAME ✓ *BLY DIESELFDE* (1)
- 2.4.2 DECREASES ✓ *VERLAAG* (1)
- 2.5 There is an external force of friction exerted ✓ on the body and not only gravitational force. ✓

Daar word 'n eksterne krag van wrywing toegepas op die voorwerp en nie slegs gravitasiekrag nie.

(2)
[15]

QUESTION/VRAAG 3

- 3.1 A moving object that is only influenced by gravity and starts with an initial velocity. ✓✓

'n Bewegende voorwerp wat slegs onder die invloed van gravitasie beweeg en begin met 'n aanvanklike snelheid.

(2)

3.2 $v_f^2 = v_i^2 + 2a\Delta y$ ✓
 $= (8,4)^2 + 2(9,8)(5)$ ✓
 $v_f = 12,983 \text{ m} \cdot \text{s}^{-1}$ ✓

(3)

3.3 OPTION 1:/OPSIE 1

$$\Delta y = v_i \Delta t + \frac{1}{2} a \Delta t^2$$
 ✓

$$5 \text{ ✓} = 8,4t + \frac{1}{2}(9,8)t^2$$
 ✓

$$t = 0,47 \text{ s}$$

OPTION 2:/OPSIE 2

$$\Delta y = \left(\frac{v_f + v_i}{2} \right) \Delta t$$
 ✓

$$5 \text{ ✓} = \left(\frac{8,4 + 12,983}{2} \right) t$$
 ✓

$$t = 0,47 \text{ s}$$

OPTION 3:/OPSIE 3 Positive marking from 3.2/Positiewe nasien vanaf 3.2

$$v_f = v_i + a \Delta t$$
 ✓

$$12,983 \text{ ✓} = 8,4 + (9,8)t$$
 ✓

$$t = 0,47 \text{ s}$$

(3)

3.4 3.4.1 $9,8 \text{ m}\cdot\text{s}^{-2}$ ✓ Downwards ✓
Afwaarts

(2)


3.4.2 $v_f^2 = v_i^2 + 2a\Delta y$ ✓

$0 = (-4,2)^2 + 2(9,8)y$ ✓

$y = 0,9 \text{ m}$ ✓

(3)

3.4.3 Taking down as positive/Neem af as positief



Marking guidelines:


- ✓ The velocity of the ball at **A**
- ✓ The velocity of the ball as it hits the ground
- ✓ Time taken for the ball to reach the ground **and** ✓ leave the ground
- ✓ The velocity at the maximum height of point **B**
- ✓ Correct shape – parallel lines

Nasienriglyne:

- ✓ Die snelheid van die bal by **A**.
- ✓ Die snelheid van die bal wanneer dit die grond tref
- ✓ Die tyd wat dit neem vir die bal om die grond te bereik **en** ✓ verlaat die grond.
- ✓ Die snelheid by die maksimum hoogte van punt **B**.
- ✓ Korrekte vorm – parallel lyne.

Taking down as negative./Neem afwaarts as positief.

(6)



3.5.1 DECREASES ✓ AFNEEM

(1)

3.5.2 INCREASES ✓

The contact time is less, the force is more, the final velocity is more.
Higher kinetic energy ✓ (any one of the answers)

TOENEEM

*Die kontaktyd is minder, die krag is groter, die finale snelheid is meer.
Hoër kinetiese energie (enige een van die antwoorde)*

(2)

[22]

QUESTION/VRAAG 4

4.1 In an isolated system the total linear momentum remains constant. ✓✓


In 'n geïsoleerde sisteem bly die totale lineêre momentum konstant.

Marking criteria/Nasienkriteria

If any of the underlined key words/phrases in the correct context is omitted deduct 1 mark./Indien enige van die onderstreepte sleutelwoorde/frases in die korrekte konteks uitgelaat is, trek 1 punt af.

(2)



4.2 OPTION 1:/OPSIE 1:

$$\begin{aligned}\sum p_{\text{before/ voor}} &= \sum p_{\text{after/ na}} \checkmark \\ m_x v_{xi} + m_y v_{yi} &= m_x v_{xf} + m_y v_{yf} \\ m(4,5) + 2m(-2,8) \checkmark &= m(-2,1) + 2mv_{yf} \checkmark \\ v_{yf} &= 0,5 \text{ m.s}^{-1} \checkmark \text{ to the right } \checkmark\end{aligned}$$

OPTION 2:/OPSIE 2:

$$\begin{aligned}\sum p_{\text{before/ voor}} &= \sum p_{\text{after/ na}} \checkmark \\ m_x v_{xi} + m_y v_{yi} &= m_x v_{xf} + m_y v_{yf} \\ 0,05(4,5) + 0,1(-2,8) \checkmark &= 0,05(-2,1) + 0,01v_{yf} \checkmark \\ v_{yf} &= 0,5 \text{ m.s}^{-1} \checkmark \text{ to the right } \checkmark\end{aligned} \quad (5)$$

4.3

$$\begin{aligned}\Sigma E_k \text{ before/ voor} &= \frac{1}{2}mv^2 + \frac{1}{2}mv^2 \checkmark \\ &= (0,5\text{m})(4,5)^2 + (0,5(2\text{ m}))(2,8)^2 \checkmark \\ &= 10,125 \text{ m} + 7,84 \text{ m} \\ &= 17,965 \text{ m J}\end{aligned}$$

$$\begin{aligned}\Sigma E_k \text{ after/ na} &= \frac{1}{2}mv^2 + \frac{1}{2}mv^2 \\ &= (0,5\text{m})(2,1)^2 + (0,5(2\text{ m}))(0,5)^2 \checkmark \\ &= 2,205 \text{ m} + 0,25 \text{ m} \\ &= 2,455 \text{ m J}\end{aligned}$$

$\Sigma E_k \text{ before/ voor} \neq \Sigma E_k \text{ after/ na} \checkmark$
INELASTIC COLLISION ✓/ ONELASTIESE BOTSING

(5)

4.4 REMAINS THE SAME ✓

The system remains isolated, ✓ all momentum before and after the collisions will remain the same. There are no external forces. ✓

BLY DIESELFDE

Die sisteem bly geïsoleerd, al die momentum voor en na die botsings is dieselfde. Daar is geen eksterne kragte nie.

(3)

[15]

QUESTION/VRAAG 5

- 5.1 The total mechanical energy at any given point is the same and conserved in a closed system. ✓✓

Die totale meganiiese energie by enige gegewe oomblik is dieselfde en bly behoue in 'n geslote sisteem.

(2)

- 5.2 E mechanical /meganies = E mechanical /meganies

$$(E_k + E_p)_A = (E_k + E_p)_B \quad \checkmark$$

$$\left(\frac{1}{2}mv^2 + mgh\right)_A = \left(\frac{1}{2}mv^2 + mgh\right)_B$$

$$\checkmark \frac{1}{2}(0,030)v^2 + (0,03)(9,8)(0,31) = 0,12 + 0 \quad \checkmark$$

$$v = 1,387 \text{ m}\cdot\text{s}^{-1} \quad \checkmark$$

accept/ aanvaar $v = 1,39\text{m}\cdot\text{s}^{-1}$

(4)

- 5.3 5.3.1

$$E_k = \frac{1}{2}mv^2$$

$$0,12 = \frac{1}{2}(0,03)v^2$$

$$v = 2,828 \text{ m}\cdot\text{s}^{-1}$$

$$F_{\text{net}}\Delta t = \Delta p \quad \checkmark$$

$$\Delta p = m(v_f - v_i)$$

$$-0,096 \checkmark = 0,03(v_f - 2,828) \checkmark$$

$$v_f = -0,372 \text{ m}\cdot\text{s}^{-1}$$

$$v = 0,372 \text{ m}\cdot\text{s}^{-1} \quad \checkmark$$

(4)

- 5.3.2

$$F_{\text{net}}\Delta t = \Delta p \quad \checkmark$$

$$F_{\text{net}}(0,02) \checkmark = 0,096 \checkmark$$

$$F_{\text{net}} = 4,8 \text{ N} \quad \checkmark$$

OR/OF

$$F_{\text{net}}\Delta t = \Delta p \quad \checkmark$$

$$F_{\text{net}}(0,02) \checkmark = -0,096 \checkmark$$

$$F_{\text{net}} = -4,8 \text{ N}$$

$$F_{\text{net}} = 4,8 \text{ N} \quad \checkmark$$

(4)

[14]

QUESTION/VRAAG 6

- 6.1 The work done on an object by a net force is equal to the change in the object's kinetic energy. ✓✓

Die arbeid verrig op 'n voorwerp deur 'n netto krag is gelyk aan die verandering in die voorwerp se kinetiese energie.

(2)

6.2



$$K = \frac{1}{2} m v^2 \quad \checkmark$$

$$= \frac{1}{2} (600)(9,5)^2 \quad \checkmark$$

$$= 27\,075 \text{ J} \quad \checkmark$$

(3)

6.3



Accepted labels/Aanvaarde byskrifte	
w ✓	F_g/F_w /weight/mg/gravitational force F_g/F_w /gewig/ gravitasiekrag
f ✓	F_{friction}/F_f /friction/ f_k F_{wrywing}/F_f / wrywing/ f_k
N ✓	F_N/F_{normal} /normal force F_N/F_{normaal} /normaalkrag
	Deduct 1 mark for any additional force. Marks are given for both arrow and label. If everything is correct, but no arrows, deduct a mark.
	<i>Trek een punt af vir enige addisionele kragte. Punte word gegee vir beide pylpunt en byskrif. Indien alles korrek is, maar geen pyle, trek 'n punt af.</i>

(3)

6.4 OPTION 1:/OPSIE 1

$$W_{nc} = \Delta K + \Delta U \quad \text{or/of} \quad W_{nc} = \Delta E_k + \Delta E_p \quad \checkmark$$

$$W_f = -\frac{1}{2}mv_i^2 + mgh$$

$$f\Delta x \cos\theta = -27\,075 + 600(9,8)(4,1) \quad \checkmark$$

$$f\left(\frac{h}{\sin 30^\circ}\right) \checkmark \cos 180^\circ \checkmark = -27\,075 + 24\,108$$

$$f\left(\frac{4,1}{\sin 30^\circ}\right) \cos 180^\circ = -2\,967$$

$$f = \frac{-2\,967}{-8,2}$$

$$= 361,83 \text{ N} \quad \checkmark$$

OPTION 2:/OPSIE 2

$$W_{net} = \Delta K \quad \checkmark$$

$$W_N + W_f + W_w = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

$$0 + f\Delta x \cos\theta + mg\Delta x \cos 120^\circ = 0 - 27\,075 \quad \checkmark$$

$$f\left(\frac{4,1}{\sin 30^\circ}\right) \checkmark \cos 180^\circ \checkmark + (600)(9,8)\left(\frac{4,1}{\sin 30^\circ}\right) \cos 120^\circ = -27\,075$$

$$f = 361,83 \text{ N} \quad \checkmark$$

OPTION 3:/OPSIE 3

$$W_{net} = \Delta K \quad \checkmark$$

$$W_N + W_f + W_{w\perp} + W_{w\parallel} = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

$$0 + f\Delta x \cos\theta + 0 + mg\sin 30^\circ \Delta x \cos 0^\circ = 0 - 27\,075$$

$$f\left(\frac{4,1}{\sin 30^\circ}\right) \checkmark \cos 180^\circ \checkmark + (600)(9,8)\left(\frac{4,1}{\sin 30^\circ}\right) \cos 120^\circ \checkmark = -27\,075$$

$$f = 361,83 \text{ N} \quad \checkmark$$

(5)

- 6.5 6.5.1 A force for which the work done in moving an object between two points depends on the path taken. ✓✓



’n Krag waarvoor die arbeid wat verrig word om die voorwerp te beweeg tussen twee punte afhang van die padroete gevolg.

(2)

6.5.2 $W_F = F \Delta x \cos \theta$

$$= (300)(850) \cos 0^\circ \checkmark$$

$$= 255\,000 \text{ J}$$

$$P = \frac{W}{\Delta t} \checkmark$$

$$9000 = \frac{255000}{t} \checkmark$$

$$t = 28,33 \text{ s} \checkmark$$

(4)

- 6.5.3 DECREASES ✓

The work done on the carriage stays the same. ✓

The time and power are inversely proportional. If the power increases, then the time will decrease. ✓

VERLAAG

Die arbeid verrig op die karretjie bly dieselfde.

Die tyd en drywing is omgekeerd eweredig. Indien die drywing verhoog, dan sal die tyd verlaag.

(3)

[22]

QUESTION/VRAAG 7

- 7.1 An electric field is a region or space in which an electric charge experiences a force. ✓✓

’n Elektriese veld is ’n area of gebied waarin ’n elektriese lading ’n krag sal ondervind.

(2)

7.2

$$E_{\text{net}} = \frac{kQ_A}{r^2} - \frac{kQ_B}{r^2} \quad \checkmark$$

$$0 = \frac{9 \times 10^9 (3 \times 10^{-12})}{(0,05)^2} \checkmark - \checkmark \frac{(9 \times 10^9) Q_B}{(0,10)^2} \checkmark$$

$$Q_B = 1,2 \times 10^{-11} \text{ C} \quad \checkmark$$

(5)

- 7.3 The magnitude of the electrostatic force exerted by one point charge (Q_1) on another point charge (Q_2), is directly proportional to the product of the magnitudes of the charges and inversely proportional to the square of the distance (r) between them. ✓✓

Die grootte van die elektrostatische krag uitgeoefen deur een puntlading (Q_1) op ’n ander puntlading (Q_2) is direk eweredig aan die produk van die groottes van die ladings en omgekeerd eweredig aan die kwadraat van die afstand (r) tussen hulle.

(2)

7.4

$$F = \frac{kQ_A Q_B}{r^2} \quad \checkmark$$

$$= \frac{9 \times 10^9 (3 \times 10^{-12})(12 \times 10^{-12})}{(0,15)^2} \quad \checkmark$$

$$= 1,44 \times 10^{-11} \text{ N} \quad \checkmark$$

(3)

7.5 INCREASES ✓

The force is inversely proportional to the squared distance between the charges. ✓

The charges remain the same. ✓

VERHOOG

Die krag is omgekeerd eweredig aan die kwadraat van die afstand tussen die ladings.

Die ladings bly dieselfde.

(3)

[15]

QUESTION/VRAAG 8

- 8.1 The electric field at a point is the electrostatic force experienced per unit of positive charge placed at that point. ✓✓

Die elektriese veld by 'n punt is die elektrostatiese krag ondervind per eenheid van positiewe lading geplaas by daardie punt.

(2)

- 8.2 Force is directly proportional to charge. ✓
OR Charge is directly proportional to force.

Straight line through the origin. ✓

*Krag is direk eweredig aan die lading.
OF Lading is direk eweredig aan die krag.*

Reguit lyn deur die oorsprong.

(2)

- 8.3 8.3.1 Any values in the table may be used./*Enige waardes in die tabel kan gebruik word.*

$$\begin{aligned} \text{gradient/g radiënt} &= \frac{\Delta y}{\Delta x} \\ &= \frac{0,35 - 0}{53,2 \times 10^{-9} - 0} \checkmark \\ &= 6,58 \times 10^6 \text{ (N.C}^{-1}\text{)} \checkmark \end{aligned}$$

(3)


- 8.3.2 Positive marking from 8.3.1/*Positiewe nasien vanaf 8.3.1*

$$\begin{aligned} E &= \frac{F_x}{q} \checkmark \\ 6,58 \times 10^6 \checkmark &= \frac{F_x}{63,8 \times 10^{-9}} \checkmark \\ F_x &= 0,42 \text{ N} \checkmark \end{aligned}$$

(4)



8.4

Marking guidelines:

- ✓ Arrows towards the charge
- ✓ All lines touch the charge at an angle of 90°
- ✓ Correct pattern and around the charge.

Nasienriglyne:

- ✓ Pyle na die lading
- ✓ Alle lyne raak die lading teen 'n hoek van 90°
- ✓ Korrekte patroon en rondom die lading

(3)
[14]**QUESTION/VRAAG 9**

- 9.1 9.1.1 The change in the observed frequency (or pitch) of the sound detected by a listener ✓ because the sound source and the listener have different velocities relative to the medium of sound propagation. ✓

Die verandering in die waarneembare frekwensie (of toonhoogte) van die klank waargeneem deur 'n luisteraar aangesien die klankbron en die luisteraar verskillende snelhede relatief tot die klankbeweging het.

(2)

- 9.1.2 The mosquito is flying away from you. ✓ The observed frequency is ✓ lower than the actual frequency. ✓

Die muskiet vlieg weg van jou af. Die waargenome frekwensie is laer as die werklike frekwensie.

(3)

9.1.3

$$f_L = \frac{v \pm v_L}{v \pm v_s} f_s \quad \checkmark$$

$$1034 \checkmark = \frac{330 + 0}{330 + v_s} \checkmark 1050 \checkmark$$

$$v_s = 5,106 \text{ m} \cdot \text{s}^{-1}$$

$$v_s = 5,11 \text{ m} \cdot \text{s}^{-1} \quad \checkmark$$



(5)

- 9.2 9.2.1 Away from the earth ✓ *Van die aarde af weg te beweeg*

(1)

- 9.2.2 – It monitors foetal heart beat ✓ (Do not accept observing the baby.)
Dit monitor die fetus se hartklop. (Moenie aanvaar om baba te sien nie.)
- It detects flow of blood. ✓
Dit neem bloedvloei waar.

(2)

[13]

TOTAL/TOTAAL: 150