




LIMPOPO  
PROVINCIAL GOVERNMENT  
REPUBLIC OF SOUTH AFRICA


DEPARTMENT OF  
**EDUCATION**

NATIONAL  
SENIOR CERTIFICATE



MARKS: 150

TIME: 3 HOURS



MEMATHP2



This question paper consists of 10 pages and an information sheet.

**INSTRUCTIONS AND INFORMATION**

**Read the following instructions carefully before answering the questions.**

1. This question paper consists of 10 questions.
2. Answer ALL the questions.
3. Clearly show ALL calculations, diagrams, graphs, et cetera that you have used in determining your answers.
4. ANSWERS ONLY will not necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round answers off to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. Number the answers correctly according to the numbering system used in this question paper.
9. Write legibly and present your work neatly.



**QUESTION 1**

In a survey, a student asked his friends and family how many marketing phone calls they receive per month. The following table represent the data:

AGE OF FRIENDS AND FAMILY	FREQUENCY	CUMULATIVE FREQUENCY
$20 < x \leq 30$	7	7
$30 < x \leq 40$		27
$40 < x \leq 50$	25	
$50 < x \leq 60$		64
$60 < x \leq 70$		72
$70 < x \leq 80$	4	
$80 < x \leq 90$		80

- 1.1 Complete the Cumulative frequency table in the ANSWER BOOK. (3)
- 1.2 How many people took part in the survey? (1)
- 1.3 Write down the modal class. (1)
- 1.4 Draw the ogive in the ANSWER BOOK. (3)
- 1.5 Determine the percentage of marketing calls a person older than 54 will receive. (3)

[11]

**QUESTION 2**


The numbers {10, 20, 45, 58, 80} are the 5 number summary of a set of data with nine numbers. The second and third number of the data set are the same. The eighth number is 3 times the first quartile. The fourth number is equal to the interquartile range. The mean for the data set is 42.

- 2.1 Draw the box and whisker diagram in the ANSWER BOOK. (2)
- 2.2 Comment on the skewness of the data. (1)
- 2.3 Write down a possible list of nine numbers which will result in the above box and whisker plot. (6)

[9]

**QUESTION 3**

In the diagram  $P(-4; 2)$ ,  $Q(4; \alpha)$ ,  $R(2; -1)$  and  $S(-2; -3)$  are the vertices of trapezium PQRS with  $PQ \parallel SR$ .





- 3.1 Calculate the value of  $\alpha$ . (4)
- 3.2 Find the equation of the line passing through P and R. (3)
- 3.3 Find the size of  $\hat{P}RQ$ . (5)
- 3.4 If  $A(-1; t)$  and P, A and R are collinear, calculate the value of  $t$ . (2)


[14]

**QUESTION 4**

In the diagram the circle  $P$  with equation  $(x - 2)^2 + (y + 3)^2 = 34$  is drawn. The point  $A(4 ; 5)$  is a point on the tangent defined by  $5y = 3x + 13$ .  $B(a ; b)$  is the point on the circumference where the tangent touches the circle.



- 4.1 Determine the coordinates of  $P$  and the value of the radius of the circle. (2)
- 4.2 Write down the gradient of the radius  $PB$  in terms of  $a$  and  $b$ . (2)
- 4.3 Determine the gradient of the tangent  $AB$  in terms of  $a$  and  $b$ . (1)
- 4.4 Hence or otherwise, show that the values of  $a = -1$  and  $b = 2$ . (6)
- 4.5 Calculate the length of  $AB$ . Leave the answer in surd form. (2)
- 4.6 A second tangent is drawn from  $A$  to touch the circle at  $C(d ; 0)$ . Determine the value of  $d$ . (4)
- 4.7 Prove that  $\hat{BPC} = 90^\circ$ . (3)
- 4.8 If  $AB$  intersects the  $y$ -axis at  $D$  and  $CP$  produced intersects the  $y$ -axis at  $E$ , calculate the length of  $DE$ . (5)



**QUESTION 5**

 5.1 If  $\tan 35^\circ = m$ , determine the value of the following in terms of  $m$ :

5.1.1  $\sin 215^\circ$  (3)

5.1.2  $\sin 70^\circ$  (3)

5.2  $\tan \alpha = \frac{1}{3}$  and  $\tan \beta = \frac{1}{7}$ , where  $\alpha$  and  $\beta$  are acute angles.

Determine the value of:  $50 \times \sin(2\alpha + \beta)$ . (6)

5.3 Simplify: 
$$\frac{\cos(-x) \tan(180^\circ - x) \cos(90^\circ - x)}{\sin(540^\circ + x) \sin(180 - x)}$$
 (6)

5.4 5.4.1 Prove that: 
$$\frac{\sin x}{1 - \sin x} + \frac{\sin x}{1 + \sin x} = \frac{2 \tan x}{\cos x}$$
 (4)

5.4.2 For which value(s) of  $x$  in the interval  $[0^\circ ; 180^\circ]$  will the identity in 5.3.1 not be defined. (2)

[24]

**QUESTION 6**

$$f(x) = \cos(x - 30^\circ) \text{ and } g(x) = \sin x, \quad x \in [-90^\circ ; 90^\circ]$$

6.1 If  $x \in [-90^\circ ; 90^\circ]$ , solve the equation  $f(x) = g(x)$ . (4)

6.2 6.2.1 Draw the graphs of  $f(x)$  and  $g(x)$  on the same set of axes provided in the ANSWER BOOK. (4)


Hence use the graphs to answer the following questions:

6.2.2 Write down the range of  $f$ . (2)

6.2.3 For which values of  $x$  will  $f(x)$  decrease if  $x$  increase. (2)

6.2.4 If  $x \in [-90^\circ ; 0^\circ]$ , for which values of  $x$  will  $f(x) \cdot g(x) \geq 0$ ? (3)


6.2.5 Write down the value or of  $x$  where  $g(x) > f(x)$ . (2)



[17]

**QUESTION 7**

In the diagram AF is a tangent. AEC is a line intersecting the circle at E. BD is a diameter. EB, CB and CD are joined.  $\hat{A}BE = 2x$  and  $AB = EB = h$ .




- 7.1      7.1.1     Write down, with reasons, the size of  $\hat{A}$  in terms of  $x$ . (2)  
 7.1.2    Write down, with reasons, the size of  $A\hat{C}B$  in terms of  $x$ . (2)

7.2      Show that  $BC = \frac{h}{2\sin x}$ . (3)

7.3      Hence show with reasons that  $BD = \frac{h}{\sin 2x}$ . (3)

**QUESTION 8**

In the diagram PT and PQ are tangents to the circle at T and Q.  $KP \parallel QA$ . Chords TQ and PK intersect at R.  $\hat{T_2} = 50^\circ$



Find, with reasons, the sizes of:


- 8.1  $\hat{A}$  (2)
- 8.2  $\hat{K}_1$  (2)
- 8.3  $\hat{Q}_3$  (2)
- 8.4 Give a reason why TKQP a cyclic quadrilateral is. (1)
- 8.5  $\hat{Q}_1$  (3)

[10]

### QUESTION 9


9.1 In the diagram O is the centre of the circle and KLMN is a cyclic quadrilateral.

Prove the theorem stating that  $\hat{K} + \hat{M} = 180^\circ$ .



(5)

9.2 In the diagram below, WEG and VFH are tangents to the circle. EJ is drawn such that  $EJ = EF$ . WF cuts the circle at K. EK produced meets VH at V, JK cuts EF at L.



Prove that:

$$9.2.1 \quad \hat{EJF} = \hat{K}_2 \quad (4)$$


$$9.2.2 \quad \hat{K}_1 = \hat{K}_2 \quad (2)$$

9.2.3 KLFV is a cyclic quadrilateral. (3)

|14|

**QUESTION 10**


In the diagram PQ is a tangent to the circle at Q. T, R, A and B are points in the circumference of the circle and S is a point outside the circle. RT produced meets the tangent at P. TAS and RBS are straight lines.  $PT = 4$  units,  $TR = 5$  units and  $PS \parallel AB$ .



- 10.1 If  $\hat{S}_1 = x$ , name with reasons, 2 other angles equals to  $x$ . (4)  
10.2 Prove that  $\triangle PQT \sim \triangle PRQ$ . (3)  
10.3 Calculate the length of PQ. (3)  
10.4 Prove that  $\triangle PTS \sim \triangle PSR$ . (3)  
10.5 Prove that  $PQ = PS$  (3)

**[16]****GRAND TOTAL:** **150**

INFORMATION SHEET: MATHEMATICS

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n-1)d$$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r-1} ; r \neq 1$$

$$S_r = \frac{a}{1-r} ; -1 < r < 1$$

$$F = \frac{x[(1+i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1+i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \quad \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2} ab \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$


$$\bar{x} = \frac{\sum fx}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$



**LIMPOPO**  
PROVINCIAL GOVERNMENT  
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF  
**EDUCATION**

NATIONAL  
SENIOR CERTIFICATE

**GRADE/GRAAD 12**

MATHEMATICS P2/WISKUNDE 2

JUNE 2024

MARKING GUIDELINES/NASIENRIGLYNE

*Stanmorephysics.com*

**MARKS/PUNTE: 150**

This marking guidelines consist of 19 pages/Hierdie nasienriglyne bestaan uit 19 bladsye.



**QUESTION/VRAAG 1**

1.1	AGE OF FRIENDS AND FAMILY	FREQUENCY	CUMULATIVE FREQUENCY		
	$20 < x \leq 30$	7	7		
	$30 < x \leq 40$	<b>20</b>	27		
	$40 < x \leq 50$	25	<b>52</b>		
	$50 < x \leq 60$	<b>12</b>	64		
	$60 < x \leq 70$	<b>8</b>	72		
	$70 < x \leq 80$	4	<b>76</b>		
	$80 < x \leq 90$	<b>4</b>	80		
1.2	80			✓ 20 & 52	
1.3	$40 < x \leq 50$			✓ 12 & 8	
1.4	<p style="text-align: center;"><b>Ages of family and friends</b></p>				✓ 76 & 4 (3)
1.5	$\frac{21}{80} \times 100 = 26,25\%$			✓ answer (1)	
				✓ answer (1)	
				✓ grounding ✓ end point ✓ 2 points (3)	
					[11]

## QUESTION/VRAAG 2

2.1	<p>A box plot on a number line from 0 to 90. The minimum is at 10, the Q1 is at 20, the median is at 45, the Q3 is at 55, and the maximum is at 80.</p>	✓✓ plotting values	(2)
2.2	Skew to the left OR negatively skew	✓ answer	(1)
2.3	$2^{\text{nd}} = 20$ $4^{\text{th}} = 38$ $8^{\text{th}} = 60$ $7^{\text{th}} = 56$ $6^{\text{th}} : \frac{10 + 20 + 20 + 38 + 45 + x + 56 + 60 + 80}{9} = 42$ $\frac{329 + x}{9} = 42$ $329 + x = 378$ $x = 49$	✓ 20 ✓ 32 ✓ 60 ✓ 56 ✓ $\frac{329 + x}{9} = 42$ ✓ answer	(6)
			[9]


QUESTION/VRAAG 3




3.1	$m(PQ) = m(SR)$ $\frac{2-a}{-4-4} = \frac{-3+1}{-2-2}$ $\frac{2-a}{-8} = \frac{1}{2}$ $4-2a = -8$ $-2a = -12$ $a = 6$	✓ $m(PQ)$ ✓ $m(SR)$ ✓ equating ✓ answer	(4)
3.2	$m(PR) = \frac{2+1}{-4-2} = -\frac{1}{2}$ $y = mx + c$ $2 = \left(-\frac{1}{2}\right)(-4) + c$ $c = 0$ $y = -\frac{1}{2}x$  OR	✓ $m(PR)$ ✓ subst gradient and point ✓ answer	(3)

	$m(PR) = \frac{2+1}{-4-2} = -\frac{1}{2}$ $y - y_1 = m(x - x_1)$ $y - 2 = \left(-\frac{1}{2}\right)(x + 4)$ $y - 2 = -\frac{1}{2}x - 2$ $y = -\frac{1}{2}x$	✓ $m(PR)$  ✓ subst gradient and point  ✓ answer	(3)
3.3	$m(PR) = -\frac{1}{2}$ $m(QR) = \frac{6+1}{4-2} = \frac{7}{2}$ $\tan \alpha = -\frac{1}{2}$ $\tan \beta = \frac{7}{2}$ $\alpha = 180^\circ - 26,57^\circ$ $\beta = 74,05^\circ$ $\alpha = 153,43^\circ$ $\therefore \hat{PQR} = 79,38^\circ$	✓ tan ratio $\alpha$ ✓ $\alpha = 153,43^\circ$ ✓ $m(QR)$ ✓ $\beta = 74,05^\circ$  ✓ $\hat{PQR}$	(5)
3.4	$y = -\frac{1}{2}x$ $(-1; t): t = -\frac{1}{2}(-1)$ $t = \frac{1}{2}$  OR  $m(PA) = m(PR)$ $\frac{2-t}{-4+1} = -\frac{1}{2}$ $\frac{2-t}{-3} = -\frac{1}{2}$ $4-2t = 3$ $-2t = -1$ $t = \frac{1}{2}$	✓ equation  ✓ answer	(2)
		✓ equating  ✓ answer	(2)
			[14]


**QUESTION/VRAAG 4**



4.1	P(2 ; -3) Radius = $\sqrt{34}$	✓ centre ✓ radius	(2)
4.2	$m(PB) = \frac{b+3}{a-2}$	✓ subst in formula ✓ answer	(2)
4.3	$m(AB) = -\frac{a-2}{b+3}$  OR  $m(AB) = \frac{b-5}{a-4}$	✓ answer	(1)
4.4	Eq AB: $5y = 3x + 13$ through B( $a$ ; $b$ ) $5b = 3a + 13$ $b = \frac{3}{5}a + \frac{13}{5}$ .....1  Eq PB: $y = mx + c$ $m = -\frac{5}{3}$ and P(2 ; -3) $-3 = -\frac{5}{3}(2) + c$ $c = \frac{1}{3}$ $y = -\frac{5}{3}x + \frac{1}{3}$ through B( $a$ ; $b$ ) $b = -\frac{5}{3}a + \frac{1}{3}$ .....2	✓ subst B( $a$ ; $b$ ) in line equation  ✓ $m(PB) = -\frac{5}{3}$ ✓ subst gradient and P(2 ; -3)  ✓ equation PB	



 <p>Subs 1 in 2: <math>\frac{3}{5}a + \frac{13}{5} = -\frac{5}{3}a + \frac{1}{3}</math></p> $9a + 39 = -25a + 5$ $34a = -34$ $a = -1$ $b = 2$ <p>OR</p> <p>Eq AB: <math>5y = 3x + 13</math> through B(<math>a ; b</math>)</p> $5b = 3a + 13$ $b = \frac{3}{5}a + \frac{13}{5} \quad \dots\dots 1$ $m(PB) \times m(BA) = -1$ $\frac{b+3}{a-2} \times \frac{b-5}{a-4} = -1$ $(b+3)(b-5) = -1(a-2)(a-4) \quad \dots\dots 2$ <p>Subs 1 in 2: <math>\left(\frac{3}{5}a + \frac{13}{5} + 3\right)\left(\frac{3}{5}a + \frac{13}{5} - 5\right) = -1(a-2)(a-4)</math></p> $\left(\frac{3}{5}a + \frac{28}{5}\right)\left(\frac{3}{5}a + \frac{12}{5}\right) = -a^2 + 6a - 8$ $\frac{9}{25}a^2 + \frac{48}{25}a - \frac{336}{25} = -a^2 + 6a - 8$ $34a^2 - 102a - 136 = 0$ $a^2 - 3a - 4 = 0$ $(a-4)(a+1) = 0$ $a = 4 \text{ or/of } a = -1$ <p>NA</p> $b = 2$	<ul style="list-style-type: none"> <li>✓ equate equations</li> <li>✓ simplification</li> </ul> <p>(6)</p>
--	---

	<p>OR</p> <p>Subst B(<math>a</math> ; <math>b</math>) in <math>5y = 3x + 13</math>: <math>5b = 3a + 13</math> .....1</p> $y = \frac{3}{5}x + \frac{13}{5}$ $\therefore \frac{3}{5} = -\frac{a-2}{b+3}$ $3(b+3) = -5(a-2)$ $3b+9 = -5a+10$ $3b = -5a+1$ $b = -\frac{5}{3}a + \frac{1}{3} \quad \dots\dots 2$ <p>Subst 2 in 1: <math>5\left(-\frac{5}{3}a + \frac{1}{3}\right) = 3a + 13</math></p> $-\frac{25}{3}a + \frac{5}{3} = 3a + 13$ $-25a + 5 = 9a + 39$ $-34a = 34$ $a = -1$ $b = 2$	<ul style="list-style-type: none"> <li>✓ subst B(<math>a</math> ; <math>b</math>) in line equation</li> <li>✓ equate gradients</li> <li>✓ simplification</li> <li>✓ equation 2</li> <li>✓ subst</li> <li>✓ simplification</li> </ul>	(6)
4.5	$AB = \sqrt{(-1-4)^2 + (2-5)^2}$ $AB = \sqrt{34}$	<ul style="list-style-type: none"> <li>✓ subst in dist formula</li> <li>✓ answer</li> </ul>	(2)
4.6	$(x-2)^2 + (y+3)^2 = 34 \text{ through } (d ; 0)$ $(d-2)^2 + (0+3)^2 = 34$ $(d-2)^2 + 9 = 34$ $(d-2)^2 = 25$ $d-2 = \pm 5$ $d = 7 \quad \text{or} \quad d = -3$ <p style="text-align: center;">NA</p>	<ul style="list-style-type: none"> <li>✓ Subst in formula</li> <li>✓ simpification</li> <li>✓ factors</li> <li>✓ answers</li> </ul>	(4)

	<p>OR</p> $(x-2)^2 + (y+3)^2 = 34 \text{ through } (d ; 0)$ $(d-2)^2 + (0+3)^2 = 34$ $d^2 - 4d + 4 + 9 = 34$ $d^2 - 4d - 21 = 0$ $(d-7)(d+3) = 0$ $d = 7 \quad \text{or} \quad d = -3$ <p style="text-align: center;">NA</p>	<ul style="list-style-type: none"> <li>✓ subst in formula</li> <li>✓ standard form</li> <li>✓ factors</li> <li>✓ select answer</li> </ul> <p style="text-align: right;">(4)</p>
	<p>OR</p>  $m(PC) \times m(AC) = -1$ $\frac{-3-0}{2-d} \times \frac{5-0}{4-d} = -1$ $\frac{-3}{2-d} \times \frac{5}{4-d} = -1$ $\frac{-15}{8-6d+d^2} = -1$ $-15 = -8 + 6d - d^2$ $d^2 - 6d - 7 = 0$ $(d-7)(d+1) = 0$ $d = 7 \quad \text{or} \quad d = -1$ <p style="text-align: center;">NA</p>	<ul style="list-style-type: none"> <li>✓ subst in formula</li> <li>✓ standard form</li> <li>✓ factors</li> <li>✓ select answer</li> </ul> <p style="text-align: right;">(4)</p>
4.7	$m(PB) = \frac{2+3}{-1-2} = -\frac{5}{3}$ $m(PC) = \frac{-3-0}{2-7} = \frac{3}{5}$ $\therefore m(PB) \times m(PC) = -\frac{5}{3} \times \frac{3}{5} = -1$ $\therefore \hat{BPC} = 90^\circ$	<ul style="list-style-type: none"> <li>✓ <math>m(PB)</math></li> <li>✓ <math>m(PC)</math></li> <li>✓ subst in formula</li> </ul> <p style="text-align: right;">(3)</p>

4.8	<p>y-intercept of AB: Let <math>x = 0</math> in <math>5y = 3x + 13</math></p> $5y = 13$ $y = \frac{13}{5}$ $D\left(0 ; \frac{13}{5}\right)$ <p>Equation of CP: <math>y = mx + c</math></p> $-3 = \frac{3}{5}(2) + c$ $c = -\frac{21}{5}$ $y = \frac{3}{5}x - \frac{21}{5}$ $E\left(0 ; -\frac{21}{5}\right)$ $DE = \frac{34}{5} = 6,8$	<p>✓ <math>y</math> - value</p> <p>✓ subst <math>m</math> and point</p> <p>✓ equation of CP</p> <p>✓ <math>y</math> - value of E</p> <p>✓ answer</p>	<p>(5)</p>
			<p>[25]</p>

QUESTION/VRAAG 5


5.1.1	$\begin{aligned} \sin 215^\circ &= -\sin 35^\circ \\ &= -\frac{m}{\sqrt{m^2 + 1}} \end{aligned}$ 	$\checkmark -\sin 35^\circ$ $\checkmark \sqrt{m^2 + 1}$ $\checkmark$ answer	(3)
5.1.2	$\begin{aligned} \sin 70^\circ &= \sin 2 \times 35^\circ \\ &= 2 \sin 35^\circ \cos 35^\circ \\ &= 2 \left( \frac{m}{\sqrt{m^2 + 1}} \right) \left( \frac{1}{\sqrt{m^2 + 1}} \right) \\ &= \frac{2m}{m^2 + 1} \end{aligned}$	$\checkmark$ double $\angle$ $\checkmark$ subst $\checkmark$ answer	(3)
5.2	 $\begin{aligned} &50 \times \sin(2\alpha + \beta) \\ &= 50 [\sin 2\alpha \cos \beta + \cos 2\alpha \sin \beta] \\ &= 50 [2 \sin \alpha \cos \alpha \cos \beta + (2 \cos^2 \alpha - 1) \sin \beta] \\ &= 50 \left[ 2 \left( \frac{1}{\sqrt{10}} \right) \left( \frac{3}{\sqrt{10}} \right) \left( \frac{7}{\sqrt{50}} \right) + \left( 2 \left( \frac{3}{\sqrt{10}} \right)^2 - 1 \right) \left( \frac{1}{\sqrt{50}} \right) \right] \\ &= 50 \left[ \frac{42}{10\sqrt{50}} + \frac{8}{10\sqrt{50}} \right] \\ &= 50 \left[ \frac{42}{10\sqrt{50}} + \frac{8}{10\sqrt{50}} \right] \\ &= 50 \left[ \frac{50}{10\sqrt{50}} \right] \\ &= 25\sqrt{2} \end{aligned}$	$\checkmark \sqrt{10} \text{ & } \sqrt{50}$ $\checkmark$ compound $\angle$ $\checkmark$ double $\angle$ 's $\checkmark$ subst $\checkmark$ simplification $\checkmark$ answer	(6)

5.3	$\begin{aligned} & \frac{\cos(-x) \tan(180^\circ - x) \cos(90^\circ - x)}{\sin(540^\circ + x) \sin(180 - x)} \\ &= \frac{(\cos x)(-\tan x)(\sin x)}{(-\sin x)(\sin x)} \\ &= \frac{-\tan x}{-\tan x} \\ &= 1 \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>\cos x</math></li> <li>✓ <math>-\tan x</math></li> <li>✓ <math>\sin x</math></li> <li>✓ <math>-\sin x</math></li> <li>✓ <math>\sin x</math></li> </ul> <p>✓ answer</p>	(6)
5.4.1	$\begin{aligned} \frac{\sin x}{1 - \sin x} + \frac{\sin x}{1 + \sin x} &= \frac{2 \tan x}{\cos x} \\ \text{LHS} &= \frac{\sin x}{1 - \sin x} + \frac{\sin x}{1 + \sin x} \\ &= \frac{\sin x(1 + \sin x) + \sin x(1 - \sin x)}{(1 - \sin x)(1 + \sin x)} \\ &= \frac{\sin x + \sin^2 x + \sin x - \sin^2 x}{1 - \sin^2 x} \\ &= \frac{2 \sin x}{\cos^2 x} \\ &= \frac{2 \sin x}{\cos x \cos x} \\ &= \frac{2 \tan x}{\cos x} = \text{RHS} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ LCM</li> <li>✓ simplify numerator</li> <li>✓ <math>1 - \sin^2 x</math></li> <li>✓ <math>\cos^2 x</math></li> </ul>	(4)
5.4.2	$x = 90^\circ$ and $x = 270^\circ$	<ul style="list-style-type: none"> <li>✓ <math>90^\circ</math></li> <li>✓ <math>270^\circ</math></li> </ul>	(2)
			[24]

**QUESTION/VRAAG 6**


6.1	$\cos(x - 30^\circ) = \sin x$ $\cos(x - 30^\circ) = \cos(90^\circ - x)$ $x - 30^\circ = 90^\circ - x + k \cdot 360^\circ \text{ or } x - 30^\circ = 360^\circ - (90^\circ - x) + k \cdot 360^\circ$ $2x = 120^\circ + k \cdot 360^\circ \quad x - 30^\circ = 360^\circ - 90^\circ + x + k \cdot 360^\circ$ $x = 60^\circ + k \cdot 180^\circ, k \in \mathbb{Z} \quad \text{NA}$ $x = 60^\circ$	✓ co-function ✓ both equations ✓ $x = 60^\circ + k \cdot 180^\circ, k \in \mathbb{Z}$ ✓ $x = 60^\circ$	(4)
6.2.1		✓✓ f ✓✓ g	(4)
6.2.2	$y \in [-1 ; 1]$	✓ critical values ✓ notation	(2)
6.2.3	$x \in (30^\circ ; 90^\circ]$	✓ critical values ✓ notation	(2)
6.2.4	$x \in [-90^\circ ; -60^\circ], x = 0^\circ$	✓✓ $x \in [-90^\circ ; -60^\circ]$ ✓ $x = 0^\circ$	(3)
6.2.5	$x \in (60^\circ ; 90^\circ]$	✓ critical values ✓ notation	(2)
			[17]

QUESTION/VRAAG 7



7.1.1	$\hat{A} = \hat{AEB}$ ∠'s opp equal sides $2\hat{A} = 180^\circ - 2x$ ∠'s of $\Delta$ $\hat{A} = 90^\circ - x$	✓ S ✓ R	(2)
7.1.2	$\hat{ACB} = 2x$ tan- chord theorem	✓ S ✓ R	(2)
7.2	In $\Delta ECB$ : $\frac{BC}{\sin(90^\circ + x)} = \frac{h}{\sin 2x}$ $BC = \frac{h \cos x}{2 \sin x \cos x}$ $BC = \frac{h}{2 \sin x}$	✓ subst in sine rule ✓ co-function ✓ double ∠	(3)
7.3	In $\Delta BCD$ : $\hat{C} = 90^\circ$ $\frac{BD}{\sin 90^\circ} = \frac{h}{\sin(90^\circ - x)}$ $BD = \frac{h}{2 \sin x \cos x}$ $BD = \frac{h}{\sin 2x}$	✓ $\hat{C} = 90^\circ$ ✓ subst in sine rule ✓ co-function	(3)  OR $\sin(90^\circ - x) = \frac{h}{BD}$ $BD = \frac{h}{2 \sin x \cos x} = \frac{h}{\sin 2x}$
			[10]


QUESTION/VRAAG 8



8.1	$\hat{T}_2 = \hat{A} = 50^\circ$ tan-chord theorem	$\checkmark S \checkmark R$	(2)
8.2	$\hat{A} = \hat{K}_3 = 50^\circ$ corresp $\angle's$ , $AQ \parallel KP$	$\checkmark S \checkmark R$	(2)
8.3	$\hat{A} = \hat{Q}_3 = 50^\circ$ tan-chord theorem  OR $\hat{T}_2 = \hat{Q}_3 = 50^\circ$ $\angle's$ opp equal tangents	$\checkmark S \checkmark R$	(2)
8.4	Line subtend equal $\angle's$ OR converse $\angle's$ in the same segment	$\checkmark R$	(1)
8.5	$\hat{T}_2 = \hat{K}_2 = 50^\circ$ $\angle's$ in same segment $\hat{K}_2 = \hat{Q}_1 = 50^\circ$ alt $\angle's$ , $AQ \parallel KP$	$\checkmark S \checkmark R$ $\checkmark R$	(3)
			[10]




QUESTION/VRAAG 9




9.1	<p>Construction: Draw radii ON and OL</p> <p><math>\hat{O}_1 = 2 \times \hat{K}</math> midpt <math>\angle = 2 \times</math> circumf <math>\angle</math></p> <p><math>\hat{O}_2 = 2 \times \hat{M}</math> midpt <math>\angle = 2 \times</math> circumf <math>\angle</math></p> <p><math>\hat{O}_1 + \hat{O}_2 = 360^\circ</math> revolution</p> <p><math>2\hat{K} + 2\hat{M} = 360^\circ</math></p> <p><math>\hat{K} + \hat{M} = 180^\circ</math></p>	<p>✓ Construction</p> <p>✓ S ✓ R</p> <p>✓ S/R</p> <p>✓ S/R</p>	(5)
-----	--	--	-----





9.2.1	$\hat{J} = \hat{F}_3$ $\angle's$ opp equal sides $\hat{F}_3 = \hat{K}_2$ $\angle's$ in the same segment $\therefore \hat{J} = \hat{K}_2$	$\checkmark S \checkmark R$ $\checkmark S \checkmark R$ $\checkmark R$	(4)
9.2.2	$\hat{K}_1 = \hat{J}$ ext $\angle$ of cyclic quad $\hat{J} = \hat{K}_2$ proven $\hat{K}_1 = \hat{K}_2$	$\checkmark S \checkmark R$	(2)
9.2.3	$\hat{F}_1 + \hat{F}_2 = \hat{J}$ tan-chord theorem $\hat{J} = \hat{K}_2$ proven $\hat{F}_1 + \hat{F}_2 = \hat{K}_2$ $\therefore KLFV$ is a cyclic quad      converse ext $\angle$ of cyclic quad	$\checkmark S \checkmark R \checkmark$ $\checkmark R$	(3)

QUESTION/VRAAG 10



10.1	$\hat{S}_1 = \hat{A}_1 = x$ $\hat{A}_1 = \hat{R}_1 = x$ <p>alt <math>\angle's</math>, PS <math>\parallel</math> AB          ext <math>\angle</math> of cyclic quad</p>	$\checkmark$ S $\checkmark$ R $\checkmark$ S $\checkmark$ R	(4)
10.2	<p>In <math>\Delta PQT</math> and <math>\Delta PRQ</math>:</p> <p>(i) <math>\hat{P}_1 = \hat{P}_1</math> common          (ii) <math>\hat{Q}_1 = \hat{R}_2</math> tan-chord theorem  <math>\therefore \Delta PQT \equiv \Delta PRQ</math> 3 <math>\angle's</math></p> <p>OR</p> <p>In <math>\Delta PQT</math> and <math>\Delta PRQ</math>:</p> <p>(i) <math>\hat{P}_1 = \hat{P}_1</math> common          (ii) <math>\hat{Q}_1 = \hat{R}_2</math> tan-chord theorem          (iii) <math>\hat{T}_1 = \hat{PQR}</math> 3<sup>rd</sup> <math>\angle</math>  <math>\therefore \Delta PQT \equiv \Delta PRQ</math> 3 <math>\angle's</math></p>	$\checkmark$ S $\checkmark$ S/R $\checkmark$ R	(3)
10.3	$\frac{PQ}{PR} = \frac{PT}{PQ} \quad \parallel \Delta's$ $\frac{PQ}{9} = \frac{4}{PQ}$ $PQ^2 = 36$ $\therefore PQ = 6$	$\checkmark$ R $\checkmark$ subst $\checkmark$ answer	(3)

10.4	<p>In <math>\Delta PTS</math> and <math>\Delta PSR</math> :</p> <p>(i) <math>\hat{P}_2 = \hat{P}_2</math> common</p> <p>(ii) <math>\hat{S}_1 = \hat{R}_1 = x</math> proven</p> <p><math>\therefore \Delta PTS \parallel \Delta PSR</math> 3 <math>\angle's</math></p> <p>OR</p> <p>In <math>\Delta PTS</math> and <math>\Delta PSR</math> :</p> <p>(i) <math>\hat{P}_2 = \hat{P}_2</math> common</p> <p>(ii) <math>\hat{S}_1 = \hat{R}_1 = x</math> proven</p> <p>(iii) <math>\hat{P}_2 = \hat{S}</math> 3<sup>rd</sup> <math>\angle</math></p> <p><math>\therefore \Delta PTS \parallel \Delta PSR</math> 3 <math>\angle's</math></p>	<p>✓ S</p> <p>✓ S</p> <p>✓ R</p> <p>✓ S</p> <p>✓ S/R</p> <p>✓ S</p>	(3)
10.5	<p><math>\frac{PT}{PS} = \frac{PS}{PR} \parallel \Delta's</math></p> <p><math>\therefore PS^2 = PT \cdot PR</math></p> <p><math>PQ^2 = PT \cdot PR</math> from 10.3</p> <p><math>PS^2 = PQ^2</math></p> <p><math>\therefore PS = PQ</math></p>	<p>✓ S/R</p> <p>✓ S</p> <p>✓ S</p>	(3)
			[16]

**TOTAL/TOTAAL : 150**