



education

MPUMALANGA PROVINCE  
REPUBLIC OF SOUTH AFRICA

**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 12**

**MATHEMATICS P2 (FINAL)**

*Stanmorephysics.com*  
**JUNE 2024**

**MARKS: 150**

**TIME: 3 HOURS**



**This question paper consists of 14 pages including 1 information sheet  
and an answer book is provided.**

**INSTRUCTIONS AND INFORMATION**

Read the following instructions carefully before answering the questions.

1. The question paper consists of 10 questions.
2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
3. Clearly show ALL calculations, diagrams, graphs, etc. which you have used in determining the answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical) unless stated otherwise.
6. If necessary, round off answers to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. Write neatly and legibly.



### QUESTION 1

Nine learners wrote a test out of 30 their marks are as follows

7	9	9	13	17	21	24	26	27
---	---	---	----	----	----	----	----	----


- 1.1 Write down the five number summary of the data. (4)
- 1.2 Draw a box and whisker diagram to represent the data. (3)
- 1.3 Comment on the skewness of the data. (1)
- 1.4 If a learner's mark lies below one standard deviation from the mean, the learner must attend an intervention class. How many learners will not attend the intervention class. (2)
- 1.5 The teacher omitted a question with a total of two marks. Determine the actual
  - 1.5.1 Standard deviation (1)
  - 1.5.2 Mean (1)
  - 1.5.3 Upper quartile (1)

**[13]**



### QUESTION 2

The ogive below shows the money collected by parents during a fundraising event at a school.




- 2.1 How many parents were present. (1)
- 2.2 Write down the modal class. (1)
- 2.3 Estimate the number of parents that can contribute R1 900. (2)
- 2.4 Estimate the semi-interquartile range. (3)

[7]



**QUESTION 3**


In the diagram below, PQRS is parallelogram with  $SP \parallel QR$  and vertices  $P(a;5)$ ,  $Q(3;0)$ ,  $R(1;-8)$  and  $S(-1;-3)$ .  $SE \perp QR$ .



- 3.1 Determine the length of QR. (2)
  - 3.2 Determine the gradient of QR. (2)
  - 3.3 Determine the equation of SE (3)
  - 3.4 Determine the equation of a circle passing through SER (3)
  - 3.5 Calculate the value of  $a$ . (3)
  - 3.6 Calculate the value of:
    - 3.6.1  $\theta$  (2)
    - 3.6.2  $\widehat{SRQ}$  (5)
  - 3.7 Determine the area of PSRQ (3)
- [23]**

**QUESTION 4**

A circle with equation  $x^2 + 2x + y^2 + 2y = 4$  has centre C. FAE is a tangent to the circle at  $A(-3;1)$  with  $\hat{EAB} = 33^\circ$



- 4.1 Determine the coordinates of C. (2)
- 4.2 Determine the equation of the tangent to the circle at A. (5)
- 4.3 Determine the value of  $k$  if  $y = x + k$  will be a secant to the circle. (3)
- 4.4 Calculate the size of  $\hat{ACB}$ . (4)
- 4.5 Another circle with equation  $(x + 3)^2 + (y + 2)^2 = 4$  is drawn, will the two circles intersect (4)

**[18]**

**QUESTION 5**

5.1 Given  $\cos 11^\circ = k$ . Determine the following in terms of  $k$ , **without using a calculator**.

5.1.1  $\sin 11^\circ$  (2)

5.1.2  $\sin 22^\circ$  (2)

5.1.3  $\cos 19^\circ$  (3)

5.1.4  $\cos 5,5^\circ$  (3)

5.2 Simplify the following expression without a calculator:

$$\frac{\sin 25^\circ \cos(x - 180^\circ) \sin 2x}{\sin^2(90 + x) \cos 65^\circ}$$
 (5)

5.3 Given:  $\frac{\sin x - \cos x}{\sin x + \cos x} - \frac{\sin x + \cos x}{\cos x - \sin x} = 2 \tan 2x$

5.3.1 Prove the identity. (5)

5.3.2 For which values of  $x$  will  $\frac{\sin x - \cos x}{\sin x + \cos x} - \frac{\sin x + \cos x}{\cos x - \sin x}$  be undefined? (4)

**[24]**



**QUESTION 6**

Consider the functions:  $f(x) = \cos 2x$  and  $g(x) = \sin(x - 45^\circ)$ , where  $x \in [-90^\circ; 180^\circ]$

6.1 On the same set of axis sketch the graphs of  $f$  and  $g$  (5)

6.2 Determine the general solution of  $f(x) = g(x)$ . (4)

6.3 Determine the  $x$ -value(s) in the interval  $x \in [0^\circ; 180^\circ]$  for which:

6.3.1  $f(x) > g(x)$  (2)

6.3.2  $x f(x) \cdot g(x) > 0$  (2)



**[13]**





**QUESTION 7**

AD represents a vertical tower, with A at the top and D at the foot of the tower. D is a point on the tower below A. On the same horizontal level as C is point B,  $AB = k$  metres.  $\hat{CDB} = 2\theta$  and  $\hat{ABD} = \theta$ .



7.1 Show that  $CB = k \cdot \sin 2\theta$  (7)


7.2 If  $k = 10$  units and  $\theta = 30^\circ$  calculate the area of  $\triangle CBD$  (4)  
**[11]**



Give reasons for your statements in QUESTIONS 8, 9 and 10.

**QUESTION 8**



8.1 In the diagram below O is the centre of the circle



Prove that  $\widehat{BOC} = 2\widehat{BAC}$

(5)

8.2 In the diagram below O is a centre of the circle with  $FE = AB$  and  $BO \parallel CD$ .  $\hat{O}_2 = 88^\circ$  and  $\hat{D}_2 = 23^\circ$ . Tangent PAS touches the circle at A.



Determine with reasons the size of the following angles:

8.2.1  $\hat{O}_3$  (2)

8.2.2  $\hat{D}_1$  (3)

8.2.3  $\hat{A}_1$  (2)

8.2.4  $\hat{C}_1$  (2)



8.2.5  $\hat{C}_2$  (2)

**[16]**



**QUESTION 9**

In the diagram below TWSK lies on the circumference of the circle, such that  $TW \parallel SK$  with O the centre on the circle.  $\angle K_1 = x$




- 9.1 Determine with reasons, THREE other angles equal to  $x$ . (4)
- 9.2 Prove that SOFT is a cyclic quadrilateral (4)
- 9.3 Prove that  $ST = 2SO \cdot \sin x$  (4)

**[12]**



**QUESTION 10**

In the diagram below BCED is a cyclic quadrilateral with  $AB = AD$ ,  $AB = 2$  cm,  $BC = 3$  cm and  $BD = 4$  cm.



- 10.1 Prove that  $BD \parallel CE$  (3)
- 10.2 Calculate the length of DE (2)
- 10.3 Prove that  $\triangle ADB \sim \triangle AEC$  (3)
- 10.4 Calculate the length of EC (2)
- 10.5 Calculate the value of  $\frac{\text{Area of } \triangle ECD}{\text{Area of } \triangle ABD}$  (3)

[13]

**TOTAL: 150**



**FORMULA SHEET**

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d]$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; -1 < r < 1$$

$$F = \frac{x[(1 + i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1 + i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}; \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c \quad y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cdot \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2}ab \cdot \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cdot \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cdot \cos \beta + \sin \alpha \sin \beta$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$= 1 - 2\sin^2 \alpha$$

$$= 2\cos^2 \alpha - 1$$

$$\bar{x} = \frac{\sum fx}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ of } B) = P(A) + P(B) - P(A \text{ en } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum(x - \bar{x})(y - \bar{y})}{\sum(x - \bar{x})^2}$$



**education**

**MPUMALANGA PROVINCE  
REPUBLIC OF SOUTH AFRICA**

**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 12**

**MATHEMATICS PAPER 2**

**JUNE 2024**

**MARKING GUIDELINE**

*Stanmorephysics.com*

**MARKS: 150**

**This marking guideline consist of 18 pages**

NSC Marking Guideline

**NOTE:**

- If a candidate answered a question TWICE, mark only the FIRST attempt.
- If a candidate crossed out an answer and did not redo it, mark the crossed-out answer.
- Consistent accuracy applies to ALL aspects of the marking guidelines.
- Assuming values/answers in order to solve a problem is unacceptable.

**QUESTION 1**

7	9	9	13	17	21	24	26	27
---	---	---	----	----	----	----	----	----

1.1	$Minimum = 7$ $Q_1 = 9$ $Q_2 = 17$ $Q_3 = 25$ $Maximum = 27$	✓ 7 ✓ 9 ✓ 17 ✓ 25 ✓ 27  (5)
1.2		✓ Whiskers ✓ $Q_1, Q_3$ ✓ $Q_2$  (3)
1.3	Equally distributed/ Centrally skewed	✓ Answer.  (1)
1.4	$\bar{x} = 17$ $\sigma_x = 7,38$ $\bar{x} - \sigma_x = 17 - 7,38$ $= 9,62$ Only 8 learners	✓ $\bar{x} - \sigma_x$ ✓ Answer  (2)
1.5.1	$\sigma_x = 7,38$	✓ Answer.  (1)
1.5.2	$\bar{x} = 19$	✓ Answer.  (1)
1.5.3	$Q_3 = 27$	✓ Answer.  (1)
		<b>[13]</b>




**QUESTION 2**

2.1	60 parents	✓ Answer. (1)
2.2	$1000 < x < 1500$	✓ Answer. (1)
2.3	46	✓ ✓ Answer. (2)
2.4	$\begin{aligned} \text{IQR} &= 1850 - 1000 \\ &= 850 \\ \therefore \text{Semi-IQR} &= \frac{850}{2} \\ &= 425 \end{aligned}$	✓ IQR ✓ $\frac{850}{2}$ ✓ Answer. (3)
		<b>[7]</b>






**QUESTION 3**




3.1	$QR = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ $QR = \sqrt{((3-1)^2 + (0+8)^2)}$ $QR = \sqrt{68}$ $QR = 2\sqrt{17}$ $QR = 8,25$	✓ correct subst into the distance formula.  ✓ $2\sqrt{17}$  (2)
3.2	$m_{QR} = \frac{0 - (-8)}{3 - 1}$ $m_{QR} = \frac{8}{2}$ $m_{QR} = 4$	✓ Subst into the formula.  ✓ Answer. (2)
3.3	$m_{SE} = -\frac{1}{4}$ $-3 = -\frac{1}{4}(-1) + c$ $c = -\frac{13}{4}$ $y = -\frac{1}{4}x - \frac{13}{4}$	✓ $m_{SE}$  ✓ substitution $S(-1; -3)$  ✓ Answer. (3)

NSC Marking Guideline

<p>3.4</p>	 $M_{SE} \left( \frac{-1+1}{2}; \frac{-3-8}{2} \right)$ $= M_{SE} \left( 0; -\frac{11}{2} \right)$ $(x-a)^2 + (y-b)^2 = r^2$ $(x-0)^2 + \left( y + \frac{11}{2} \right)^2 = (\sqrt{17})^2$ $x^2 + \left( y + \frac{11}{2} \right)^2 = 17$	<p>✓ Midpoint                  ✓ Substitution                  ✓ Answer.</p> <p>(3)</p>
<p>3.5</p>	$m_{RQ} = m_{SP} = 4$ $\frac{4}{1} = \frac{5+3}{a+1}$ $8 = 4a + 4$ $4 = 4a$ $a = 1$ <p style="text-align: center;">OR</p>  $-3 = 4(-1) + c$ $c = 1$ $y = 4x + 1$ $5 = 4a + 1$ <p style="text-align: center;">OR</p>	<p>✓ Gradient                  ✓ Substitution                  ✓ Answer</p> <p>(3)</p> <p>OR</p> <p>✓ Gradient                  ✓ Substitution                  ✓ Answer</p> <p>(3)</p>
<p>3.6.1</p>	$m_{PS} = 4$ $\tan \theta = 4$ $\theta = 75,96^\circ$	<p>✓ <math>\tan \theta = 4</math>                  ✓ <math>75,96^\circ</math></p> <p>(2)</p>
<p>3.6.2</p>	$m_{SR} = \frac{-8 - (-3)}{1 - (-1)}$ $= -\frac{5}{2}$ $\tan \beta = -\frac{5}{2}$ $= 180^\circ - 68,20^\circ$ $= 111,80^\circ$	 <p>✓ <math>-\frac{5}{2}</math>                  ✓ <math>\tan \beta = -\frac{5}{2}</math></p> <p>✓ <math>111,80^\circ</math></p>



**QUESTION 4**




4.1	$C(-1; -1)$	✓ $x$ - value. ✓ $y$ - value. (2)
4.2	$m_{\text{tangent}} = \frac{1+1}{-3+1} = -1$ $1 = 1(-3) + c$ $c = 4$ $y + x + 4$	✓ Substitution ✓ $m_{\text{tangent}}$ ✓ Substitution $A(-3; 1)$ ✓ Answer (4)
4.3	$D(1; -3)$ $-3 = 1(1) + c$ $c = -4$ $y = x - 4$ $-4 < k < 4$	✓ Substitution $D(1; -3)$ ✓ Equation ✓ Notation ✓ Critical Values (4)
4.4	$\hat{D} = 33^\circ$ [Tan-Chord Theorem] $\hat{ACB} = 66^\circ$ [ $\angle$ at centre = $2 \times \angle$ at circumference]	✓ S    ✓ R ✓ S    ✓ R (4)

NSC Marking Guideline

<p>4.5</p>	<p><math>C(-3; -2) \quad r = 2</math></p> $d_{CD} = \sqrt{(-1+3)^2 + (-1+2)^2}$ $= \sqrt{5}$ $= 2,24$ $r_{SM} = 2 + \sqrt{6}$ $= 4,45$ <p><math>\therefore d_{CD} &lt; r_{SM}</math>, the circles intersect twice.</p>	<p>✓ Substitution                  ✓ <math>d_{CD}</math>                  ✓ <math>r_{SM}</math>                  ✓ Answer</p> <p>(4)</p>
		<p>[18]</p>

**QUESTION 5**

<p>5.1.1</p>	 <p><math>(k)^2 + y^2 = (1)^2</math>  <math>k^2 + y^2 = 1</math>  <math>y = \pm\sqrt{1-k^2}</math>  <math>\therefore y = \sqrt{1-k^2}</math></p> <p><math>\sin 11^\circ = \sqrt{1-k^2}</math></p>	<p>✓ <math>\sqrt{1-k^2}</math>                  ✓ Answer</p> <p>(2)</p>
<p>5.1.2</p>	<p><math>\sin 2(11^\circ) = 2 \sin(11^\circ) \cos(11^\circ)</math>  <math>= 2(\sqrt{1-k^2})(k)</math>  <math>= 2k\sqrt{1-k^2}</math></p>	<p>✓ Expansion                  ✓ <math>2k\sqrt{1-k^2}</math></p> <p>(2)</p>

NSC Marking Guideline

5.1.3	$\begin{aligned} \cos 19^\circ &= \cos(30^\circ - 11^\circ) \\ &= \cos 30^\circ \cos 11^\circ + \sin 30^\circ \sin 11^\circ \\ &= \left(\frac{\sqrt{3}}{2}\right)\left(\frac{k}{1}\right) + \left(\frac{1}{2}\right)\left(\frac{\sqrt{1-k^2}}{1}\right) \\ &= \frac{\sqrt{3}k}{2} + \frac{\sqrt{1-k^2}}{2} \\ &= \frac{\sqrt{3}k + \sqrt{1-k^2}}{2} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>-\cos 60^\circ</math></li> <li>✓ <math>\tan 45^\circ</math></li> <li>✓ <math>\left(-\frac{1}{2}\right)(1)</math></li> <li>✓ <math>-4</math></li> </ul> <p style="text-align: right;">(4)</p>
5.1.4	$\begin{aligned} \cos 11^\circ &= \cos 2(5,5^\circ) = k \\ 2 \cos^2 5,5^\circ - 1 &= k \\ \frac{2 \cos^2 5,5^\circ}{2} &= \frac{k+1}{2} \\ \sqrt{\cos^2 5,5^\circ} &= \sqrt{\frac{k+1}{2}} \\ \cos 5,5^\circ &= \sqrt{\frac{k+1}{2}} \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>2 \cos^2(5,5^\circ) - 1</math></li> <li>✓ Simplification</li> <li>✓ Answer</li> </ul> <p style="text-align: right;">(3)</p>
5.2	$\begin{aligned} &\frac{\sin 25^\circ (-\cos x) 2 \sin x \cos x}{\cos^2 x \cdot \sin 25^\circ} \\ &= \frac{-2 \cos^2 x \sin x}{\cos^2 x} \\ &= -2 \sin x \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>-\cos x</math></li> <li>✓ <math>2 \sin x \cos x</math></li> <li>✓ <math>\cos^2 x</math></li> <li>✓ <math>\sin 25^\circ</math></li> <li>✓ Answer.</li> </ul> <p style="text-align: right;">(3)</p>
5.3.1	$\begin{aligned} &\frac{\sin x - \cos x}{\sin x + \cos x} - \frac{\sin x + \cos x}{\sin x - \cos x} \\ &= \frac{(\sin x - \cos x)^2 - (\sin x + \cos x)^2}{(\sin x + \cos x)(\sin x - \cos x)} \\ &= \frac{\sin^2 x - 2 \sin x \cos x + \cos^2 x - (\sin^2 x + 2 \sin x \cos x + \cos^2 x)}{\sin^2 x - \cos^2 x} \\ &= \frac{-4 \sin x \cos x}{-(\cos^2 x - \sin^2 x)} \\ &= \frac{2 \sin 2x}{\cos 2x} \\ &= 2 \tan 2x \end{aligned}$	<ul style="list-style-type: none"> <li>✓ <math>(\sin x - \cos x)^2</math></li> <li>✓ <math>(\sin x + \cos x)^2</math></li> <li>✓ Simplification</li> <li>✓ <math>2 \sin 2x</math></li> <li>✓ <math>\cos 2x</math></li> </ul> <p style="text-align: right;">(5)</p>

NSC Marking Guideline

<p>5.3.2</p>	$\cos x + \sin x = 0$ $\cos x = -\sin x$ $\tan x = -1$ $x = 135^\circ + k180^\circ, k \in \mathbb{Z}$ $\cos x - \sin x = 0$ $\cos x = \sin x$ $\tan x = 1$ $x = 45^\circ + k180^\circ, k \in \mathbb{Z}$	<p>✓ <math>\cos x + \sin x = 0</math>                  ✓ <math>x = 135^\circ</math></p> <p>✓ <math>\cos x - \sin x = 0</math>                  ✓ <math>x = 45^\circ</math></p> <p>(4)</p>
		<p>[24]</p>


QUESTION 6

<p>6.1</p>		<p><math>f(x)</math></p> <p>✓ <math>x</math> – intercepts                  ✓ <math>y</math> – intercept                  ✓ Shape.</p> <p><math>g(x)</math></p> <p>✓ <math>x</math> and <math>y</math> – intercept                  ✓ Shape.</p> <p>(5)</p>
<p>6.2</p>	$\sin(x - 45^\circ) = \cos 2x$ $\sin(x - 45^\circ) = \sin(90^\circ - 2x)$ $x - 45^\circ = 90^\circ - 2x$ $x = 45^\circ - 120^\circ k, k \in \mathbb{Z}$ $x - 45^\circ = 180^\circ - (90^\circ - 2x)$ $x - 45^\circ = 90^\circ + 2x$ $-x = 135^\circ + k360^\circ, k \in \mathbb{Z}$ $x = -135^\circ - k360^\circ, k \in \mathbb{Z}$	<p>✓ <math>\sin(90^\circ - 2x)</math>                  ✓ <math>x = 45^\circ</math>                  ✓ <math>-120^\circ k, k \in \mathbb{Z}</math></p> <p>✓ <math>x = -135^\circ</math></p> <p>(4)</p>
<p>6.3.1</p>	$0^\circ < x < 45^\circ$	<p>✓ <math>0^\circ &lt; x &lt; 45^\circ</math></p>



	$165^\circ < x < 180^\circ$	✓ $165^\circ < x < 180^\circ$ (2)
6.3.2	$0^\circ < x < 45^\circ$	✓ Notation ✓ Endpoints. (2)
		<b>[13]</b>

**QUESTION 7**




7.1	$\frac{DB}{k} = \cos\theta$ $BD = k\cos\theta$ <p>In <math>\triangle CDB</math></p> $CD = DB = k\cos\theta$ $CB^2 = CD^2 + DB^2 - 2CD \cdot DB \cos\hat{D}$ $CB^2 = (k\cos\theta)^2 + (k\cos\theta)^2 - 2(k\cos\theta)(k\cos\theta)\cos 2\theta$ $CB^2 = k^2\cos^2\theta + k^2\cos^2\theta - 2k^2\cos^2\theta \cdot \cos 2\theta$ $CB^2 = 2k^2\cos^2\theta(1 - \cos 2\theta)$ $CB^2 = 2k^2\cos^2\theta(1 - (1 - 2\sin^2\theta))$ $= 2k^2\cos^2\theta \cdot 2\sin^2\theta$ $CD = \sqrt{4k^2\cos^2\theta \cdot 2\sin^2\theta}$ $= 2k\cos\theta \cdot \sin\theta$ $CD = k\sin 2\theta$	✓ Trig ratio ✓ BD  ✓ Correct substitution ✓ Simplification  ✓ $1 - \cos 2\theta$ ✓ $1 - \sin^2\theta$  ✓ $2k\cos\theta \cdot \sin\theta$ (7)
-----	---	--

NSC Marking Guideline

7.2	$CD = (10) \cos 2(30^\circ)$ $= 5$ $\text{Area} = \frac{1}{2} CD \cdot BD \sin \hat{D}$ $= \frac{1}{2} (5)(5) \sin 60^\circ$ $= \frac{25\sqrt{3}}{4}$ $= 10,83$	✓ Substitution ✓ CD  ✓ Substitution ✓ Answer  (4)
		[11]

QUESTION 8

8.1	 <p>Construction: Draw OA produced to E</p> $\hat{A}_1 + \hat{B} = \hat{O}_1 \quad [\text{Ext } \angle \text{ of } \Delta]$ $OA = OB \quad [\text{Radii}]$ $\hat{A}_1 = \hat{B} \quad [\angle \text{ opp} = \text{sides.}]$ $\therefore \hat{O}_1 = 2\hat{A}_1$ <p>Similarly <math>\hat{O}_2 = 2\hat{A}_2</math></p> $\therefore \hat{O}_1 + \hat{O}_2 = 2\hat{A}_1 + 2\hat{A}_2$ $\therefore \hat{O}_1 + \hat{O}_2 = 2(\hat{A}_1 + \hat{A}_2)$	✓ Construction. ✓ S/R ✓ S ✓ R  ✓ S/R  (5)
8.2.1.	$\hat{O}_3 = 46$ $[\angle \text{ at centre} = 2 \times \angle \text{ at circumference}]$	✓ S ✓ R (2)

NSC Marking Guideline

8.2.2.	$\widehat{D}_1 = \widehat{B}_1$ $\widehat{B}_1 = 23^\circ$	$[\angle opp = sides]$ $[sum\ of\ \angle\ 's\ in\ \Delta]$	$\checkmark\ S\ \checkmark\ R$ $\checkmark\ S/R$	(3)
8.2.3.	$\widehat{A}_1 = 23^\circ$	$[tan\ chord\ theorem]$	$\checkmark\ S\ \checkmark\ R$	(2)
8.2.4.	$\widehat{C}_1 = 23^\circ$	$[= chords = \angle\ 's]$	$\checkmark\ S\ \checkmark\ R$	(2)
8.2.5.	$\widehat{OCD} = 46$ $\widehat{C}_2 = 46^\circ - \widehat{C}_1$ $\widehat{C}_2 = 23^\circ$	$[Alt\ \angle\ 's,\ BO\ //\ CD]$	$\checkmark\ S/R$  $\checkmark\ S$	  (2)
				<b>[16]</b>


QUESTION 9

9.1.	$\widehat{T}_3 = \widehat{K}_1 = x$ $\widehat{W}_1 = \widehat{K}_1 = x$ $\widehat{WSK} = \widehat{W}_1 = x$	$[Alt\ \angle\ 's\ TW\ \parallel\ SK]$ $[\angle\ 's\ in\ the\ same\ seg]$ $[Alt\ \angle\ 's\ TW\ \parallel\ SK]$	$\checkmark\ S\ /R$ $\checkmark\ S\ \checkmark\ R$ $\checkmark\ S\ /R$	(4)
9.2.	$\widehat{O}_1 = 2x$ $\widehat{F}_4 = 2x$ $SOFT\ is\ a\ cyclic\ quad$	$[\angle\ at\ centre = 2 \times \angle\ at\ circumference]$ $[Ext\ \angle\ 's\ of\ a\ \Delta]$ $[converse\ \angle\ 's\ in\ the\ same\ seg]$	$\checkmark\ S\ \checkmark\ R$ $\checkmark\ S\ /R$ $\checkmark\ R$	(4)

NSC Marking Guideline

<p>9.3.</p>	$SO = TO \quad \text{[Radii]}$ $\hat{O}_1 = 2x \quad \text{[Proven]}$ $ST = \sqrt{2(SO)^2 - 2(SO)^2 \cos^2 2x}$ $ST = \sqrt{2(SO)^2(1 - \cos^2 2x)}$ $ST = \sqrt{2(SO)^2[1 - (1 - 2 \sin^2 x)]}$ $ST = \sqrt{2(SO)^2(2 \sin^2 x)}$ $ST = \sqrt{4(SO)^2 \sin^2 x}$ $ST = 2SO \sin x$ <p style="text-align: center;"><b>OR</b></p> <p><i>in</i> <math>\triangle STO</math>:</p> $SO = TO \quad \text{[Radii]}$ $\hat{T}_1 = \hat{S}O = 90^\circ - x \quad \text{[sum of } \angle\text{'s in } \triangle]$ $\frac{ST}{\sin 2x} = \frac{SO}{\sin(90^\circ - x)}$ $ST \cos x = 2SO \sin x \cos x$ $\frac{ST \cos x}{\cos x} = \frac{2SO \sin x \cos x}{\cos x}$ $ST = 2SO \sin x$	<p>✓ S/R</p> <p>✓ Sub in correct formula</p> <p>✓ Double angle</p> <p>✓ Square root</p> <p style="text-align: right;">(4)</p> <p>✓ S/R</p> <p>✓ Sub in correct formula</p> <p>✓ Double angle</p> <p>✓ Dividing by <math>\cos x</math></p> <p style="text-align: right;">(4)</p> <p style="text-align: right;"><b>[12]</b></p>
-------------	--	---

**QUESTION 10**

		
<p>10.1.</p>	$\hat{B}_1 = \hat{D}_1 \quad \text{[}\angle\text{'s opp = sides]}$ $\hat{B}_1 = \hat{E} \quad \text{[Ext } \angle\text{'s of cyclic quad]}$ $\hat{D}_1 = \hat{E} \quad \text{[Each = } \hat{B}_1 \text{]}$ $BD // CE \quad \text{[corresp } \angle\text{'s =]}$	<p>✓ S/R</p> <p>✓ S/R</p> <p>✓ R</p> <p style="text-align: right;">(3)</p>
<p>10.2.</p>	$\frac{2}{DE} = \frac{2}{3}$ $\frac{2DE}{2} = \frac{6}{2}$ $DE = 3 \text{ cm}$ <p style="text-align: center;">[prop theorem, <math>BD // CE</math>]</p>	<p>✓ S/R</p> <p>✓ Answer</p> <p style="text-align: right;">(2)</p>

NSC Marking Guideline

10.3.	<p><i>in</i> <math>\triangle ADB</math> &amp; <math>\triangle AEC</math></p> <p><math>\hat{A} = \hat{A}</math> [common <math>\sphericalangle</math>]</p> <p><math>\hat{D}_1 = \hat{E}</math> [corresp <math>\sphericalangle</math>'s, <math>BD \parallel CE</math>]</p> <p><math>\hat{B}_1 = \hat{C}</math> [corresp <math>\sphericalangle</math>'s, <math>BD \parallel CE</math>]</p> <p><math>\triangle ADB \parallel \triangle AEC</math> [A,A,A]</p>	<p>✓ S/R</p> <p>✓ S/R</p> <p>✓ S/R</p> <p>(3)</p>
10.4.	<p><math>\frac{AD}{AE} = \frac{BD}{EC}</math> [///'s]</p> <p><math>\frac{2}{5} = \frac{4}{EC}</math></p> <p><math>\frac{2EC}{2} = \frac{20}{2}</math></p> <p><math>EC = 10 \text{ cm}</math></p>	<p>✓ S/R</p> <p>✓ Answer (2)</p>
10.5.	<p><math>\frac{\text{Area } \triangle ECD}{\text{Area } \triangle ABD} = \frac{\text{Area } \triangle ECD}{\text{Area } \triangle ACE} \times \frac{\text{Area } \triangle ACE}{\text{Area } \triangle ABD}</math></p> <p><math>= \frac{\frac{1}{2}EC \cdot ED \sin \hat{E}}{\frac{1}{2}EC \cdot AE \sin \hat{E}} \times \frac{\frac{1}{2}AB \cdot AD \sin \hat{A}}{\frac{1}{2}AC \cdot AE \sin \hat{A}}</math></p> <p><math>= \frac{3}{5} \times \frac{(2)(2)}{(5)(5)}</math></p> <p><math>= \frac{12}{125}</math></p>	<p>✓ Formula</p> <p>✓ Substitution</p> <p>✓ Answer (3)</p>
		<b>[13]</b>

<b>TOTAL:</b>	<b>150</b>
---------------	------------

