



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS

PHYSICAL SCIENCES: CHEMISTRY (P2)

MAY/JUNE 2024

Stanmorephysics.com

MARKS: 150

TIME: 3 hours

This question paper consists of 16 pages and 4 data sheets.

INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. Show ALL formulae and substitutions in ALL calculations.
8. Round off your FINAL numerical answers to a minimum of TWO decimal places.
9. Give brief motivations, discussions, etc. where required.
10. You are advised to use the attached DATA SHEETS.
11. Write neatly and legibly.

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, e.g. 1.11 E.

1.1 The functional group for an ALDEHYDE is a ...

- A formyl group.
- B carboxyl group.
- C carbonyl group.
- D hydroxyl group. (2)


1.2 Which ONE of the following equations represents the reaction for the IDENTIFICATION of an UNSATURATED organic compound in the laboratory?

- A $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHCH}_2 + \text{H}_2 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$
- B $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHCH}_2 + \text{HBr} \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CHBrCH}_3$
- C $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHCH}_2 + \text{Br}_2 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CHBrCH}_2\text{Br}$
- D $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3 + \text{Br}_2 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br} + \text{HBr}$ (2)

1.3 Which ONE of the following is the EMPIRICAL formula of ethyl ethanoate?


- A $\text{C}_2\text{H}_4\text{O}$
- B $\text{C}_2\text{H}_2\text{O}$
- C $\text{C}_4\text{H}_8\text{O}$
- D $\text{C}_4\text{H}_8\text{O}_2$ (2)

1.4 In an experiment 5 g of calcium carbonate, $\text{CaCO}_3(\text{s})$, reacts with EXCESS hydrochloric acid, $\text{HCl}(\text{aq})$, at a temperature of $40\text{ }^\circ\text{C}$.



The volume of $\text{CO}_2(\text{g})$ produced versus time is shown by CURVE P in the graph below.


The experiment is repeated with 10 g of the same $\text{CaCO}_3(\text{s})$ sample and an excess of $\text{HCl}(\text{aq})$ with the same concentration at $40\text{ }^\circ\text{C}$. Which ONE of the curves will now be obtained?



- A Curve I
- B Curve II
- C Curve III
- D Curve IV

(2) ...

1.5 The balanced equation below represents a reaction at equilibrium.




Which statement(s) is/are TRUE when a few drops of concentrated hydrochloric acid, HCl(conc), are added to the mixture?

- (i) The reverse reaction will be favoured.
- (ii) The concentration of $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ increases.
- (iii) The colour of the solution changes from yellow to orange.

- A (i) only
- B (i) and (ii) only
- C (i) and (iii) only
- D (ii) and (iii) only (2)

1.6 Consider the equation below for a hypothetical reaction.



If the equilibrium constant $K_c = 1 \times 10^{-4}$, then ...

- A $[\text{A}][\text{B}] < [\text{C}][\text{D}]$
- B $[\text{A}][\text{B}] > [\text{C}][\text{D}]$
- C $[\text{A}] > [\text{C}][\text{D}]$
- D $[\text{A}] < [\text{C}][\text{D}]$ (2)

1.7 Which ONE of the following shows the PRODUCTS for the reaction of oxalic acid with sodium hydroxide?

- A $(\text{COO})_2\text{Na}_2(\text{aq}) + \text{H}_2\text{O}(\ell) + \text{CO}_2(\text{g})$
- B $(\text{COO})_2\text{Na}_2(\text{aq}) + \text{H}_2\text{O}(\ell)$
- C $\text{CH}_3\text{COONa}(\text{aq}) + \text{H}_2\text{O}(\ell)$
- D $\text{CH}_3\text{COONa}(\text{aq}) + \text{H}_2\text{O}(\ell) + \text{CO}_2(\text{g})$ (2)

1.8 Four solutions of different acids of the same concentration are compared.

Which ONE of the following K_a values represents the WEAKEST acid at 25 °C?

A $4,5 \times 10^{-6}$

B $2,5 \times 10^{-5}$


C $1,8 \times 10^{-2}$

D $6,5 \times 10^{-2}$

(2)

1.9 Copper is purified by electrolysis.





Which ONE of the following combinations is CORRECT for the changes occurring at the anode, cathode and in the electrolyte when the cell is in operation?



	MASS OF THE ANODE	MASS OF THE CATHODE	COLOUR OF THE ELECTROLYTE
A	Increases	Decreases	No change
B	Decreases	Increases	No change
C	Increases	Decreases	Becomes darker
D	Decreases	Increases	Becomes lighter

(2)

1.10 Which ONE of the following redox reactions is SPONTANEOUS under standard conditions?



(2)
 [20]

QUESTION 2 (Start on a new page.)

The letters **A** to **H** in the table below represent eight organic compounds.

A	Butan-2-ol	B	$\text{CH}_3\text{C}(\text{CH}_3)_2(\text{CH}_2)_2\text{CH}_3$
C	3-ethylpent-1-yne	D	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$
E	$\begin{array}{c} \text{CH}_3 - \text{CH}_2 - \text{CH} \\ \quad \quad \quad \parallel \\ \text{CH}_3 - \text{CH}_2 - \text{C} - \text{CH}_2 - \text{CH}_3 \end{array}$	F	Butan-1-ol
G	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CH}_2 - \text{CH} - \text{CH}_3 \\ \quad \quad \\ \text{Cl} \quad \quad \text{Cl} \end{array}$	H	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{CH}_3 \\ \\ \text{C} = \text{O} \\ \\ \text{H} \end{array}$

- 2.1 Define the term *hydrocarbon*. (2)
- 2.2 Write down the letter(s) for:
- 2.2.1 TWO compounds that are UNSATURATED hydrocarbons (1)
- 2.2.2 TWO compounds that are CHAIN ISOMERS of each other (2)
- 2.2.3 A secondary alcohol (1)
- 2.3 Write down the:
- 2.3.1 STRUCTURAL formula of the FUNCTIONAL ISOMER of compound **D** (2)
- 2.3.2 General formula of the homologous series to which compound **B** belongs (1)
- 2.3.3 STRUCTURAL formula of compound **C** (2)
- 2.4 Write down the IUPAC name of compound:
- 2.4.1 **E** (3)
- 2.4.2 **G** (3)
- 2.4.3 **H** (2)
- 2.5 Compound **B** undergoes complete combustion. Using MOLECULAR FORMULAE, write down the balanced equation for this reaction. (3)

[22]

QUESTION 3 (Start on a new page.)

The boiling points of some organic compounds are shown in the table below. The atmospheric pressure is 101,3 kPa.

	ORGANIC COMPOUND	BOILING POINT (°C)
A	CH ₃ CH ₂ CH ₂ CH ₂ Cl	78
B	CH ₃ CH(CH ₃)CH ₂ Cl	46
C	CH ₃ CH ₂ CH ₂ CH ₂ OH	118
D	CH ₃ CH ₂ CH ₂ CHO	X


- 3.1 Define the term *boiling point*. (2)
- 3.2 Which ONE of compounds **A**, **B** or **C** is mainly in the liquid phase at 100 °C? (1)
- 3.3 Explain the difference in the boiling points of compounds **A** and **B**. (3)
- 3.4 Consider the boiling points below.

75 °C	120 °C	126 °C
-------	--------	--------

- 3.4.1 Which ONE of these values represents **X**, the boiling point of compound **D**? (1)
- 3.4.2 Fully explain the answer to QUESTION 3.4.1. (2)
- 3.5 The atmospheric pressure is now changed to 83 kPa.
- How will the boiling points of these organic compounds be affected? Choose from INCREASE, DECREASE or REMAIN THE SAME. (1)
- [10]**

QUESTION 4 (Start on a new page.)

- 4.1 In an experiment, a test tube containing methanol, propanoic acid and a catalyst is heated in a water bath.




Write down:

- 4.1.1 The NAME or FORMULA of the catalyst (1)
- 4.1.2 The type of reaction taking place (1)
- 4.1.3 TWO reasons why the use of a water bath is preferred in this experiment (2)
- 4.1.4 The balanced equation for this reaction using STRUCTURAL FORMULAE (5)
- 4.1.5 The IUPAC name of the organic product for this reaction (2)

4.2 Compound **A**, a six-carbon branched haloalkane, is used in a two-step reaction to prepare compound **C**.

Reaction 2 is an ADDITION reaction.




Write down:

- 4.2.1 The NAME or FORMULA of the inorganic reactant in reaction 2 (1)
- 4.2.2 The IUPAC name of compound **B** (2)
- 4.2.3 The type of reaction represented by reaction 1 (1)

Compound **B** is now used in two-step reaction to prepare compound **A**.

Reaction 4 is a SUBSTITUTION reaction.




Write down:

- 4.2.4 The NAME or FORMULA of the catalyst used in reaction 3 (1)
- 4.2.5 The IUPAC name of compound **D** (2)
- 4.2.6 The type of reaction represented by reaction 3 (1)
- 4.2.7 The type of haloalkane represented by compound **A** (Choose from primary, secondary or tertiary.) (1)

[20]

QUESTION 5 (Start on a new page.)

The reaction between aluminium and EXCESS sulphuric acid is used to investigate factors affecting rates of reactions.




5.1 INVESTIGATION I

The effect of a catalyst on the rate of reaction is determined.

Aluminium powder of mass 5 g reacts with excess $0,1 \text{ mol}\cdot\text{dm}^{-3} \text{ H}_2\text{SO}_4$ at $60 \text{ }^\circ\text{C}$.

Consider the following energy diagrams (not drawn to scale) for this investigation. **X** and **Y** in diagram **B** represent the activation energies.



- 5.1.1 Is the reaction between $\text{Al}(s)$ and dilute $\text{H}_2\text{SO}_4(aq)$ ENDOTHERMIC or EXOTHERMIC? Give a reason for the answer by referring to the above diagrams. (2)
- 5.1.2 What does the shaded area to the right of line **P** represent? (1)
- 5.1.3 Determine the numerical value represented by the letter **X** on diagram **B**. (2)

5.2 **INVESTIGATION II**


The investigation is now repeated at 30 °C using the same reactants (5g Al powder and excess 0,1 mol·dm⁻³ H₂SO₄) and catalyst.

How will this affect EACH of the following when compared to INVESTIGATION I? Choose from INCREASES, DECREASES or REMAINS THE SAME.

- 5.2.1 The size of the shaded area (diagram B) (1)
- 5.2.2 The value of Y (1)
- 5.2.3 The TOTAL volume of hydrogen gas produced (1)

5.3 **INVESTIGATION III**

In this investigation, 5 g of the same sample of IMPURE aluminium powder reacts with an EXCESS diluted H₂SO₄ at 60 °C in each of three runs. The table below summarises the conditions and the results obtained. (Assume that the impurities do not react.)



RUN	CONCENTRATION H ₂ SO ₄ (aq) (mol·dm ⁻³)	AVERAGE RATE OF VOLUME H ₂ (g) PRODUCED (cm ³ ·s ⁻¹)
1	0,1	15
2	0,2	19
3	0,4	40


- 5.3.1 Write down the independent variable for this investigation. (1)
- 5.3.2 Use the collision theory to explain how the average rate of the reaction is affected in this investigation. (3)
- 5.3.3 The time for the reaction to reach completion in RUN 3 is 2,6 minutes.

Calculate the percentage purity of the aluminium. Take the molar gas volume at 60 °C to be 27 000 cm³·mol⁻¹.

(6)
[18]


QUESTION 6 (Start on a new page.)

Consider the balanced equation for a hypothetical reaction that takes place in 2 dm^3 sealed containers.



The graphs below, not drawn to scale, are obtained for the same reaction at two different temperatures.

Graph **A** is obtained at 298 K and graph **B** at 398 K.



- 6.1 State Le Chatelier's principle. (2)
- 6.2 What do the parallel lines after $t = 5$ minutes in graph **A** represent? (1)
- 6.3 Is the forward reaction EXOTHERMIC or ENDOTHERMIC? (1)
- 6.4 Explain the answer to QUESTION 6.3. (2)
- 6.5 How does the value of the equilibrium constant, K_c , for the reaction in graph **B** compare to that in graph **A**? Choose from GREATER THAN, LESS THAN or EQUAL TO. (1)
- 6.6 The equilibrium constant, K_c , is 0,49 at 398 K (graph **B**).
 Calculate the initial number of moles of P_2Q . (8)
- 6.7 Describe the change made to the equilibrium system at $t = 8$ minutes, as shown in graph **B**, at a constant temperature. (1)
- 6.8 Explain by using Le Chatelier's principle how the system reacts to the change in QUESTION 6.7. (2)

[18]

QUESTION 7 (Start on a new page.)

7.1 A standard solution is prepared by dissolving 10 g of sodium carbonate, $\text{Na}_2\text{CO}_3(\text{s})$, in $0,7 \text{ dm}^3$ of water.

7.1.1 Calculate the concentration of the solution. (3)

7.1.2 Will the pH of the solution be GREATER THAN or LESS THAN 7? (1)


7.1.3 Write an equation that explains the answer to QUESTION 7.1.2. (2)

The sodium carbonate solution is titrated with dilute hydrochloric acid, $\text{HCl}(\text{aq})$. The following indicators are available for this titration.

INDICATOR	pH RANGE
P	3,4–4,5
Q	6,8–7,2
R	8,3–10

7.1.4 Which ONE of the indicators (**P**, **Q** or **R**) is most suitable for this titration? Give a reason for the answer by referring to the data in the table. (2)

7.2 When 0,01 moles of *dilute* sulphuric acid, $\text{H}_2\text{SO}_4(\text{aq})$, is mixed with 0,024 moles of potassium hydroxide, $\text{KOH}(\text{aq})$, the total volume of the final solution is $0,2 \text{ dm}^3$.




7.2.1 What is meant by a *dilute* acid? (2)

7.2.2 Calculate the pH of the final solution. (8)


[18]

QUESTION 8 (Start on a new page.)

The relationship between the concentration of the electrolyte and the cell potential is investigated using the following electrochemical cell represented by the cell notation:



The concentration of M^{2+} is changed and the corresponding emf is measured. The concentration of $\text{Al}^{3+}(\text{aq})$ and the temperature are at standard conditions. The graph below shows the results of this investigation.



- 8.1 Identify the reducing agent in this cell. (1)
- 8.2 Determine the concentration of $\text{M}^{2+}(\text{aq})$ that will produce an emf of 1,87 V. (2)
- 8.3 How will the concentration of $\text{M}^{2+}(\text{aq})$ be affected as the cell operates? Choose from INCREASES, DECREASES or REMAINS THE SAME. (2)
- Give a reason for the answer.
- 8.4 Potassium nitrate, $\text{KNO}_3(\text{aq})$, is used in the salt bridge of this cell. (1)
- To which electrode will the K^+ ions move in the salt bridge (Al or M)?

8.5 Identify metal **M** with the aid of a calculation. (6)


8.6 Metal **M** is now replaced with magnesium, Mg.

8.6.1 Which electrode, Al or Mg, will be the anode? (1)

8.6.2 Refer to the relative strengths of the oxidising agents to explain the answer. (2)
[15]

QUESTION 9 (Start on a new page.)

The simplified diagram below represents the cell used for electroplating ornaments with silver, Ag. **P** and **Q** are the two terminals of the battery.



9.1 State the energy conversion that takes place in this cell. (1)

9.2 Which terminal of the battery (**P** or **Q**) is negative? (1)

9.3 Write down the equation for the half-cell reaction that takes place at the cathode. (2)

9.4 Calculate the current needed to electroplate the ornament with 3,25 g of silver in 30 minutes. (5)
[9]

TOTAL: 150



**DATA FOR PHYSICAL SCIENCES GRADE 12
 PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12
 VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at/by 298 K	
$E_{\text{cell}}^\theta = E_{\text{cathode}}^\theta - E_{\text{anode}}^\theta$ / $E_{\text{sel}}^\theta = E_{\text{katode}}^\theta - E_{\text{anode}}^\theta$ or/of $E_{\text{cell}}^\theta = E_{\text{reduction}}^\theta - E_{\text{oxidation}}^\theta$ / $E_{\text{sel}}^\theta = E_{\text{reduksie}}^\theta - E_{\text{oksidasie}}^\theta$ or/of $E_{\text{cell}}^\theta = E_{\text{oxidisingagent}}^\theta - E_{\text{reducingagent}}^\theta$ / $E_{\text{sel}}^\theta = E_{\text{oksideermiddel}}^\theta - E_{\text{reduseermiddel}}^\theta$	
$I = \frac{Q}{\Delta t}$	$n = \frac{Q}{q_e}$ where n is the number of electrons/ waar n die aantal elektrone is

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE



KEY/SLEUTEL

Atomic number
Atoomgetal

Electronegativity
Elektronegatiwiteit

Symbol
Simbool

29
1,9 Cu
63,5

Approximate relative atomic mass
Benaderde relatiewe atoommassa

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)
2,1 1 H 1																	2 He 4
1,0 3 Li 7	1,5 4 Be 9											2,0 5 B 11	2,5 6 C 12	3,0 7 N 14	3,5 8 O 16	4,0 9 F 19	10 Ne 20
0,9 11 Na 23	1,2 12 Mg 24											1,5 13 Al 27	1,8 14 Si 28	2,1 15 P 31	2,5 16 S 32	3,0 17 Cl 35,5	18 Ar 40
0,8 19 K 39	1,0 20 Ca 40	1,3 21 Sc 45	1,5 22 Ti 48	1,6 23 V 51	1,6 24 Cr 52	1,5 25 Mn 55	1,8 26 Fe 56	1,8 27 Co 59	1,8 28 Ni 59	1,9 29 Cu 63,5	1,6 30 Zn 65	1,6 31 Ga 70	1,8 32 Ge 73	2,0 33 As 75	2,4 34 Se 79	2,8 35 Br 80	36 Kr 84
0,8 37 Rb 86	1,0 38 Sr 88	1,2 39 Y 89	1,4 40 Zr 91	1,6 72 Hf 179	1,8 42 Nb 92	1,9 43 Mo 96	2,2 44 Ru 101	2,2 45 Rh 103	2,2 46 Pd 106	1,9 47 Ag 108	1,7 48 Cd 112	1,7 49 In 115	1,8 50 Sn 119	1,9 51 Sb 122	2,1 52 Te 128	2,5 53 I 127	54 Xe 131
0,7 55 Cs 133	0,9 56 Ba 137	57 La 139					76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	1,8 81 Tl 204	1,8 82 Pb 207	1,9 83 Bi 209	2,0 84 Po	2,5 85 At	86 Rn
0,7 87 Fr	0,9 88 Ra 226	89 Ac															
58 Ce 140	59 Pr 141	60 Nd 144	61 Pm	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175				
90 Th 232	91 Pa	92 U 238	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr				

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE



Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels



Half-reactions/Halfreaksies	E^θ (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- \rightleftharpoons Pt$	+ 1,20
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^- \rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	- 2,36
$Na^+ + e^- \rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	- 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	- 2,93
$Li^+ + e^- \rightleftharpoons Li$	- 3,05

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels



TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE



Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies	E^θ (V)
$\text{Li}^+ + e^- \rightleftharpoons \text{Li}$	- 3,05
$\text{K}^+ + e^- \rightleftharpoons \text{K}$	- 2,93
$\text{Cs}^+ + e^- \rightleftharpoons \text{Cs}$	- 2,92
$\text{Ba}^{2+} + 2e^- \rightleftharpoons \text{Ba}$	- 2,90
$\text{Sr}^{2+} + 2e^- \rightleftharpoons \text{Sr}$	- 2,89
$\text{Ca}^{2+} + 2e^- \rightleftharpoons \text{Ca}$	- 2,87
$\text{Na}^+ + e^- \rightleftharpoons \text{Na}$	- 2,71
$\text{Mg}^{2+} + 2e^- \rightleftharpoons \text{Mg}$	- 2,36
$\text{Al}^{3+} + 3e^- \rightleftharpoons \text{Al}$	- 1,66
$\text{Mn}^{2+} + 2e^- \rightleftharpoons \text{Mn}$	- 1,18
$\text{Cr}^{2+} + 2e^- \rightleftharpoons \text{Cr}$	- 0,91
$2\text{H}_2\text{O} + 2e^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$	- 0,83
$\text{Zn}^{2+} + 2e^- \rightleftharpoons \text{Zn}$	- 0,76
$\text{Cr}^{3+} + 3e^- \rightleftharpoons \text{Cr}$	- 0,74
$\text{Fe}^{2+} + 2e^- \rightleftharpoons \text{Fe}$	- 0,44
$\text{Cr}^{3+} + e^- \rightleftharpoons \text{Cr}^{2+}$	- 0,41
$\text{Cd}^{2+} + 2e^- \rightleftharpoons \text{Cd}$	- 0,40
$\text{Co}^{2+} + 2e^- \rightleftharpoons \text{Co}$	- 0,28
$\text{Ni}^{2+} + 2e^- \rightleftharpoons \text{Ni}$	- 0,27
$\text{Sn}^{2+} + 2e^- \rightleftharpoons \text{Sn}$	- 0,14
$\text{Pb}^{2+} + 2e^- \rightleftharpoons \text{Pb}$	- 0,13
$\text{Fe}^{3+} + 3e^- \rightleftharpoons \text{Fe}$	- 0,06
$2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+ 0,14
$\text{Sn}^{4+} + 2e^- \rightleftharpoons \text{Sn}^{2+}$	+ 0,15
$\text{Cu}^{2+} + e^- \rightleftharpoons \text{Cu}^+$	+ 0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2e^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+ 0,17
$\text{Cu}^{2+} + 2e^- \rightleftharpoons \text{Cu}$	+ 0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4e^- \rightleftharpoons 4\text{OH}^-$	+ 0,40
$\text{SO}_2 + 4\text{H}^+ + 4e^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$	+ 0,45
$\text{Cu}^+ + e^- \rightleftharpoons \text{Cu}$	+ 0,52
$\text{I}_2 + 2e^- \rightleftharpoons 2\text{I}^-$	+ 0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2\text{O}_2$	+ 0,68
$\text{Fe}^{3+} + e^- \rightleftharpoons \text{Fe}^{2+}$	+ 0,77
$\text{NO}_3^- + 2\text{H}^+ + e^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+ 0,80
$\text{Ag}^+ + e^- \rightleftharpoons \text{Ag}$	+ 0,80
$\text{Hg}^{2+} + 2e^- \rightleftharpoons \text{Hg}(\ell)$	+ 0,85
$\text{NO}_3^- + 4\text{H}^+ + 3e^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+ 0,96
$\text{Br}_2(\ell) + 2e^- \rightleftharpoons 2\text{Br}^-$	+ 1,07
$\text{Pt}^{2+} + 2e^- \rightleftharpoons \text{Pt}$	+ 1,20
$\text{MnO}_2 + 4\text{H}^+ + 2e^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+ 1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4e^- \rightleftharpoons 2\text{H}_2\text{O}$	+ 1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6e^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+ 1,33
$\text{Cl}_2(\text{g}) + 2e^- \rightleftharpoons 2\text{Cl}^-$	+ 1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5e^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+ 1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2e^- \rightleftharpoons 2\text{H}_2\text{O}$	+ 1,77
$\text{Co}^{3+} + e^- \rightleftharpoons \text{Co}^{2+}$	+ 1,81
$\text{F}_2(\text{g}) + 2e^- \rightleftharpoons 2\text{F}^-$	+ 2,87

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**SENIOR CERTIFICATE EXAMINATIONS/
NATIONAL SENIOR CERTIFICATE EXAMINATIONS
SENIORSERTIFIKAAT-EKSAMEN/
NASIONALE SENIORSERTIFIKAAT-EKSAMEN**

**PHYSICAL SCIENCES: CHEMISTRY (P2)
FISIESE WETENSKAPPE: CHEMIE (V2)**

MAY/JUNE/MEI/JUNIE 2024

MARKING GUIDELINES/NASIENRIGLYNE

Stanmorephysics.com

MARKS/PUNTE: 150

**These marking guidelines consist of 20 pages./
Hierdie nasienriglyne bestaan uit 20 bladsye.**

Approved
M. Koen
DBE Moderator
22-5-24

Approved
[Signature]
DBE Moderator
22-5-24

Approved
[Signature]
Umahlali Ext. Mod.
22/5/24

Approved!
[Signature]
Umahlali Ext. Moderator
22/5/24


QUESTION 1/VRAAG 1

- 1.1 A ✓✓ (2)
- 1.2 C ✓✓ (2)
- 1.3 A ✓✓ (2)
- 1.4 D ✓✓ (2)
- 1.5 D ✓✓ (2)
- 1.6 C ✓✓ (2)
- 1.7 B ✓✓ (2)
- 1.8 A ✓✓ (2)
- 1.9 B ✓✓ (2)
- 1.10 B ✓✓ (2)
- [20]**


QUESTION 2/VRAAG 2

- 2.1 Organic compounds that consist of hydrogen and carbon only. ✓✓ (2 or 0)
 Organiese verbindings wat slegs uit waterstof en koolstof bestaan. (2 of 0) (2)
- 2.2.1 C and/en E ✓ (1)
- 2.2.2 D and/en H ✓✓ (2 or/of 0) (2)
- 2.2.3 A ✓ (1)

2.3	Marking criteria/Nasienkriteria:	IFIINDIEN:
2.3.1	<ul style="list-style-type: none"> Functional group. ✓ Funksionele groep. Whole structure correct. ✓ Hele struktuur korrek. 	<ul style="list-style-type: none"> More than one functional group/wrong functional group: Meer as een funksionele groep/foutiewe funksionele groep: $\frac{0}{2}$ If condensed structural formulae used/Indien gekondenseerde struktuurformules gebruik: Max/Maks. $\frac{1}{2}$




OR/OF



(2)

2.3.2 C_nH_{2n+2} ✓

2.3.3



Marking criteria/Nasienkriteria:

- Functional group $-C\equiv C-$. ✓
Funksionele groep $-C\equiv C-$.
- Whole structure correct. ✓
Hele struktuur korrek.

IF/INDIEN

- More than one functional group/wrong functional group:
Meer as een funksionele groep/foutiewe funksionele groep: $\frac{0}{2}$
- If condensed structural formulae used/*Indien gekondenseerde struktuurformules gebruik:*
 Max/Maks. $\frac{1}{2}$

(1)

(2)

2.4.1 3-ethylhex-3-ene ✓✓✓/3-ethyl-3-hexene/3-etielheks-3-ene/3-etiel-3-hekseen

Marking criteria:

- Correct stem i.e. hexene. ✓
- Substituent (ethyl) correctly identified. ✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam d.i. hekseen. ✓
- Substituent (etiel) korrek geïdentifiseer. ✓
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekens en kommas. ✓

(3)

2.4.2 2,5-dichloro-2,4-dimethylhexane ✓✓✓/ 2,5-dichloro-2,4-dimietielheksaan

Marking criteria:

- Correct stem i.e. hexane. ✓
- All substituents (dichloro and dimethyl) correctly identified. ✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam d.i. heksaan. ✓
- Alle substituenten (dichloro en dimietiel) korrek geïdentifiseer. ✓
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekens en kommas. ✓

(3)

2.4.3 2,2-dimethyl✓propanal✓/dimethylpropanal

2,2-dimietielpropanaal/dimietielpropanaal

NOTE/NOTA:

2,2-dimethyl✓propan-1-al (Max/Maks: $\frac{1}{2}$)

(2)

N. V. S. P. K.

2.5

Marking criteria/Nasienkriteria:

- Correct molecular formula: C_7H_{16} ✓
Korrekte molekulêre formula: C_7H_{16}
- Correct molecular formula of inorganic reactant and products. ✓
Korrekte molekulêre formule vir die anorganiese reaktans en produkte.
- Balancing/Balansering ✓



Notes/Aantekeninge:

- Ignore double arrows and phases. *Ignoreer dubbelpyle en fases.*
- Marking rule 6.3.10/Nasienreël 6.3.10.
- If condensed structural formulae used: *Indien gekondenseerde struktuurformules gebruik:* Max/Maks. $\frac{2}{3}$
- **ACCEPT:** multiple coefficients for this exam.
AANVAAR: veelvoude van koëffisiënte vir hierdie eksamen.

(3)
[22]

QUESTION 3/VRAAG 3

3.1

Marking criteria/Nasienkriteria

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark. *Indien enige van die onderstreepte frases in die korrekte konteks uitgelaat is, trek 1 punt af.*

The underlined phrases must be in the correct context. *Die onderstreepte frases moet in die korrekte konteks wees.*

The temperature at which the vapour pressure (of a substance) equals atmospheric pressure. ✓✓

Die temperatuur waarby die dampdruk (van die stof) gelyk is aan atmosferiese druk.

(2)

3.2

C ✓

(1)

N.
Vig
G
K

3.3

Marking criteria:

- Compare structures. ✓
- Compare the strength of intermolecular forces. ✓
- Compare the energy required to overcome intermolecular forces. ✓

Nasienkriteria:

- Vergelyk strukture. ✓
- Vergelyk die sterkte van intermolekulêre kragte. ✓
- Vergelyk die energie benodig om intermolekulêre kragte te oorkom. ✓

Accept: IMF for this exam/**Aanvaar:** IMK vir hierdie eksamen

A/CH₃CH₂CH₂CH₂Cl /1-chlorobutane

- **Structure:**
Longer chain length/larger surface area (over which intermolecular forces act). ✓
- **Intermolecular forces:**
Stronger/more intermolecular forces/Van der Waals forces/London forces/dipole-dipole forces. ✓
- **Energy:**
More energy needed to overcome or break intermolecular forces/Van der Waals forces/dipole-dipole forces. ✓

OR

B/CH₃CH(CH₃)CH₂Cl/1-chloro-2-methylpropane

- **Structure:**
Shorter chain length / branched / compact / more spherical / smaller surface area (over which intermolecular forces act). ✓
- **Intermolecular forces:**
Weaker/less intermolecular forces/Van der Waals forces/London forces/dipole-dipole forces. ✓
- **Energy:**
Less energy needed to overcome or break intermolecular forces/Van der Waals forces/dipole-dipole forces. ✓

A/CH₃CH₂CH₂CH₂Cl /1-chlorobutaan

- **Struktuur:**
Langer kettinglengte/groter oppervlak (waaroor intermolekulêre kragte werk). ✓
- **Intermolekulêre kragte:**
Sterker/meer intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte. ✓
- **Meer energie benodig om intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte te oorkom/breek.** ✓

OF

B/CH₃CH(CH₃)CH₂Cl/1-chloro-2-metielpropan

- **Struktuur:**
Korter kettinglengte / vertak / kompak / meer sferies / kleiner oppervlak (waaroor intermolekulêre kragte werk). ✓
- **Intermolekulêre kragte:**
Swakker/minder intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte. ✓
- **Energie:**
Minder energie benodig om intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte te oorkom/breek. ✓

N.
V.
G.
K.

(3)

3.4.1 75 (°C) ✓

(1)

3.4.2

Marking criteria:

- Compare the strength of intermolecular forces. ✓
- Compare the energy required to overcome intermolecular forces. ✓

Nasienkriteria:

- Vergelyk die sterkte van intermolekulêre kragte. ✓
- Vergelyk die energie benodig om intermolekulêre kragte te oorkom. ✓

• **Intermolecular forces:**

C ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ /butanol) has stronger intermolecular forces than D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ /butanal). ✓

• **Energy:**

More energy needed to overcome or break intermolecular forces. ✓

Accept: Boiling point of C will be more (in relation to C and D/118°C vs 75°C).

OR

• **Intermolecular forces:**

D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ /butanal) has weaker intermolecular forces than C ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ /butanol)

• **Energy:**

Less energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to C and D/118°C vs 75°C).

OR

• **Intermolecular forces:**

A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is a more polar molecule than D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$) increasing the intermolecular forces

• **Energy:**

More energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to A and D).

OR

• **Intermolecular forces:**

Electron density of A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is greater than D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$) increasing the intermolecular forces

• **Energy:**

More energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to A and D).

• **Intermolekulêre kragte:**

C ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ /butanol) het sterker intermolekulêre kragte as D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ /butanaal). ✓

• **Meer energie benodig om intermolekulêre kragte te oorkom/breek.** ✓

Aanvaar: Kookpunt van D sal minder wees (met betrekking tot C en D)

OR

• **Intermolekulêre kragte:**

D ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ /butanaal) het swakker intermolekulêre kragte as C ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ /butanol).

• **Minder energie benodig om intermolekulêre kragte/te oorkom/breek.**

Aanvaar: Kookpunt van C sal meer wees (met betrekking tot C en D)

N.
Vus
G
K



OF

• **Intermolekulêre kragte:**

A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is 'n meer polêre molekule as D wat sterker intermolekulêre kragte tot gevolg het.

• Meer energie benodig om intermolekulêre kragte te oorkom/breek.

Aanvaar: Kookpunt van D sal minder wees (met betrekking tot A en D)

OF

• **Intermolekulêre kragte:**

Elektrondigtheid van A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is groter wat sterker intermolekulêre kragte tot gevolg het.

• Meer energie benodig om intermolekulêre kragte te oorkom/breek.

• Aanvaar: Kookpunt van D sal minder wees (met betrekking tot A en D)

(2)

3.5 Decreases/Neem af ✓

(1)

[10]

QUESTION 4/VRAAG 4

4.1

4.1.1 (Concentrated) sulphuric acid/ $\text{H}_2\text{SO}_4(\text{aq})$ ✓
(Gekonsentreerde) swawelsuur

(1)

4.1.2 Esterification / Condensation ✓ / Verestering / Esterifikasie / Kondensasie

(1)

4.1.3 **ANY TWO/ENIGE TWEE:**

• Alcohol/methanol/reactant is flammable/catches fire easily. ✓

Alkohol/metanol/reaktans is vlambaar/slaan maklik aan die brand.

• To heat evenly/A steady/controlled/gradual increase in temperature. ✓

Om eweredig/gekontroleerd/gelydelik te verhit/'n Eweredige toename in temperatuur.

• Alcohol/methanol will evaporate too quickly/is volatile.

Alkohol/metanol sal te vinnig verdamp/is vlugtig.

(2)

N.
V.
G.
K.

4.1.4

Marking criteria:


- Whole structural formula correct for propanoic acid. ✓
- Whole structural formula correct for methanol. ✓
- Functional group of ester correct. ✓
- Whole structural formula of ester correct. ✓
- H₂O ✓

Nasienkriteria:

- Hele struktuurformule vir propanoësuur korrek. ✓
- Hele struktuurformule vir metanol korrek. ✓
- Funksionele groep van ester korrek. ✓
- Hele struktuurformule van ester korrek. ✓
- H₂O ✓

IF/INDIEN

- Any error e.g. omission of all H atoms, condensed or semi structural formula/Enige fout bv. weglating van alle H-atome, gekondenseerde of semi-struktuurformule: Max/Maks. 2/5 (Functional group, H₂O/Funksionele groep, H₂O)
- Any additional reactants or products /Enige addisionele reaktanse of produkte: Subtract 1 mark./Trek 1 punt af.
- Molecular formulae used:/Molekulêre formule gebruik: Max/Maks. 1/5 (water)
- No arrows: The first two structures given are considered as reactants and can be marked/Geen pyltjie: die eerste twee strukture geskryf, word beskou as reaktanse en kan gemerk word.



(5)

4.1.5 Methyl ✓ propanoate ✓ /Metielpropanoaat

(2)

4.2.1 Hydrogen/H₂ ✓ /Waterstof(gas)

(1)

4.2.2 3,3-dimethyl ✓ but-1-ene ✓ /3,3-dimethyl-1-butene
3,3-dimetiel but-1-een /3,3-dimetiel-1-buteen

(2)

4.2.3 elimination **OR** dehydrohalogenation ✓ eliminasi **OF** dehidrohalogenering

(1)

4.2.4 H₂SO₄/H₃PO₄ **OR/OF** Sulphuric acid/Phosphoric acid ✓
Swawelsuur/Fosforsuur

(1)

4.2.5 3,3-dimethyl ✓ butan-2-ol ✓ /3,3-dimethyl-2-butanol
3,3-dimetiel butan-2-ol /3,3-dimetiel-2-butanol

(2)

4.2.6 Addition/hydration ✓ Addisie/hidrasie

(1)

4.2.7 Secondary ✓ /Sekondêr

(1)

[20]

N.
V.
K.

QUESTION 5/VRAAG 5

5.1.1 Exothermic/Eksotermies ✓

Lower (potential) energy of the products than reactants. $\Delta H < 0/\Delta H$ negative / $\Delta H = -121,7$ kJ/More energy is released than absorbed. ✓

Laer (potensiële) energie van produkte as die reaktanse. $\Delta H < 0/\Delta H$ negatief / $\Delta H = -121,7$ kJ/Meer energie word afgegee as wat opgeneem is. (2)

5.1.2 (The number of) particles with sufficient/enough (kinetic) energy (with a catalyst) OR $E_K \geq E_A$ (which can undergo effective collisions.) ✓

(Die hoeveelheid) deeltjies met genoeg/voldoende (kinetiese) energie (met 'n katalisator) OF $E_K \geq E_A$ (om effektiewe botsings te ondergaan). (1)

5.1.3 $240,8 - 208,2 = 32,6$ (kJ) ✓

IF: only answer award 2 marks//INDIEN: slegs antwoord gee 2 punte (2)

5.2

5.2.1 Decreases/Afneem ✓ (1)

5.2.2 Remains the same/Bly dieselfde ✓ (1)

5.2.3 Remains the same/Bly dieselfde ✓ (1)

5.3.1 Concentration (of sulphuric acid/ $H_2SO_4(aq)$)/Konsentrasie (van swawelsuur) ✓ (1)

5.3.2 • More (H_2SO_4) particles per unit volume. ✓

• More effective collisions per unit time./Higher frequency of effective collisions. ✓

• Higher reaction rate. ✓

OR

• Less (H_2SO_4) particles per unit volume. ✓

• Less effective collisions per unit time./Lower frequency of effective collisions. ✓

• Lower reaction rate ✓

• Meer (H_2SO_4) deeltjies per eenheid volume. ✓

• Meer effektiewe botsings per eenheidtyd./Hoër frekwensie van effektiewe botsings. ✓

• Hoër reaksietempo. ✓

OF

• Minder (H_2SO_4)-deeltjies per eenheid volume. ✓

• Minder effektiewe botsings per eenheidtyd./Laer frekwensie van effektiewe botsings. ✓

• Laer reaksietempo. ✓ (3)

N.
V.
S.
K.

5.3.3

Marking criteria:	Nasienkriteria:
<p>(a) Substitute (2,6)(60)(40) cm³ OR (156)(40) in rate formula ✓</p> <p>(b) Substitute 27 000 cm³ / 27 dm³ and volume in $n(\text{H}_2) = \frac{V}{V_m}$ ✓</p> <p>(c) USE mole ratio $n(\text{Al}) = \frac{2}{3}n(\text{H}_2)$ ✓</p> <p>(d) Substitution 27 and reacting mole in $n(\text{Al}) = \frac{m}{M}$ ✓</p> <p>(e) Substitution of $\frac{4,05}{5}(100)$ ✓</p> <p>(f) Final answer: 83,2 % ✓ Range: 81 – 83,3 %</p>	<p>(a) Vervang 2,6(60)(40) cm³ OF (156)(40) in tempo formule ✓</p> <p>(b) Vervang 27 000 cm³ / 27 dm³ en volume in $n(\text{H}_2) = \frac{V}{V_m}$ ✓</p> <p>(c) GEBRUIK molverhouding $n(\text{Al}) = \frac{2}{3}n(\text{H}_2)$ ✓</p> <p>(d) Vervang 27 en mol gereageer in $n(\text{Al}) = \frac{m}{M}$ ✓</p> <p>(e) Vervang van $\frac{4,05}{5}(100)$ ✓</p> <p>(f) Finale antwoord: 81 % ✓ Gebied: 81 – 83,3 %</p>
<p>OPTION 1/OPSIE 1:</p> <p>Rate/Tempo = $\frac{\Delta V_{\text{H}_2}}{\Delta t}$</p> <p>$40 = \frac{\Delta V_{\text{H}_2}}{2,6(60)}$ ✓ (a)</p> <p>$V(\text{H}_2) = 6\,240 \text{ cm}^3$</p> <p>$n(\text{H}_2) = \frac{V}{V_m}$</p> <p>$= \frac{6\,240}{27\,000}$ ✓ (b)</p> <p>$= 0,23 \text{ mol}$</p> <p>$n(\text{Al}) = \frac{2}{3}n(\text{H}_2)$</p> <p>$n(\text{Al}) = \frac{2}{3}(0,23)$ ✓ (c)</p> <p>$= 0,15 \text{ mol}$</p> <p>$n(\text{Al}) = \frac{m}{M}$</p> <p>$0,15 = \frac{m}{27}$ ✓ (d)</p> <p>$m = 4,05 \text{ g}$</p> <p>% purity/suiwerheid = $\frac{4,05}{5}(100)$ ✓ (e)</p> <p>$= 81 \%$ ✓ (f)</p>	<p>OPTION 2/OPSIE 2:</p> <p>rate H₂ = 40 cm³·s⁻¹</p> <p>Rate in $n(\text{H}_2) = \frac{V}{V_m}$</p> <p>$= \frac{40}{27\,000}$ ✓ (b)</p> <p>$= 0,00148 \text{ mol} \cdot \text{s}^{-1}$</p> <p>Rate(Al) = $\frac{2}{3}n(\text{H}_2)$</p> <p>$= \frac{2}{3}(0,00148)$ ✓ (c)</p> <p>$= 9,88 \times 10^{-4} \text{ mol} \cdot \text{s}^{-1}$</p> <p>$n(\text{Al}) = \frac{m}{M}$</p> <p>$9,88 \times 10^{-4} = \frac{m}{27}$ ✓ (d)</p> <p>$m = 0,0267 \text{ g} \cdot \text{s}^{-1}$</p> <p>Rate/Tempo = $\frac{\Delta m_{\text{Al}}}{\Delta t}$</p> <p>$0,0267 = \frac{\Delta m_{\text{Al}}}{2,6(60)}$</p> <p>$m(\text{Al}) = 4,16 \text{ g}$</p> <p>% purity/suiwerheid = $\frac{4,16}{5}(100)$ ✓ (e)</p> <p>$= 83,2 \%$ ✓ (f)</p>

N. Vind

QUESTION 6/VRAAG 6

6.1

Marking criteria/Nasienkriteria:

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark. / Indien enige van die onderstreepte frases in die **korrekte konteks** uitgelaat is, trek 1 punt af.

The underlined phrases must be in the correct context. / Die onderstreepte frases moet in die korrekte konteks wees.

When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will cancel/oppose the disturbance. ✓✓

Wanneer die ewewig in 'n geslote sisteem versteur word, sal die sisteem 'n nuwe ewewig instel deur die reaksie te bevoordeel wat die versteuring kanselleer/teenwerk.

IF "isolated" system -1/ **INDIEN:** "geïsoleerde" sisteem -1)

(2)

6.2

(Chemical) equilibrium/Concentrations of reactants and products remain constant./Rate of the forward and reverse reactions are equal. ✓

(Chemiese) ewewig/Konsentrasies van reaktante en produkte bly konstant./ Tempo van voorwaartse en terugwaartse reaksie is gelyk.

(1)

6.3

OPTION 1/OPSIE 1:

Exothermic/Eksotermies ✓



6.4

- With an increase in temperature the endothermic reaction is favoured. ✓
- The reverse reaction is favoured./ Equilibrium shifts to the left. / Reactants / $[P_2Q]$ increases OR Products / $[PQ_2]$ decreases ✓
- 'n Toename in temperatuur bevoordeel die endotermiese reaksie.
- Die terugwaartse reaksie word bevoordeel./ Ewewig skuif na links. / Reaktante / $[P_2Q]$ neem toe OF Produkte / $[PQ_2]$ neem af



6.5

Less than/Kleiner as ✓

OPTION 2/OPSIE 2:

Endothermic/Endotermies ✓



- With an increase in temperature the endothermic reaction is favoured. ✓
- The forward reaction is favoured./ Equilibrium shifts to the right. Reactants / $[PQ_2]$ increases OR Products / $[P_2Q]$ decreases ✓
- 'n Toename in temperatuur bevoordeel die endotermiese reaksie.
- Die voorwaartse reaksie word bevoordeel./ Ewewig skuif na regs./ Reaktante / $[PQ_2]$ neem toe OF Produkte / $[P_2Q]$ neem af



Greater than/Groter as ✓

(1)

(2)

(1)

N.
V.
G.
K.

6.6

METHOD 1/METODE 1: Using lines/Gebruik lyne

CALCULATIONS USING CONCENTRATION

Marking criteria:

- (a) Correct K_c expression (formulae in square brackets). ✓✓
(If solid is included deduct 1 mark)
- (b) Substitute 0,49 into K_c expression. ✓
- (c) Substitute equilibrium concentration (0,35) into correct K_c expression. ✓
- (d) Change in concentration/mole ✓
- (e) **USE** ratio: $P_2Q : 2PQ_2 = 1 : 2$ ✓
- (f) Substitute 2 dm^3 in $n = cV$. ✓
- (g) Final answer = 0,85 (mol) OR 1,11 (mol) OR 3,09 (mol) ✓

Nasienkriteria:

- (a) Korrekte K_c uitdrukking (formules in vierkantige hakies). ✓✓
(Indien vastestof invervang is, trek 1 punt af)
- (b) Vervang 0,49 in K_c -uitdrukking. ✓
- (c) Vervang ewewigkonsentrasie (0,35) in korrekte K_c -uitdrukking. ✓
- (d) Verandering in konsentrasie/mol ✓
- (e) **GEBRUIK** verhouding: $P_2Q : PQ_2 = 1 : 2$ ✓
- (f) Vervang 2 dm^3 in $n = cV$. ✓
- (g) Finale antwoord = 0,85 (mol) OF 1,11 (mol) OF 3,09 (mol) ✓

OPTION 1/OPSIE 1:

	P_2Q	PQ_2
Initial concentration ($\text{mol} \cdot \text{dm}^{-3}$) Aanvangskonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)	x	0
Change in concentration ($\text{mol} \cdot \text{dm}^{-3}$) Verandering in konsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)	0,175 ✓(e)	0,35
Equilibrium concentration ($\text{mol} \cdot \text{dm}^{-3}$) Ewewigskonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)	(d) x - 0,175	0,35

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$0,49 \quad \checkmark \text{ (b)} = \frac{(0,35)^2 \quad \checkmark \text{ (c)}}{(x - 0,175)}$$

$$x = 0,425 \text{ mol} \cdot \text{dm}^{-3}$$

$$n(P_2Q) = cV$$

$$= 0,425 \times 2 \quad \checkmark \text{ (f)}$$


$$= 0,85 \text{ mol} \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. $\frac{6}{8}$

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. $\frac{5}{8}$

N.
Vid
G
K

OPTION 2/OPSIE 2:



$$K_c = \frac{[PQ_2]^2}{[P_2Q]}$$

$$0,49 = \frac{(0,35)^2}{P_2Q}$$

$$P_2Q = 0,25 \text{ mol}\cdot\text{dm}^{-3}$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. $\frac{6}{8}$

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. $\frac{5}{8}$

	P ₂ Q	PQ ₂
Initial concentration (mol·dm ⁻³) Aanvangskonsentrasie (mol·dm ⁻³)	0,425 ✓ (d)	0
Change in concentration (mol·dm ⁻³) Verandering in konsentrasie (mol·dm ⁻³)	-0,175	0,35 ✓ (e)
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,25	0,35

$$n(P_2Q) = cV$$

$$= 0,425(2) \quad \checkmark \text{ (f)}$$

$$= 0,85 \text{ mol} \quad \checkmark \text{ (g)}$$

CALCULATIONS USING NUMBER OF MOLES

OPTION 3/OPSIE 3:

	P ₂ Q	PQ ₂
Initial quantity (mol) Aanvangshoeveelheid (mol)	x	0
Change (mol) Verandering (mol)	0,35 ✓ (e)	0,7
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	x - 0,35 ✓ (d)	0,7
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	$\frac{x - 0,35}{2}$ ✓ (f)	0,35

$$K_c = \frac{[PQ_2]^2}{[P_2Q]}$$

$$0,49 = \frac{(0,35)^2}{\left(\frac{x - 0,35}{2}\right)}$$

$$x = 0,85 \text{ mol} \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. $\frac{6}{8}$

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. $\frac{5}{8}$

N.
Ves
h
K

OPTION 4/OPSIE 4:

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$0,49 \checkmark \text{ (b)} = \frac{(0,35)^2}{[P_2Q]} \quad \checkmark \text{ (c)}$$

$$[P_2Q] = 0,25 \text{ mol} \cdot \text{dm}^{-3}$$

Wrong K_c expression/

Verkeerde K_c -uitdrukking: Max./Maks. $\frac{5}{8}$

No K_c expression, correct substitution/Geen K_c -
 uitdrukking, korrekte substitusie: Max./Maks. $\frac{6}{8}$

	P ₂ Q	PQ ₂
Initial quantity (mol) Aanvangshoeveelheid (mol)	$\checkmark \text{ (g)}$ 0,85	0
Change (mol) Verandering (mol)	$\checkmark \text{ (e)}$ -0,35	0,7 $\checkmark \text{ (d)}$
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	0,5	0,7
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,25	0,35

METHOD 2/METODE 2: Using labels/Gebruik byskrifte

OPTION 1/OPSIE 1:

	P ₂ Q	PQ ₂
Initial concentration (mol·dm ⁻³) Aanvangskonsentrasie (mol·dm ⁻³)	x	0
Change in concentration (mol·dm ⁻³) Verandering in konsentrasie (mol·dm ⁻³)	$\checkmark \text{ (e)}$ -0,207	0,414 $\checkmark \text{ (d)}$
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,35	0,414

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$\text{(b)} \quad \checkmark \quad 0,49 = \frac{[PQ_2]^2}{(0,35)} \quad \checkmark \text{ (c)}$$

$$[PQ_2] = 0,414 \text{ mol} \cdot \text{dm}^{-3}$$

$$\text{initial } n(P_2Q) = (0,35 + 0,207)(2) \quad \checkmark \text{ (f)}$$

$$= 1,11 \text{ mol} \quad \checkmark \text{ (g)}$$

No K_c expression, correct
 substitution/Geen K_c -
 uitdrukking,
 korrekte substitusie: Max./Maks. $\frac{6}{8}$

Wrong K_c expression/

Verkeerde K_c -uitdrukking: Max./Maks. $\frac{5}{8}$

N.
V.
K.

OPTION 2/OPSIE 2:

	P ₂ Q	PQ ₂
Initial quantity (mol) Aanvangshoeveelheid (mol)	x	0
Change (mol) Verandering (mol)	0,414	0,828 ✓ (e)
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	0,7	0,828
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,35 ✓ (f)	0,414

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$(b) \checkmark 0,49 = \frac{[PQ_2]^2}{(0,35)} \quad \checkmark \text{ (c)}$$

$$[PQ_2] = 0,414 \text{ mol} \cdot \text{dm}^{-3}$$

$$x - 0,414 = 0,7 \quad \checkmark \text{ (d)}$$


$$x = 1,11 \text{ mol } P_2Q \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c-uitdrukking, korrekte substitusie: Max./Maks. 6/8

Wrong K_c expression/

Verkeerde K_c-uitdrukking: Max./Maks. 5/8

METHOD 3/METODE 3: (Equation written as reverse/Vergelyking omgekeerd geskryf)



	PQ ₂	P ₂ Q
Initial quantity (mol) Aanvangshoeveelheid (mol)	x	0
Change (mol) Verandering (mol)	1,4	0,7 ✓ (e)
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol) (d) ✓	x - 1,4	0,7
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	$\frac{x - 1,4}{2}$	0,35

$$K_c = \frac{[P_2Q]}{[PQ_2]^2} \quad \checkmark \checkmark \text{ (a)}$$

$$(b) \checkmark 0,49 = \frac{(0,35)}{[PQ_2]^2} \quad \checkmark \text{ (c)}$$

$$\frac{x - 1,4}{2} = 0,845 \quad \checkmark \text{ (f)}$$

$$x = 3,09 \text{ mol } P_2Q \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c-uitdrukking, korrekte substitusie: Max./Maks. 6/8

Wrong K_c expression/

Verkeerde K_c-uitdrukking: Max./Maks. 5/8

METHOD 4/METODE 4: Reading from graph/Af lees van grafiek

OPTION 1/OPSIE 1

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} = 0,49 \quad \text{Initial } [P_2Q] = 0$$

$$n = 0 \text{ (mol)} \quad \checkmark \checkmark \checkmark \checkmark \checkmark \checkmark \checkmark \checkmark \quad (8/8)$$

N.
Vind
dit

OPTION 2/OPSIE 2:

	P ₂ Q	PQ ₂
Initial concentration (mol·dm ⁻³) Aanvangskonsentrasie (mol·dm ⁻³)	0	y
Change in concentration (mol·dm ⁻³) Verandering in konsentrasie (mol·dm ⁻³)	-0,207	0,414
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,35	0,414

n = 0 (mol) ✓✓✓✓✓✓✓✓ (8/8)

(8)

6.7 Pressure was decreased/volume of the container was increased. ✓
Druk is verlaag/volume van die houer is vergroot.

(1)

6.8 **OPTION 1/OPSIE 1:** Using labels/Gebruik byskrifte

- Favours the reaction that increases the number of moles (of gas) ✓/
Bevoordeel die reaksie wat aantal mol (gas) laat toeneem
- [P₂Q] increased/neem toe ✓ **OR/OF** [PQ₂] decreased/neem af

OPTION 2/OPSIE 2: Using lines/Gebruik lyne

- Favours the reaction that increases the number of moles (of gas) ✓/
Bevoordeel die reaksie wat aantal mol (gas) laat toeneem
- [PQ₂] increased/neem toe ✓ **OR/OF** [P₂Q] decreased/neem af

(2)

[18]

N.
Vid
U
K

QUESTION 7/VRAAG 7

7.1	<p>Marking criteria:</p> <ul style="list-style-type: none"> Any formula $c = \frac{m}{MV}$ or $n = \frac{m}{M}$ or $c = \frac{n}{V}$ ✓ Substitute <u>10, 106 and 0,7</u> into formula ✓ Final answer: $0,13 \text{ mol}\cdot\text{dm}^{-3}$ ✓ 	<p>Nasienkriteria:</p> <ul style="list-style-type: none"> Enige formule $c = \frac{m}{MV}$ of $n = \frac{m}{M}$ of $c = \frac{n}{V}$ ✓ Vervang <u>10, 106 and 0,7</u> in formula ✓ Finale antwoord: $0,13 \text{ mol}\cdot\text{dm}^{-3}$ ✓
7.1.1	<p>OPTION 1/OPSIE 1:</p> $c = \frac{m}{MV} \checkmark$ $= \frac{10}{(106)(0,7)} \checkmark$ $= 0,13 \text{ mol}\cdot\text{dm}^{-3} \checkmark$	<p>OPTION 2/OPSIE 2:</p> <p>$n = \frac{m}{M}$ Any one/Enige een ✓</p> $= \frac{10}{106} \checkmark$ $= 0,09 \checkmark$ <p>$c = \frac{n}{V}$</p> $= \frac{0,09}{0,7} \checkmark$ $= 0,13 \text{ mol}\cdot\text{dm}^{-3} \checkmark$

(3)

7.1.2 Greater than/Groter as ✓

(1)

7.1.3 $\text{CO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\ell) \checkmark \rightleftharpoons \text{HCO}_3^-(\text{aq}) + \text{OH}^-(\text{aq}) \checkmark$

OR/OF

$\text{CO}_3^{2-}(\text{aq}) + 2\text{H}_2\text{O}(\ell) \checkmark \rightleftharpoons \text{H}_2\text{CO}_3(\text{aq}) + 2\text{OH}^-(\text{aq}) \checkmark$

OR/OF

$\text{Na}_2\text{CO}_3(\text{aq}) + \text{H}_2\text{O}(\ell) \checkmark \rightleftharpoons \text{NaHCO}_3(\text{aq}) + \text{NaOH}(\text{aq}) \checkmark$

OR/OF

$\text{Na}_2\text{CO}_3(\text{aq}) + 2\text{H}_2\text{O}(\ell) \checkmark \rightleftharpoons \text{H}_2\text{CO}_3(\text{aq}) + 2\text{NaOH}(\text{aq}) \checkmark$

Marking criterial/Nasienkriteria:

- Reactants ✓ Products ✓
- Reaktanse ✓ Produkte ✓
- Ignore/Ignoreer → and phases/en fases
- Marking rule 6.3.10/Nasienreël 6.3.10

(2)

7.1.4 P ✓



(Titration of) weak base and a strong acid./The equivalence point is lower than pH 7. ✓

(Titrasië van) 'n swak basis en 'n sterk suur./ Die ekwivalente punt is laer as 'n pH van 7.

(2)

7.2

7.2.1 Dilute acid contains small amount/number of moles of acid in proportion to the volume of water. ✓✓ **(2 or/of 0)**

Verdunde sure bevat 'n klein hoeveelheid/getal mol suur in verhouding met die volume water.

(2)

N.
V.
K.

7.2.2

Marking criteria:	Nasienkriteria:
<p>(a) USE of ratio: $n(\text{KOH})_{\text{reacted}} = 2n(\text{H}_2\text{SO}_4)_{\text{reacted}}$ $[\text{KOH}]_{\text{reacted}} = 2n[\text{H}_2\text{SO}_4]_{\text{reacted}}$ ✓</p> <p>(b) Subtract: $n(\text{KOH})_{\text{initial}} - n(\text{KOH})_{\text{reacted}}$ $[\text{KOH}]_{\text{initial}} - [\text{KOH}]_{\text{reacted}}$ ✓✓</p> <p>(c) Divide n by 0,20 dm³ in c = $\frac{n}{V}$ ✓</p> <p>(d) Either formulae: $\text{pH} = -\log[\text{H}_3\text{O}^+]$ / $\text{pH} = -\log[\text{H}^+] / \text{pOH} = -\log[\text{OH}^-]$ AND $[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$ / $\text{pH} + \text{pOH} = 14$ ✓</p> <p>(e) Substitute calculated $[\text{OH}^-]$ in $[\text{H}_3\text{O}^+][\text{OH}^-]$ / in $\text{pOH} = -\log[\text{OH}^-]$ ✓</p> <p>(f) Substitute calculated $[\text{H}_3\text{O}^+]$ in pH formula/ pOH in $\text{pH} + \text{pOH} = 14$ ✓</p> <p>(g) Final answer: 12,3 ✓</p>	<p>(a) GEBRUIK verhouding: $n(\text{KOH})_{\text{gereageer}} = 2n(\text{H}_2\text{SO}_4)_{\text{gereageer}}$ $[\text{KOH}]_{\text{gereageer}} = 2n[\text{H}_2\text{SO}_4]_{\text{gereageer}}$ ✓</p> <p>(b) Aftrek: $n(\text{KOH})_{\text{aanvanklik}} - n(\text{KOH})_{\text{gereageer}}$ $[\text{KOH}]_{\text{aanvanklik}} - [\text{KOH}]_{\text{gereageer}}$ ✓✓</p> <p>(c) Deel n deur 0,20 dm³ in c = $\frac{n}{V}$ ✓</p> <p>(d) Enige een v formules: $\text{pH} = -\log[\text{H}_3\text{O}^+]$ / $\text{pH} = -\log[\text{H}^+] / \text{pOH} = -\log[\text{OH}^-]$ EN $[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$ / $\text{pH} + \text{pOH} = 14$ ✓</p> <p>(e) Vervang berekende $[\text{OH}^-]$ in $[\text{H}_3\text{O}^+][\text{OH}^-]$ / in $\text{pOH} = -\log[\text{OH}^-]$ ✓</p> <p>(f) Vervang berekende $[\text{H}_3\text{O}^+]$ in pH formule/ pOH in $\text{pH} + \text{pOH} = 14$ ✓</p> <p>(g) Finale antwoord: 12,3 ✓</p>
<p>OPTION 1/OPSIE 1:</p> <p>$n(\text{KOH})_{\text{reacted}} = 2n(\text{H}_2\text{SO}_4)_{\text{reacted}}$ $= 2(0,01)$ ✓ (a) $= 0,02$</p> <p>$n(\text{KOH})_{\text{excess}} = 0,024 - 0,02$ ✓✓ (b) $= 0,004 \text{ mol}$</p> <p>$[\text{OH}^-] = \frac{n}{V}$ $= \frac{0,004}{0,20}$ ✓ (c) $= 0,02 \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$ $[\text{H}_3\text{O}^+] (0,02) = 1 \times 10^{-14}$ ✓ (e) $[\text{H}_3\text{O}^+] = 5 \times 10^{-13} \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$\text{pH} = -\log[\text{H}_3\text{O}^+]$ $= -\log(5 \times 10^{-13})$ ✓ (f) $= 12,3$ ✓ (g)</p> <p style="text-align: right;">Either/ Enige een ✓ (d)</p>	<p>OPTION 2/OPSIE 2:</p> <p>$[\text{KOH}] = \frac{n}{V}$ $= \frac{0,024}{0,20}$ ✓ (c) $= 0,12 \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$[\text{H}_2\text{SO}_4] = \frac{n}{V}$ $= \frac{0,01}{0,20}$ $= 0,05 \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$[\text{KOH}]_{\text{reacted}} = 2[\text{H}_2\text{SO}_4]_{\text{reacted}}$ ✓ (a) $= 2(0,05)$ $= 0,1 \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$[\text{KOH}]_{\text{excess}} = 0,12 - 0,1$ ✓✓ (b) $= 0,02 \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14}$ $[\text{H}_3\text{O}^+] (0,02) = 1 \times 10^{-14}$ ✓ (e) $[\text{H}_3\text{O}^+] = 5 \times 10^{-13} \text{ mol}\cdot\text{dm}^{-3}$</p> <p>$\text{pH} = -\log[\text{H}_3\text{O}^+]$ $= -\log(5 \times 10^{-13})$ ✓ (f) $= 12,3$ ✓ (g)</p> <p style="text-align: right;">Both/ Beide ✓ (d)</p> <p style="text-align: right;">Either/ Enige een ✓ (d)</p>
<p>OPTION 3/OPSIE 3</p> <p>$\text{pOH} = -\log[\text{OH}^-]$ $\text{pOH} = -\log(0,02)$ ✓ (e) $\text{pOH} = 1,7$</p> <p>$\text{pH} + \text{pOH} = 14$ $\text{pH} + 1,7 = 14$ ✓ (f) $\text{pH} = 12,3$ ✓ (g)</p> <p style="text-align: right;">Any one/Enige een ✓ (d)</p>	

N. Vieg
 H
 K

QUESTION 8/VRAAG 8

8.1 Aluminium/Al ✓ (1)

8.2 0,325 (mol·dm⁻³) ✓✓
 Range/Gebied: 0,32 – 0,33 (mol·dm⁻³) (2)

8.3 Decreases / *Neem af* ✓
 M²⁺ is reduced/ M²⁺ used up/ M²⁺ is the oxidising agent. ✓ (2)
M²⁺ word gereduseer/ M²⁺ opgebruik/ M²⁺ is die oksideermiddel.

8.4 M ✓ (1)

<p>OPTION 1/OPTION 1</p> <p>$E_{\text{cell}}^{\theta} = E_{\text{reduction}}^{\theta} - E_{\text{oxidation}}^{\theta}$ ✓</p> <p>$2 \checkmark \checkmark = E_{\text{cathode}}^{\theta} - (-1,66)$ ✓</p> <p>$E_{\text{cathode}}^{\theta} = 0,34 \text{ (V)}$ ✓</p> <p>M is copper/Cu/<i>koper</i> ✓</p>	<p>NOTE/LET WEL</p> <ul style="list-style-type: none"> • Accept any other correct formula from the data sheet. <i>Aanvaar enige ander korrekte formule vanaf gegewensblad.</i> • Any other formula using unconventional abbreviations, e.g. $E^{\circ}_{\text{cell}} = E^{\circ}_{\text{OA}} - E^{\circ}_{\text{RA}}$ followed by correct substitutions: <i>Enige ander formule wat onkonvensionele afkortings gebruik, bv. $E^{\circ}_{\text{sel}} = E^{\circ}_{\text{OM}} - E^{\circ}_{\text{RM}}$ gevolg deur korrekte vervangings</i> ^{5/6} 						
<p>OPTION 2/OPSIE 2</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 2px;">$M^{2+}(\text{aq}) + 2e^{-} \rightarrow M(\text{aq})$</td> <td style="padding: 2px;">$E = +x \text{ V}$</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">$Al(\text{s}) \rightarrow Al^{3+}(\text{aq}) + 3e^{-}$</td> <td style="padding: 2px;">$E = +1,66 \text{ V}$ ✓</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">$2Al(\text{s}) + 3M^{2+}(\text{aq}) \rightarrow 2Al^{3+}(\text{aq}) + 3M(\text{s})$</td> <td style="padding: 2px;">$E = 2,00 \text{ (V)}$ ✓✓</td> </tr> </table> <p>$x = 0,34 \text{ (V)}$ ✓</p> <p>M is copper/Cu/<i>koper</i> ✓</p>		$M^{2+}(\text{aq}) + 2e^{-} \rightarrow M(\text{aq})$	$E = +x \text{ V}$	$Al(\text{s}) \rightarrow Al^{3+}(\text{aq}) + 3e^{-}$	$E = +1,66 \text{ V}$ ✓	$2Al(\text{s}) + 3M^{2+}(\text{aq}) \rightarrow 2Al^{3+}(\text{aq}) + 3M(\text{s})$	$E = 2,00 \text{ (V)}$ ✓✓
$M^{2+}(\text{aq}) + 2e^{-} \rightarrow M(\text{aq})$	$E = +x \text{ V}$						
$Al(\text{s}) \rightarrow Al^{3+}(\text{aq}) + 3e^{-}$	$E = +1,66 \text{ V}$ ✓						
$2Al(\text{s}) + 3M^{2+}(\text{aq}) \rightarrow 2Al^{3+}(\text{aq}) + 3M(\text{s})$	$E = 2,00 \text{ (V)}$ ✓✓						

8.6.1 Magnesium/Mg ✓ (1)

8.6.2 Al³⁺ is a stronger oxidising agent than Mg²⁺ ✓, therefore, Mg will be oxidised (to Mg²⁺).
 Mg²⁺ is a weaker oxidising agent than Al³⁺ ✓, therefore, Mg will be oxidised (to Mg²⁺).

Al³⁺ is 'n sterker oksideermiddel as Mg²⁺, daarom sal Mg geoksideer word (tot Mg²⁺)./

Mg²⁺ is 'n swakker oksideermiddel as Al³⁺, daarom sal Mg geoksideer word (tot Mg²⁺).

ACCEPT/AANVAAR:

Mg ion and Al ion/*Mg ioon en Al ioon*

(2)
[15]

*N.
 Vis
 G.
 K.*




QUESTION 9/VRAAG 9

9.1 Electrical to chemical (energy)/Elektriese na chemiese (energie) ✓ (1)

9.2 P ✓ (1)

9.3 $Ag^+ + e^- \rightarrow Ag$ ✓✓

Marking criteria/Nasienkriteria:



Ignore if charge omitted on electron. // Ignoreer indien lading weggelaat op elektron. (2)

9.4

Marking criteria:	Nasienkriteria:
<p>(a) Substitute 3,25 and 108 in the formula $n = \frac{m}{M}$ ✓</p> <p>(b) Substitute $6,02 \times 10^{23}$ in $n(e^-) = \frac{N}{N_A}$ ✓</p> <p>(c) Substitute 0,03 mol in $n(e^-) = \frac{N}{N_A}$ ✓ (Substitute 96 500 in formula $Q = nF$)</p> <p>(d) Substitute 30(60) OR 1 800 ✓</p> <p>(e) Final answer: 1,61 A ✓</p>	<p>(a) Vervang 3,25 en 108 in die formule $n = \frac{m}{M}$ ✓</p> <p>(b) Vervang $6,02 \times 10^{23}$ in $n(e^-) = \frac{N}{N_A}$ ✓</p> <p>(c) Vervang 0,03 mol in $n(e^-) = \frac{N}{N_A}$ ✓ (Vervang 96 500 in formule $Q = nF$)</p> <p>(d) Vervang 30(60) OF 1 800 ✓</p> <p>(e) Finale antwoord: 1,61 A ✓</p>
<p>OPTION 1/OPSIE 1:</p> $n(Ag) = \frac{m}{M}$ $= \frac{3,25}{108} \quad \checkmark(a)$ $= 0,03 \text{ mol}$ $n(e^-) = \frac{N}{N_A}$ $(c) \checkmark 0,03 = \frac{N}{6,02 \times 10^{23}} \quad \checkmark(b)$ $N e^- = 1,81 \times 10^{22}$ $N e^- = \frac{Q}{e} \quad \text{OF/OR} \quad \frac{Q}{q_e}$ $1,81 \times 10^{22} = \frac{Q}{1,6 \times 10^{-19}}$ $Q = 2\,889,6 \text{ C}$ $I = \frac{Q}{\Delta t}$ $= \frac{2\,889,6}{30(60)} \quad \checkmark(d)$ $= 1,61 \text{ A} \quad \checkmark(e)$	<p>OPTION 2/OPSIE 2:</p> $n(Ag) = \frac{m}{M}$ $= \frac{3,25}{108} \quad \checkmark(a)$ $= 0,03 \text{ mol} = n e^-$ $Q = 0,03 \times 96\,500 \quad \checkmark(c)$ $= 2\,895 \text{ C}$ $I = \frac{Q}{\Delta t}$ $= \frac{2\,895}{30(60)} \quad \checkmark(d)$ $= 1,61 \text{ A} \quad \checkmark(e)$

N. V. S. P. K.

(5)
 [9]
TOTAL/TOTAAL: 150