



NATIONAL
SENIOR CERTIFICATE

GRADE 12

PHYSICAL SCIENCES
CONTROL TEST 1
17 MARCH 2025

MARKS : 100

TIME : 2 HOURS

This question paper consists of 19 pages including data sheets.

INSTRUCTIONS AND INFORMATION

1. Write your name on the top of your ANSWER SCRIPT.
2. This question paper consists of 7 questions. Answer ALL the questions on your ANSWER SCRIPT.
3. You may use a non-programmable calculator.
4. You may use appropriate mathematical instruments.
5. Number the answers correctly according to the numbering system used in this QUESTION PAPER.
6. YOU ARE ADVISED TO USE THE ATTACHED DATA SHEETS.
7. Give brief motivations, discussions, et cetera where required.
8. Round off your final numerical answers to a minimum of TWO decimal places.

QUESTION 1 MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A – D) next to the question numbers. (1.1 – 1.10) in the ANSWER BOOK, e.g. 1.11 D.


1.1 Which one of the following physical quantities is a vector?

- A Impulse
- B Kinetic energy
- C Time
- D Mass



(2)

1.2 Which of the following position-time graphs represent the motion of an object moving with constant, non-zero velocity?



(2)

- 1.3 A person stands on a bathroom scale in a stationary elevator. The reading on the scale is 490 N. When the elevator is in motion, the reading on the scale changes to 470 N.

Which ONE of the following combinations best describes the DIRECTION OF THE MOTION and the DIRECTION OF THE ACCELERATION of the elevator during the motion?

	Direction of motion	Direction of acceleration
A	Upwards	Upwards
B	Downwards	Downwards
C	Upwards	Downwards
D	Downwards	Upwards


(2)

- 1.4 The gravitational acceleration, g , on or near the surface of the earth depends on the...

- A mass of the earth and the diameter of the earth
- B mass of the earth and the distance from its centre
- C mass of the earth only
- D distance from its centre

(2)

- 1.5 An object is thrown upwards with unknown velocity. The velocity time graph below shows the entire motion of the object.



At which point A – D will the object have the greatest momentum.

- A. A
- B. B
- C. C
- D. D

(2)

- 1.6 A ball is dropped downwards from the height, h and hits the ground with a speed v . The speed of the ball when it reaches half its initial height is...

- A v
- B $\frac{v}{\sqrt{2}}$
- C $\frac{v}{2}$
- D $\frac{v}{4}$

(2)

1.7 Consider the compound below:



The IUPAC name for the above compound is...

A 2,2,5-trimetielhex-3-yn

B 2,5,5-trimetielhex-3-yn

C 2,2-dimetiel hept-3-yn

D 2,5- dimetielhept-3-yn

(2)

1.8 Which one of the following reactions will produce an alkene?

A Esterification reaction

B Substitution reaction

C Elimination reaction

D Addition reaction

(2)

1.9 Which one of the following compounds has the lowest vapour pressure?

A Propane

B Propan-1-ol

C Propanoic acid

D Propanone

(2)

- 1.10 List of reactants and reaction conditions are given below for a learner to use so that he/she can produce a primary alcohol from **3-ethyl-3-iodo-2-methylpentane**.

Concentrated H₂SO₄, water, heat, concentrated KOH and dilute H₂SO₄


How will the learner use the above reactants and reaction condition to produce a primary alcohol in 4 steps?

	STEP 1	STEP 2	STEP 3	STEP 4
A	Heat and concentrated KOH	Water and dilute H ₂ SO ₄	Heat and concentrated H ₂ SO ₄	Water and dilute H ₂ SO ₄
B	Heat and Concentrated H ₂ SO ₄	Water and dilute H ₂ SO ₄	Heat and concentrated KOH	Water and dilute H ₂ SO ₄
C	Water and dilute H ₂ SO ₄	Heat and concentrated KOH	Heat and concentrated H ₂ SO ₄	Water and dilute H ₂ SO ₄
D	Water and dilute H ₂ SO ₄	Heat and concentrated KOH	Water and dilute H ₂ SO ₄	Heat and concentrated H ₂ SO ₄

(2)
[20]

QUESTION 2

A block of mass 3 kg is pulled from rest on a straight horizontal surface as shown below by a force $F = 10 \text{ N}$ which make an angle of 15° to the horizontal.




Path **AB** is $x \text{ m}$ and is frictionless and path **BC** is 4 m and is rough. The coefficient of kinetic friction between the block and the surface **BC** is 0,53. The block comes to rest at point **C**.

- 2.1 Draw a force diagram showing all the forces acting on the body as it slides on path **AB**. (3)
- 2.2 Calculate the speed of the block at point **B**. (5)
- 2.3 How will the speed calculated in 2.2 be affected if the angle between the surface and the force (F) was reduced to 10° ? Write INCREASE, DECREASE or REMAIN THE SAME. Give a reason for your answer. (2)
- 2.4 Draw the graph of F_{net} vs time for the entire motion of the block. No numerical values are required. (3)
[13]

QUESTION 3


Two blocks are moving to the right along a frictionless surface. Block X has a mass of 2 kg and is moving at $3 \text{ m}\cdot\text{s}^{-1}$. Block Y has a mass of 5 kg and is travelling at $1,5 \text{ m}\cdot\text{s}^{-1}$. The blocks collide and block Y continues in its original direction with a speed of $2,5 \text{ m}\cdot\text{s}^{-1}$ after their collision.



- 3.1 State the principle or law which is applicable AT THE POINT OF COLLISION. (2)
- 3.2 Give a reason why the speed of the 5 kg block increases after collision. (1)
- 3.3 Calculate the impulse experienced by the 5 kg block. (3)
- 3.4 Determine the magnitude of the velocity of block X after collision. (4)
[10]

QUESTION 4

A ball of mass 1.5 kg is thrown down from the top of the building with a velocity of 12 m.s^{-1} . The ball hit the ground and bounces off to the maximum height of 7 m. Ignore the effect of air resistance.




- 4.1 Define the term *free fall* in words. (2)
- 4.2 Calculate the velocity which the ball leaves (bounces) the ground with. (4)
- 4.3 When the ball was in contact with the ground, 344,1 J of its kinetic energy was converted into the other forms of energy.

Calculate the:

- 4.3.1 speed that the ball hit the ground with. HINT: Use ENERGY FORMULAE. (4)
- 4.3.2 time the ball will take to travel distance H. (3)
[13]

QUESTION 5

The letters A to F in the table below represent six organic compounds.

A $\begin{array}{ccccccc} & \text{CH}_2 & - & \text{CH} & - & \text{CH}_2 & - \text{CH}_2 \\ & & & & & & \\ & \text{CH}_3 & & \text{Cl} & & \text{CH}_3 & \end{array}$	B $\begin{array}{ccccc} & & \text{CH}_3 & & \\ & & & & \\ & & \text{CH}_3 & - \text{CH}_2 & - \text{CH}_2 & - \text{C} & - \text{CH}_3 \\ & & & & & & \\ & & & & & \text{OH} & \end{array}$
C  $\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}=\text{CH}-\text{CH}_3 \\ \\ \text{CH}_3 \end{array}$	D 4-methylpentanoic acid
E $\begin{array}{ccccccccc} & \text{H} & & \text{H} & & \text{H} & \text{O} & & \text{H} \\ & & & & & & \parallel & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - \text{C} & - & \text{C} & - \text{H} \\ & & & & & & & & \\ & \text{H} & & \text{C} & - & \text{H} & & & \text{H} \\ & & & & & & & & \\ & & & \text{H} & & & & & \end{array}$	F $\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_2 \\ \\ \text{O} \\ \\ \text{C}=\text{O} \\ \\ \text{CH}_2 \\ \\ \text{CH}_2 \\ \\ \text{CH}_3 \end{array}$

5.1 Write down the letter(s) that represent(s) each of the following:

5.1.1 compounds that can **PRODUCE** unsaturated hydrocarbon (2)

5.1.2 compound that uses platinum/palladium/nickel as a catalyst when it reacts. (1)

5.2 Write the IUPAC name of the following:

5.2.1 compound **A**. (2)

5.2.2 compound **E**. (3)

5.3 Draw the structural formula of functional group of compound **D**. (2)

5.4 Write the general formula of compound **C**. (1)

5.5 Consider compound **F**.





5.5.1 Write down the NAME of the catalyst used during preparation of compound **F**. (1)

5.5.2 Is the reaction for preparation of compound **F** ENDOTHERMIC or EXOTHERMIC? (1)

[13]

QUESTION 6


The relationship between strength of intermolecular forces and the boiling point is investigated using five organic compounds. The compounds and their boiling points are given in the table below.

Experiment	Compound/Molecular Formulae	Boiling point (°C)
1	A 	C ₅ H ₁₂ 36
	B 	C ₅ H ₁₂ 28
	C 	C ₅ H ₁₂ 10
2	D 	C ₄ H ₈ O 54
	E 	C ₄ H ₈ O 68

- 6.1 Define the term *boiling point*. (2)
- 6.2 Consider experiment 2.
- To which homologous series does compound **D** and **E** belong? (1)
 - Name the type of intermolecular force found in compound **D** and **E**. (1)
 - Write down the IUPAC name of compound **D**. (2)
- 6.3 Fully explain the difference in boiling point between compound **A** and compound **B** in experiment 1. (3)
- 6.4 Draw the structural formula of compound **C**. (2)
- 6.5 From the table above, which compound is a gas at room temperature?
Write only the letter (A – E). (1)
- [12]**

QUESTION 7

The flow diagram below has reactions 1 – 7. Compounds A – G represent organic compounds. Study the flow diagram and answer the questions that follow.



7.1 Write down the **type** of:

7.1.1 reaction 3. (1)

7.1.2 reaction 4. (1)

7.2 Write down the **name/formula** of:

7.2.1 substance X. (1)

7.2.2 inorganic product of reaction 4. (1)

7.3 Compound C and compound G are structural isomers.

Is the above UNDERLINED statement TRUE or FALSE?

Write TRUE or FALSE and give a reason for your answer. (2)

7.4 Write down the IUPAC name of compound D. (2)

- 7.5 Using structural formula write down a balance chemical reaction for reaction 1. (5)
- 7.6 Write down the structural formula of the positional isomer of the organic PRODUCT of reaction 4. (2)
- 7.7 Compound **G** can be converted to compound **B**.
Write down TWO reaction conditions required to convert compound **G** to compound **B**. (2)
- 7.8 Draw the structural formula of the functional isomer of compound **E**. (2)
[19]

TOTAL: 100



Stanmorephysics.com

DATA FOR PHYSICAL SCIENCES GRADE 12 PAPER 1 (PHYSICS)

TABLE 1: PHYSICAL CONSTANTS

NAME	SYMBOL	VALUE
Acceleration due to gravity	g	$9,8 \text{ m}\cdot\text{s}^{-2}$
Universal gravitational constant	G	$6,67 \times 10^{-11} \text{ N}\cdot\text{m}^2\cdot\text{kg}^{-2}$
Speed of light in a vacuum	c	$3,0 \times 10^8 \text{ m}\cdot\text{s}^{-1}$
Planck's constant	h	$6,63 \times 10^{-34} \text{ J}\cdot\text{s}$
Coulomb's constant	k	$9,0 \times 10^9 \text{ N}\cdot\text{m}^2\cdot\text{C}^{-2}$
Charge on electron	e	$1,6 \times 10^{-19} \text{ C}$
Electron mass	m_e	$9,11 \times 10^{-31} \text{ kg}$
Mass of the Earth	M	$5,98 \times 10^{24} \text{ kg}$
Radius of the Earth	R_E	$6,38 \times 10^6 \text{ m}$

TABLE 2: FORMULAE

MOTION

$v_f = v_i + a\Delta t$	$\Delta x = v_i \Delta t + \frac{1}{2}a\Delta t^2$ OR $\Delta y = v_i \Delta t + \frac{1}{2}a\Delta t^2$
$v_f^2 = v_i^2 + 2a\Delta x$ OR $v_f^2 = v_i^2 + 2a\Delta y$	$\Delta x = \left(\frac{v_i + v_f}{2}\right)\Delta t$ OR $\Delta y = \left(\frac{v_i + v_f}{2}\right)\Delta t$

FORCE

$F_{net} = ma$	$p = mv$
$f_s^{max} = \mu_s N$	$f_k = \mu_k N$
$F_{net}\Delta t = \Delta p$ $\Delta p = mv_f - mv_i$	$w = mg$
$F = \frac{Gm_1 m_2}{d^2}$ OR $F = \frac{Gm_1 m_2}{r^2}$	$g = \frac{GM}{d^2}$ OR $g = \frac{GM}{r^2}$

WORK, ENERGY AND POWER/ARBEID, ENERGIE EN DRYWING

$W = F\Delta x \cos \theta$	$U = mgh$ OR $E_p = mgh$
$K = \frac{1}{2}mv^2$ OR $E_k = \frac{1}{2}mv^2$	$W_{net} = \Delta K$ OR $W_{net} = \Delta E_k$ $\Delta K = K_f - K_i$ OR $\Delta E_k = E_{kf} - E_{ki}$
$W_{nc} = \Delta K + \Delta U$ OR $W_{nc} = \Delta E_k + \Delta E_p$	$P = \frac{W}{\Delta t}$
$P_{ave} = Fv_{ave}$	

WAVES, SOUND AND LIGHT

$v = f\lambda$	$T = \frac{1}{f}$
$f_L = \frac{v \pm v_L}{v \pm v_s} f_s$	$E = hf$ OR $E = h\frac{c}{\lambda}$
$E = W_o + E_{k(max)}$ OR $E = W_o + K_{max}$ where	$E = hf$ and $W_0 = hf_0$ and $E_{k(max)} = \frac{1}{2}mv_{max}^2$ OR $K_{max} = \frac{1}{2}mv_{max}^2$

ELECTROSTATICS

$F = \frac{kQ_1 Q_2}{r^2}$	$E = \frac{kQ}{r^2}$
$V = \frac{W}{q}$	$E = \frac{F}{q}$
$n = \frac{Q}{e}$ OR $n = \frac{Q}{q_e}$	

ELECTRIC CIRCUITS

$R = \frac{V}{I}$	$\text{emf } (\varepsilon) = I(R + r)$
$R_s = R_1 + R_2 + \dots$	
$\frac{1}{R_p} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$	$q = I \Delta t$
$W = Vq$	$P = \frac{W}{\Delta t}$
$W = VI\Delta t$	$P = VI$
$W = I^2R\Delta t$	$P = I^2R$
$W = \frac{V^2\Delta t}{R}$	$P = \frac{V^2}{R}$

ALTERNATING CURRENT

$I_{\text{rms}} = \frac{I_{\text{max}}}{\sqrt{2}}$	$P_{\text{ave}} = V_{\text{rms}} I_{\text{rms}}$
$V_{\text{rms}} = \frac{V_{\text{max}}}{\sqrt{2}}$	$P_{\text{ave}} = I_{\text{rms}}^2 R$

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

KEY/SLEUTEL

Atomic number
Atoomgetal

Symbol
Simbool

Electronegativity
Elektronegatiwiteit

Approximate relative atomic mass
Benaderde relatiewe atoommassa

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)	2 (I)									
1 H 1	2,1 Li 7	3 Be 9	4 Na 23	5 Mg 24	6 K 39	7 Ca 40	8 Sc 45	9 Ti 48	10 V 51	11 Cr 52	12 Mn 55	13 Fe 56	14 Co 59	15 Ni 59	16 Cu 63,5	17 Zn 65	18 Ga 70	19 Ge 73	20 As 75	21 Se 79	22 Br 80	23 Kr 84					
0,8 Rb 86	1,0 Sr 88	1,2 Y 89	1,4 Zr 91	1,6 Nb 92	1,8 Mo 96	1,9 Tc 101	2,2 Ru 103	2,2 Rh 106	2,2 Pd 108	1,9 Ag 112	1,7 Cd 115	1,7 In 115	1,8 Sn 119	1,9 Sb 122	2,1 Te 128	2,5 I 127	18 Xe 131	19 At 131	20 Rn 86								
0,7 Cs 133	0,9 Ba 137	0,9 La 139	1,6 Hf 179	1,6 Ta 181	1,6 W 184	1,8 Re 186	1,8 Os 190	1,8 Ir 192	1,8 Pt 195	1,8 Au 197	1,8 Hg 201	1,8 Tl 204	1,8 Pb 207	1,9 Bi 209	2,0 Po 209	2,5 At 209	2,5 Rn 86	21 Fr 226	22 Ra 226	23 Ac 89							
58 Ce 140	59 Pr 141	60 Nd 144	61 Pm 150	62 Sm 152	63 Eu 157	64 Gd 159	65 Tb 163	66 Dy 165	67 Ho 167	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175	72 Th 232	73 Pa 238	74 U 238	75 Np 238	76 Pu 244	77 Am 243	78 Cm 247	79 Bk 247	80 Cf 251	81 Es 252	82 Fm 253	83 Md 253	84 No 253	85 Lr 103



LIMPOPO
PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF
EDUCATION



NATIONAL

SENIOR CERTIFICATE

Stanmorephysics.com

GRADE 12

PHYSICAL SCIENCES/FISIESE WETENSKAPPE

**CONTROL TEST 1 MARKING GUIDELINE/KONTROLE TOETS 1
NASIENRIGLYNE**

17 March/Maart 2025

MARKS/PUNTE : 100

**This marking guideline consists of 12 pages/Hierdie nasienriglyne bestaan uit
12 bladsye**

QUESTION/VRAAG 1





- 1.1 A ✓✓
- 1.2 D✓✓
- 1.3 B ✓✓
- 1.4 B ✓✓
- 1.5 B✓✓
- 1.6 B ✓✓
- 1.7 A ✓✓
- 1.8 C ✓✓
- 1.9 C✓✓
- 1.10 A✓✓



[20]

QUESTION/VRAAG 2

2.1



ACCEPTABLE LABELS/AANVAARDE BYSKRIFTE	NOTES/NOTAS
<p>F_N: Normal/Normaal</p> <p>F_g: w/weight/force due to gravity / gewig/krag agv gravitasie</p> <p>F_A: 10 N/Applied force/Toegepaste krag</p>	<p>ONE mark for each force represented by an arrow with a correct label/Een punt vir elke krag wat deur 'n pyl met 'n korrekte byskrif voorgestel word</p> <p><u>Penalise ONCE for each of the following:</u>/Penaliseer EEN KEER vir elk van die volgende:</p> <ul style="list-style-type: none"> • No arrow/geen pyl • There is no dot/daar is nie 'n kol nie • Gap between the line and the dot/spasie tussen die lyne en die kol • Dotted line is used/stippellyn word gebruik • A force diagram is used/kragtediagram is gebruik

(3)

2.2 $F_{net} = ma$

$F_x - f_k = ma$

$10 \cos 15^\circ - 0,53(3 \times 9,8) - 10 \sin 15^\circ \checkmark = 3a \checkmark$

$a = -1,51712 \text{ m.s}^{-2}$

$v_f^2 = v_i^2 + 2a\Delta x \checkmark$

$0 = v_i^2 + 2(-1,51712)4 \checkmark$

$v = 3,48 \text{ m.s}^{-1} (\text{right}/\text{regs}) \checkmark$

Marking criteria for choosing right as positive/Nasien kritiria as regs as positief gekies word



- 1 mark for/punt vir:
 $0,53(3(9,8) - 10\sin 15)$

- 1 mark for/punt vir:
 $10 \cos 15$ and/en $3a$

- 1 mark for/punt vir:
 $v_f^2 = v_i^2 + 2a\Delta x$

- 1 mark for/punt vir:
 $0 = v_i^2 + 2(-1,51712)4$

- 1 mark for/punt vir:
 $v = 3,48 \text{ m.s}^{-1}$ (right/regs)

(5)




Stanmorephysics.com

2.3 Increase ✓/Toeneem


The resultant force will increase/acceleration will increase causing the velocity to also increase. ✓/ Die resulterende krag sal toeneem/versnelling sal toeneem, wat veroorsaak dat die snelheid ook toeneem. (2)

2.4



Marking criteria for choosing right as positive/Nasien kriteria as regs as positief gekies word

- 1 mark for the line parallel to the x- axis and intersecting with the y axis in the first quadrant/ 1 punt vir die lyn parallel aan die x-as en sny met die y-as in die eerste kwadrant
- 1 mark for the line parallel to the x axis in the fourth quadrant/ 1 punt vir die lyn parallel aan die x-as in die vierde kwadrant



- 1 mark for the time of change in direction of the net force/ 1 punt vir die tyd van verandering in rigting van die netto krag.

Note: Two parallel lines in the fourth and the first quadrant both intersecting with the y axis (2/3)/ Let wel: Twee parallele lyne in die vierde en die eerste kwadrant wat albei met die y-as sny (2/3)

(3)
[13]



Stanmorephysics.com

QUESTION/VRAAG 3


- 3.1 When object A exerts a force on object B, object B simultaneously exert an oppositely directed force of equal magnitude on object A ✓✓
 Wanneer voorwerp A 'n krag op voorwerp B uitoefen, oefen voorwerp B gelyktydig 'n teenoorgestelde gerige krag van gelyke grootte op voorwerp A uit (2 or 0)

- 3.2 Because the direction of resultant force/acceleration is the same as the direction of motion. ✓ I Omdat die rigting van resulterende krag/versnelling dieselfde is as die rigting van beweging. (1)

3.3 $F_{net} \Delta t = mv_f - mv_i$ } ✓
 $F_{net} \Delta t = \Delta p$

$\Delta p = 5(2,5) - 5(1,5)$ ✓

$\Delta p = 5 \text{ N.s right/regs}$ ✓




(3)

3.4 $\sum p_i = \sum p_f$
 $mv_{ix} + mv_{iy} = mv_{fx} + mv_{fy}$ }

$2(3) + 5(1,5) \checkmark = 2v_{fx} + 5(2,5) \checkmark$
 $v_{fx} = 0,5 \text{ m.s}^{-1} \text{ right} \checkmark/\text{regs}$

(4)
[10]

QUESTION/VRAAG 4



4.1 The motion during which the only force acting on an object is the gravitational force. ✓✓ / Die beweging waartydens die enigste krag wat op 'n voorwerp inwerk, die gravitasiekrag is (2 or 0)

4.2 $v_f^2 = v_i^2 + 2g\Delta x$ or $v_f^2 = v_i^2 + 2g\Delta y$ ✓
 $0\checkmark = v_i^2 + 2(-9,8)(7) \checkmark$
 $v_i = 11,71 \text{ m.s}^{-1}$ (upwards/up/)✓/boontoe/op (4)


4.3.1 **Positive marking from 4.2/Positiewe nasien vanaf 4.2**

$$E_k(\text{hi the ground/}) = E_k(\text{that it bounce}) + E_k(\text{converted to other energies})$$

$$\frac{1}{2}mv_{(\text{hi})}^2 = \frac{1}{2}mv_{(\text{bounce})}^2 + \frac{1}{2}mv_{(\text{convert})}^2$$

$$\frac{1}{2}(1,5)v_{(\text{hit})}^2 \checkmark \quad (\text{m.s}^{-1} = \frac{1}{2}(1,5)(11,71)^2 \checkmark + 344,1 \checkmark)$$

$$v_{(\text{hit})} = 24,41 \text{ m.s}^{-1}$$
 (downwards)✓/(afwaarts)



(4)

4.3.2 **Positive marking from 4.3.1/Positiewe nasien vanaf 4.3.1**

$$v_f = v_i + gt \checkmark$$

$$24,41 = 12 + (9,8)t \checkmark$$

$$t = 1,27s \checkmark$$

(3)
[13]

QUESTION/VRAAG 5

5.1.1 A ✓ and/en B✓

(2)

5.1.2 C ✓

(1)

5.2.1 3-chloro✓hexane ✓ / 3-chloroheksaan

Marking criteria/Nasien kriteria

- 1 mark for 3-chloro/1 punt vir 3-chloro
- 1 mark for hexane/1 punt vir heksaan

(2)

5.2.2 4-methyl✓pentan-2-one✓✓ / 4-metielpentaan-2-oon

OR/OF

4-methyl✓-2- pentanone✓✓ / 4-metiel-2-pentanoon


Stanmorephysics.com

Marking criteria/Nasien kriteria


- 1 mark for 4-methyl/1 punt vir 4-metiel
- 1 mark for pentanone/1 punt vir pentaanoon
- 1 mark for whole name correct/1 punt vir die hele naam korrek

(3)

5.3



ACCEPT/AANVAAR



(2)

5.4 C_nH_{2n} ✓

(1)

5.5.1 (Concentrated) Sulphuric acid ✓ (No mark for H_2SO_4)/
(Gekonsentreerde) Swaelsuur (geen punt vir H_2SO_4 nie)

(1)

5.5.2 Endothermic ✓ / Endotermies

(1)

[13]

QUESTION/VRAAG 6

6.1 The temperature at which the vapour pressure of the substance is equal to the atmospheric pressure. ✓✓ / Die temperatuur waarby die dampdruk van die stof gelyk is aan die atmosferiese druk. (2 or 0)

6.2.1 Aldehyde ✓ / Aldehyd (1)

6.2.2 London force/ induced dipole force ✓ / London krag/geïnduseerde dipool krag (1)

6.2.3 methylpropanal✓✓ / metielpropaanal (2)

ACCEPT/AANVAAR


2-methylpropanal/2-metielpropaanal

- 6.3
- Compound A has larger surface area/longer chain length than compound B. ✓ / Verbinding A het 'n groter oppervlakte/langer kettinglengte as verbinding B
 - Compound A has stronger intermolecular force/London force than compound B. ✓ / Verbinding A het sterker intermolekulêre krag/Londen-krag as verbinding B
 - More energy is required to overcome the stronger intermolecular force in compound A than compound B ✓ / Meer energie word benodig om die sterker intermolekulêre krag in verbinding A as verbinding B te oorkom

Or/Of

- Compound B has smaller surface area/smaller chain length than compound A. ✓ / Verbinding B het 'n kleiner oppervlakte/kleiner kettinglengte as verbinding A.
- Compound B has weaker intermolecular force/London force than compound A. ✓ / Verbinding B het swakker intermolekulêre krag/Londen-krag as verbinding A
- Less energy is required to overcome the weaker intermolecular force/London force in compound B than in compound A ✓ / Minder energie word benodig om die swakker intermolekulêre krag/Londen krag in verbinding B te oorkom as in verbinding A

6.4



(2)

6.5

C ✓

(1)
[12]

QUESTION/VRAAG 7

7.1.1 Combustion/oxidation ✓/Verbranding/oksidasie (1)


7.1.2 Substitution ✓/Substitusie (1)

7.2.1 NaOH/KOH/H₂O/Sodium hydroxide/potassium hydroxide/water ✓/
Natriumhidroksied/kaliumhidroksied/water (1)

7.2.2 HCl/ hydrogen chloride ✓/ waterstofchloried (1)
(NO MARK FOR HYDROCHLORIC ACID/GEEN PUNT VIR
SOUTSUUR NIE)


7.3 
 - False✓/Fals
 Compound G and compound C do not have the same molecular mass/ compound G has chlorine while compound C has bromine. ✓/ Verbinding G en verbinding C het nie dieselfde molekuläre massa nie/ verbindung G het chloor terwyl verbindung C broom het (2)

7.4 1-butanol/ butan-1-ol✓✓/ 1-butaanol/butaan-1-ol (2)



Marking criteria/Nasien kriteria

- 2 mark for 8 carbons/2 punte vir 8 koolstowwe
 - 1 mark for double bonds/1 punt vir dubbelbindings
 - 1 mark for whole structure correct/1 punt vir hele struktuur korrek
 - 1 mark for butane/1 punt vir butaan
- (5)



(2 or 0)


7.7 Concentrated strong base/ concentrated NaOH/KOH ✓
/gekonsentreerde sterk basis/gekonsentreerde NaOH/KOH



Heat✓/Hitte

(2)

7.8



- 1 mark for functional group/1 punt vir funksionele groep
- 1 mark for whole structure correct/1 punt vir hele struktuur korrek

(2)
[19]

TOTAL/TOTAAL: 100