



LIMPOPO

PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA



DEPARTMENT OF
EDUCATION

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

Stanmorephysics.com
**MATHEMATICS P1
SEPTEMBER 2025**

MARKS: 150

TIME: 3 hours



EMATHP1

This question paper consists of 9 pages and 1 information sheet.

INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 10 Questions.
2. Answer **ALL** the questions.
3. Number your answers correctly according to the numbering system used in this question paper.
4. Clearly show **ALL** calculations, diagrams and graphs that you have used in determining your answers.
5. Answers only will **NOT** necessarily be awarded full marks.
6. An approved scientific calculator (non-programmable and non-graphical) may be used, unless stated otherwise.
7. If necessary, answers should be rounded off to **TWO** decimal places, unless stated otherwise.
8. Diagrams are **NOT** necessarily drawn to scale.
9. Information sheet with formulae is included at the end of the question paper.
10. Write neatly and legibly.

QUESTION 1

1.1 Solve for x :


$$1.1.1 \quad (3x+1)(x-2) = 0 \quad (2)$$

$$1.1.2 \quad 5x(x-2) = 2 \quad (\text{correct to TWO decimal places}) \quad (4)$$

$$1.1.3 \quad (3x-2)^2 > 3x \quad (4)$$


$$1.1.4 \quad x-7 = \sqrt{x-5} \quad (4)$$

$$1.1.5 \quad 3^{2x} - 12 \cdot 3^x + 27 = 0 \quad (3)$$

1.2 Solve for x and y simultaneously:

$$3^y = 9^{x-3}$$

$$y^2 = 20 - x^2 \quad (6)$$

1.3 Given: $5^n = x$ and $n = \log_2 y$ Express 50^{n+1} in terms of x and y . (4)**[27]**

QUESTION 2

- 2.1 Given: $-4 ; 0 ; 4 ; 8 ; \dots$
- 2.1.1 Write down the next three terms of the sequence. (1)
- 2.1.2 Determine the general term of the sequence. (2)
- 2.1.3 Determine the term of the sequence that is equal to 56. (2)
- 2.1.4 Calculate the value of the 12th term of the sequence. (1)


- 2.2 Given the following quadratic sequence: $x ; y ; 2 ; -2 ; -8 ; -16$
- 2.2.1 Calculate the values of x and y . (5)
- 2.2.2 The n th term of this sequence is $T_n = an^2 + bn + c$. Determine the values of a , b and c . (5)
- [16]

QUESTION 3

- Given: $3(2)^5 + 3(2)^4 + 3(2)^3 + \dots$
- 3.1 Will the series converge? Give a reason for your answer (2)
- 3.2 Calculate the sum of the first seven terms of the series. (2)
- 3.3 Calculate the sum to infinity of the series. (2)
- 3.4 Calculate: $\sum_{k=8}^{\infty} 3(2)^{6-k}$ (2)
- [8]

QUESTION 4

- 4.1 The graphs of $f(x) = -x^2 - 4x + 5$ and $g(x) = mx + c$ are given below. Points E, M and P are the intercepts of the graphs with the axes. N is the turning point of f and NTD is the axes of symmetry of f . T is a point on the graph of g .



- 4.1.1 Determine the coordinates of points E and P. (3)
- 4.1.2 Calculate the value of m and c . (2)
- 4.1.3 Calculate the length of NT. (6)
- 4.1.4 Determine the coordinates of point S, the mirror image of point M with respect to the axis of symmetry. (2)

4.2 Given:

$$f(x) = \frac{x+3}{x+1}$$

4.2.1

Show that f can be written as $f(x) = \frac{2}{x+1} + 1$. (2)

4.2.2

Determine the intercepts of f with the axes. (4)

4.2.3

Write down the equation of the axes of symmetry of f with the negative gradient. (2)

4.2.4

Sketch the graph of f (show the coordinates of the intercepts). (3)

4.2.5

For which value(s) of x will $\frac{2}{x+1} \geq -1$. (2)

[26]

QUESTION 5

Given: $f(x) = 2^{-x}$

5.1 Determine the equation of f^{-1} in the form $y = \dots\dots\dots$ (2)

5.2 Sketch the graphs of f and f^{-1} on the same set of axes (name each graph accordingly). (4)

5.3 Write down the domain of f^{-1} . (1)

[7]

QUESTION 6

Elton buys a car for R375 000. He secures a loan at an interest rate of 8% p.a., compounded monthly. The monthly instalment is R7 000. He pays the first instalment one month after the loan was secured.

6.1 Calculate the effective annual interest rate on the loan. Leave your answer correct to TWO decimal places. (2)

6.2 Calculate the number of months it will take Elton to repay the loan. (5)

6.3 Calculate the value of the final instalment. (4)

6.4 The value of the car depreciates at i % p.a. After 4 years its value is R270 000. Calculate i . (3)

[14]

QUESTION 7

7.1 Determine $f'(x)$ from first principles if $f(x) = x^2 - 3x$. (5)

7.2 Determine:

7.2.1 $D_x[(x+1)(5x+9)]$ (3)


7.2.2 $\frac{dy}{dx}$ if $y = \frac{-5}{\sqrt{x}} - x^5$ (3)

7.2.3 $f'(x)$ if $f(x) = \frac{x^3 + x^2 - 2x}{x-1}$ (4)

[15]

QUESTION 8

The diagram below represents the graph of $f(x) = x^3 + bx^2 + cx + 4$. B is the turning point and there is a relative minimum at C.



Determine:

8.1 The values of b and c . (4)


8.2 The coordinates of the point of inflection. (4)

8.3 The equation of the tangent to the curve at $x = 3$. (4)

[12]

QUESTION 9

In the figure below, $\triangle ABE$ is mounted on a rectangle $BCDE$. $\triangle ABE$ has a base of length x metres. The base and the perpendicular height of the triangle add up to 10 metres. y is the breadth of rectangle $BCDE$. The perimeter of the rectangle is 32m.



- 9.1 Show that the Area of the figure $ABCDE$ is $A(x) = -\frac{x}{2}(3x - 42)m^2$ (4)
- 9.2 Calculate the value of x for which $ABCDE$ has a maximum area. (2)
- 9.3 Calculate the maximum area of $ABCDE$. (2)

[8]

QUESTION 10

10.1 If $P(A) = 0,25$; $P(B) = 0,5$ and $P(A \text{ or } B) = 0,625$

10.1.1 Calculate $P(A \text{ and } B)$ (2)

10.1.2 Are events A and B complementary? Give reasons for your answer. (2)

10.2 In a survey, a group of 100 grade 8 learners were asked if they listen to Jazz or Soul music.

- 48 learners listened to Jazz music
- 52 learners listened to Soul music
- 10 learners listened to neither
- x learners listened to both Jazz and Soul music

10.2.1 Calculate the value of x and represents a survey of this 100 grade 8 learners on a venn diagram. (5)

10.2.2 Determine the probability of selecting a learner that listens to Soul music. (1)

10.2.3 Determine the probability of selecting a learner that listens to Jazz or Soul music. (1)

10.3 Four - digit codes (not beginning with 0), are to be constructed from the following set of digits: $\{1; 2; 3; 4; 6; 7; 8; 0\}$.

10.3.1 How many four – digit codes can be constructed, if repetition of digits is allowed? (2)

10.3.2 How many four – digit codes can be constructed, if repetition of digits is not allowed? (2)

10.3.3 Calculate the probability of randomly constructing a four – digit code which is divisible by 5 if repetition of digits is allowed. (2)

[17]

TOTAL [150]



LIMPOPO

PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF
EDUCATION

**NATIONAL SENIOR CERTIFICATE /
NASIONALE SENIOR SERTIFIKAAT**



GRADE/GRADE 12

MATHEMATICS P1/WISKUNDE V1
MARKING GUIDELINES/NASIENRIGLYNE
SEPTEMBER 2025


MARKS/PUNTE: 150



This marking guidelines consists of 14 pages/ Hierdie nasienriglyne bestaan uit 14 bladsye

NOTE/NOTA:


- If a candidate answers a question TWICE, only mark the FIRST attempt/ *As 'n kandidaat 'n vraag TWEE keer beantwoord, merk slegs die EERSTE poging*
- Consistent accuracy applies in ALL aspects of the marking guidelines/ *Konsekwente akkuraatheid is van toepassing in ALLE aspekte van die nasienriglyne*

QUESTION 1/VRAAG 1

1.1	1.1.1	$(3x+1)(x-2) = 0$ $x = -\frac{1}{3}$ or / of $x = 2$	$\checkmark x = -\frac{1}{3}$ $\checkmark x = 2$	(2)
	1.1.2	$5x(x-2) = 2$ $5x^2 - 10x - 2 = 0$ $x = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(5)(-2)}}{2(5)}$ $x = \frac{10 \pm \sqrt{140}}{10}$ $x = 2,18$ or / of $x = -0,18$	$\checkmark$ standard form/ <i>standaard vorm</i> $\checkmark$ substitution/ <i>vervanging</i> $\checkmark x = 2,18$ $\checkmark x = -0,18$	(4)
	1.1.3	$(3x-2)^2 > 3x$ $9x^2 - 12x + 4 - 3x > 0$ $9x^2 - 15x + 4 > 0$ $(3x-1)(3x-4) > 0$ CV / KW : $x = \frac{1}{3}$ or / of $x = \frac{4}{3}$  $x < \frac{1}{3}$ or / of $x > \frac{4}{3}$	$\checkmark$ standard form/ <i>standaard vorm</i> $\checkmark$ critical values/ <i>kritieke waardes</i> $\checkmark x < \frac{1}{3}$ $\checkmark x > \frac{4}{3}$	(4)

	1.1.4	$x - 7 = \sqrt{x - 5}$ $(x - 7)^2 = x - 5$ $x^2 - 14x + 49 = x - 5$ $x^2 - 15x + 54 = 0$ $(x - 6)(x - 9) = 0$ $x \neq 6$ or / of $x = 9$	✓ squaring/vierkant ✓ standard form/standaard vorm ✓ $x \neq 6$ ✓ $x = 9$	(4)
	1.1.5	$3^{2x} - 12 \cdot 3^x + 27 = 0$ $(3^x - 3)(3^x - 9) = 0$ $3^x = 3$ or $3^x = 3^2$ $x = 1$ or $x = 2$	✓ standard form /standaard vorm ✓ $x = 1$ ✓ $x = 2$	(3)
1.2	$3^y = 9^{x-3}$ $y^2 = 20 - x^2$ $3^y = (3)^{2x-6}$ $y = 2x - 6$ $(2x - 6)^2 = 20 - x^2$ $4x^2 - 24x + 36 = 20 - x^2$ $5x^2 - 24x + 16 = 0$ $(5x - 4)(x - 4) = 0$ $x = \frac{4}{5}$ or / of $x = 4$ $y = \frac{-22}{5}$ or / of $y = 2$		✓ $3^y = (3)^{2x-6}$ ✓ $y = 2x - 6$ ✓ substitution/vervanging ✓ standard form/standaard vorm ✓ x – values/waardes ✓ y – values/waardes	(6)
1.3	$n = \log_2 y$ $y = 2^n$ 50^{n+1} $= 50^n \cdot 50$ $= (5 \times 2 \times 5)^n \cdot 50$ $= 50 \cdot 5^n \cdot 2^n \cdot 5^n$ $= 50x^2 y$		✓ $y = 2^n$ ✓ $50^n \cdot 50$ ✓ $50 \cdot 5^n \cdot 2^n \cdot 5^n$ ✓ answer/antwoord	(4)
				[27]

QUESTION 2/VRAAG 2


2.1	2.1.1	12 ; 16 ; 20	✓ answer/antwoord	(1)
	2.1.2	$a = -4$ $d = 4$ $T_n = -4 + (n-1)4$ $= -4 + 4n - 4$ $= 4n - 8$	✓ $d = 4$ ✓ answer/antwoord	(2)
	2.1.3	$4n - 8 = 56$ $4n = 64$ $n = 16$	✓ equating/gelykstel ✓ answer/antwoord	(2)
	2.1.4	$T_{12} = -4 + 11(4)$ $= 40$	✓ answer/antwoord	(1)
2.2	2.2.1	 $-4 - (2 - y) = -2$ $-4 - 2 + y = -2$ $y = 4$ $2 - y - (y - x) = -2$ $2 - 4 - 4 + x = -2$ $x = 4$	✓ second difference/2e verskil ✓ equating/gelykstel ✓ y – value/waarde ✓ equating/gelykstel ✓ x – values/waarde	(5)
	2.2.2	$2a = -2$ $a = -1$ $3(-1) + b = 0$ $b = 3$ $-1 + 3 + c = 4$ $c = 2$ $T_n = -n^2 + 3n + 2$	✓ value of a /waarde van a ✓ substitution/vervanging ✓ value of b /waarde van b ✓ substitution/vervanging ✓ value of c / waarde van c	(5)
				[16]

QUESTION 3/VRAAG 3


3.1	$3(2)^5 + 3(2)^4 + 3(2)^3 + \dots$ $r = \frac{3(2)^4}{3(2)^5} = \frac{1}{2}$ $-1 < r < 1$ the series converges / reeks konvergeer	✓ value of r / waarde van r ✓ reasoning/redenasie	(2)
3.2	$S_n = \frac{a(1-r^n)}{1-r}$ $S_7 = \frac{3(2)^5 \left[1 - \left(\frac{1}{2} \right)^7 \right]}{1 - \frac{1}{2}}$ $= \frac{381}{2}$ $= 190,5$	✓ substitution/vervanging ✓ answer/antwoord	(2)
3.3	$S_\infty = \frac{a}{1-r}$ $= \frac{3(2)^5}{1 - \frac{1}{2}}$ $= 192$	✓ substitution/vervanging ✓ answer/antwoord	(2)
3.4	$\sum_{k=8}^{\infty} 3(2)^{6-k}$ $= S_\infty - S_7$ $= 192 - 190,5$ $= 1,5$	✓ substitution/vervanging ✓ answer/antwoord	(2)
			[8]

QUESTION 4/VRAAG 4

4.1	4.1.1	x - intercept/ <i>afsnit</i> ($y = 0$) $-x^2 - 4x + 5 = 0$ $x^2 + 4x - 5 = 0$ $(x + 5)(x - 1) = 0$ $x = -5$ or / <i>of</i> $x = 1$ $E(-5; 0)$ $P(1; 0)$	$\checkmark$ equating to 0/ <i>gelykstel aan 0</i> $\checkmark x = -5$ $\checkmark x = 1$	(3)
	4.1.2	$y = mx + c$ $y = mx + 5$ passing through / <i>gaan deur</i> $E(-5; 0)$ $m(-5) + 5 = 0$ $-5m = -5$ $m = 1$ $y = x + 5$	$\checkmark c = 5$ $\checkmark m = 1$	(2)
	4.1.3	$f'(x) = 0$ OR $x = -\frac{b}{2a}$ $-2x - 4 = 0$ $= -\frac{(-4)}{2(-1)}$ $x = -2$ $x = -2$ $f'(-2) = 9$ $y = x + 5$ $= -2 + 5$ $= 3$ $T(-2; 3) \therefore$ $DT = 3$ units $NT = 9 - 3$ $= 6$ units	$\checkmark$ $f'(x) = -2x - 4$ OR $= -\frac{(-4)}{2(-1)}$ $\checkmark x = -2$ $\checkmark f'(-2) = 9$ $\checkmark y = 3$ $\checkmark DT = 3$ units/ <i>eenhede</i> $\checkmark NT = 6$ units/ <i>eenhede</i>	(6)



	4.1.4	$-x^2 - 4x + 5 = 5$ $-x^2 - 4x = 0$ $-x(x + 4) = 0$ $x = 0$ or / of $x = -4$ $S(-4; 5)$	$\checkmark$ equating/gelykstel $\checkmark x = -4$	(2)
4.2	4.2.1	$f(x) = \frac{(x+1)+2}{x+1} = \frac{x+1}{x+1} + \frac{2}{x+1}$ $= \frac{2}{x+1} + 1$	$\checkmark f(x) = \frac{(x+1)+2}{x+1}$ $\checkmark f(x) = \frac{x+1}{x+1} + \frac{2}{x+1}$	(2)
	4.2.2	x - intercept/afsnit : $\frac{x+3}{x+1} = 0$ $\therefore x = -3$ y - intercept/afsnit : $= \frac{0+3}{0+1}$ $\therefore y = 3$	$\checkmark \frac{x+3}{x+1} = 0$ $\checkmark x = -3$ $\checkmark y = \frac{0+3}{0+1}$ $\checkmark y = 3$	(4)
	4.2.3	Axis of symmetry / Simmetrie – as : $y = -(x+1)+1$ $\therefore y = -x$	$\checkmark y = -(x+1)+1$ $\checkmark y = -x$	(2)
	4.2.4		$\checkmark$ shape/vorm $\checkmark$ asymptotes/asimptote $\checkmark$ intercepts/afsnitte	(3)
	4.2.5	$x \leq -3$ or $x > -1$	$\checkmark x \leq -3$ $\checkmark x > -1$	(2)
				[26]

QUESTION 5/VRAAG 5

5.1	$y = 2^{-x}$ $x = 2^{-y}$ $-y = \log_2 x$ $y = -\log_2 x$ or $y = \log_2 x^{-1}$ or $y = \log_2 \frac{1}{x}$	✓ interchange x and y / <i>ruil x en y</i> ✓ equation	(2)
5.2		Graph of f / <i>grafiek van f</i> ✓ shape/vorm ✓ (0 ; 1) Graph of f^{-1} / <i>grafiek van f^{-1}</i> ✓ shape/vorm ✓ (1 ; 0)	(4)
5.3	$x \in [0 ; \infty)$ OR $x > 0$	✓ answer/antwoord	(1)
			[7]

QUESTION 6/VRAAG 6

6.1	$1 + i_{eff} = \left(1 + \frac{i^m}{m}\right)^m$ $1 + i_{eff} = \left(1 + \frac{0,08}{12}\right)^{12}$ $i_{eff} = 0,08299950681$ $i_{eff} = 8,3\%$	✓ substitution/vervang ✓ Rate/koers = 8,3%	(2)
6.2	$P_v = \frac{x[1 - (1+i)^{-n}]}{i}$ $375\,000 = \frac{7\,000 \left[1 - \left(1 + \frac{0,08}{12}\right)^{-n}\right]}{\frac{0,08}{12}}$ $\frac{9}{14} = \left(\frac{151}{150}\right)^{-n}$ $n = -\frac{\log\left(\frac{9}{14}\right)}{\log\left(\frac{151}{150}\right)}$ $n = 66,495\,584\,57$ $n = 66,5 \text{ months}$	✓ substitution/vervang ✓ $\frac{0,08}{12}$ ✓ simplification/ vereenvoudig ✓ correct use of logs/ korrekte gebruik van logs ✓ answer/antwoord	(5)

6.3	 $P_v = \frac{7\,000 \left[1 - \left(1 + \frac{0,08}{12} \right)^{-0,49558457} \right]}{\frac{0,08}{12}} \left(1 + \frac{0,08}{12} \right)$ $= R3\,474,91$ OR $\left[375\,000 - \frac{7\,000 \left[1 - \left(1 + \frac{0,08}{12} \right)^{-66} \right]}{\frac{0,08}{12}} \right] \left(1 + \frac{0,08}{12} \right)^{67}$ $= R3\,474,91$ 	✓ value of n /waarde van n ✓ substitution/vervanging ✓ multiply/ vermenigvuldig $\left(1 + \frac{0,08}{12} \right)$ ✓ answer/antwoord ✓ value of n /waarde van n ✓ substitution/vervanging ✓ multiply/ vermenigvuldig $\left(1 + \frac{0,08}{12} \right)$ ✓ answer/antwoord	(4)
6.4	$A = P(1-i)^n$ $270\,000 = 375\,000(1-i)^4$ $(1-i)^4 = \frac{270\,000}{375\,000}$ $i = 1 - \sqrt[4]{\frac{270\,000}{375\,000}}$ $i = 7,9\%$	✓ substitution/vervanging ✓ simplification/ vereenvoudig ✓ answer/antwoord	(3)
			[14]

QUESTION 7/VRAAG 7

7.1	$f(x) = x^2 - 3x$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 3(x+h) - (x^2 - 3x)}{h}$ $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 3x - 3h - x^2 + 3x}{h}$ $= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 3h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(2x + h - 3)}{h}$ $= \lim_{h \rightarrow 0} \frac{2x + h - 3}{1}$ $= 2x - 3$	$\checkmark f(x+h) = (x+h)^2 - 3(x+h)$ $\checkmark -x^2 + 3x$ $\checkmark$ simplification/vereenvoudig $\checkmark$ common factor/gemeensk faktor $\checkmark$ answer/antwoord	(5)
7.2	7.2.1 $D_x[(x+1)(5x+9)]$ $= D_x[5x^2 + 9x + 5x + 9]$ $= D_x[5x^2 + 14x + 9]$ $= 10x + 14$	$\checkmark$ simplification/vereenvoudig $\checkmark 10x$ $\checkmark 14$	(3)
	7.2.2 $y = \frac{-5}{\sqrt{x}} - x^5$ $= -5x^{-\frac{1}{2}} - x^5$ $\frac{dy}{dx} = \frac{5}{2}x^{-\frac{3}{2}} - 5x^4$	$\checkmark -5x^{-\frac{1}{2}}$ $\checkmark \frac{5}{2}x^{-\frac{3}{2}}$ $\checkmark -5x^4$	(3)
	7.2.3 $f(x) = \frac{x^3 + x^2 - 2x}{x-1}$ $= \frac{x(x^2 + x - 2)}{x-1}$ $= \frac{x(x+2)(x-1)}{x-1}$ $= x^2 + 2x$ $f'(x) = 2x + 2$	$\checkmark$ common factor/gemeensk faktor $\checkmark x^2 + 2x$ $\checkmark 2x$ $\checkmark 2$	(4)
			[15]

QUESTION 8/VRAAG 8


8.1	$f(x) = x^3 + bx^2 + cx + 4$ $f'(0) = 0$ $f'(x) = 3x^2 + 2bx + c$ $f'(0) = c$ $\therefore c = 0$ x - intercept / afsnit ($y = 0$) $A(-1; 0)$ $\therefore x = -1$ $f(x) = x^3 + bx^2 + 0 + 4$ $f(-1) = (-1)^3 + b(-1)^2 + 0 + 4 = 0$ $-1 + b + 4 = 0$ $b = -3$ $f(x) = x^3 - 3x^2 + 4$	$\checkmark f'(x)$ $\checkmark c = 0$ $\checkmark$ substitution/vervangings $\checkmark b = -3$	(4)
8.2	Point of inflection / Buigpunt $f''(x) = 0$ $6x - 6 = 0$ $6x = 6$ $x = 1$ $f(1) = (1)^3 - 3(1)^2 + 4$ $= 2$ $(1; 2)$	$\checkmark f''(x) = 0$ $\checkmark x = 1$ $\checkmark$ substitution/vervangings $\checkmark$ answer/antwoord	(4)

8.3	$m = f'(3) = 3(3)^2 - 6(3)$ $= 9$ $\therefore$ equation of a tangent/vergelyking raaklyn $y = mx + c$ $y = 9x + c$ $f(3) = (3)^3 - 3(3)^2 + 4$ $= 4$ passing through/gaan deur (3; 4) $4 = 9(3) + c$ $c = -23$ $y = 9x - 23$	$\checkmark m = 9$ $\checkmark f(3) = 4$ $\checkmark$ substitution/vervangings $\checkmark c = -23$	(4)
			[12]

QUESTION 9/VRAAG 9

9.1	$x + h = 10$ $h = 10 - x$ $2y + 2x = 32$ $y + x = 16$ $y = 16 - x$ $A = \frac{1}{2}bh + lb$ $A(x) = \frac{1}{2}(x)(10 - x) + (x)(16 - x)$ $= 5x - \frac{x^2}{2} + 16x - x^2$ $= \frac{-3x^2}{2} + 21x$ $= -\frac{x}{2}(3x - 42)$	$\checkmark h = 10 - x$ $\checkmark y = 16 - x$ $\checkmark$ substitution/vervangings $\checkmark \frac{-3x^2}{2} - 21x$	(4)
-----	---	---	-----

QUESTION 10/VRAAG 10

10.1	10.1.1	$P(A \text{ or / of } B) = P(A) + P(B) - P(A \text{ and / en } B)$ $0.625 = 0,25 + 0,5 - P(A \text{ and / en } B)$ $P(A \text{ and / en } B) = 0,125$	✓ substitution/vervanging ✓ $P(A \text{ and / en } B) = 0,125$	(2)
	10.1.2	$P(A \text{ and / en } B) \neq 0$ $P(A \text{ and / en } B)$ are not mutually exclusive/ is nie onderling uitsluitend nie. Therefore event A and B are not complementary/daarom is gebeurtenisse A en B nie komplementêr nie	✓ $P(A \text{ and / en } B) \neq 0$ ✓ reasoning/redenasie	(2)
10.2	10.2.1	$48 - x + 52 - x + 10 + x = 100$ $100 - 2x + 10 + x = 100$ $x = 10$	✓ $100 - 2x + 10 + x = 100$ ✓ $x = 10$	(5)
			✓ 38 ✓ 10 ✓ 42	
	10.2.2	$P(\text{soul}) = \frac{52}{100} = \frac{13}{25} = 0,52$	✓ $\frac{13}{25}$	
	10.2.3	$P(\text{jazz or soul}) = \frac{48}{100} + \frac{52}{100} - \frac{10}{100} = \frac{9}{10}$	✓ $\frac{9}{10}$	(1)
10.3	10.3.1	$7 \times 8 \times 8 \times 8$ $= 3\,584 \text{ ways/maniere}$	✓ $7 \times 8 \times 8 \times 8$ ✓ answer/antwoord	(2)
	10.3.2	$7 \times 7 \times 6 \times 5$ $= 1\,470 \text{ ways / maniere}$	✓ $7 \times 7 \times 6 \times 5$ ✓ answer/antwoord	(2)
	10.3.3	$P(\text{code divide by } 5) = \frac{7 \times 8 \times 8 \times 1}{3\,584} = \frac{448}{3\,584}$ $= \frac{1}{8} = 0,125$	✓ $\frac{7 \times 8 \times 8 \times 1}{3\,584}$ ✓ answer/antwoord	(2)
				[17]

TOTAL/ TOTAAL: 150