



PHYSICAL SCIENCES P2 (CHEMISTRY)

FINAL EXAMINATION

**NOVEMBER 2025** 

Stanmorephysics.com

**GRADE 11** 

**MARKS: 150** 

TIME: 3 hours

This question paper consists of 12 pages and 4 data sheets.

Copyright reserved Please turn over

### INSTRUCTIONS AND INFORMATION

- This question paper consists of EIGHT questions. Answer ALL the questions in the ANSWER BOOK.
- 2. Start EACH question on a NEW page in the ANSWER BOOK.
- Number the answers correctly according to the numbering system used in this
  question paper.
- Leave ONE line between two sub-questions, for example between QUESTION 2.1 and QUESTION 2.2.
- You may use a non-programmable calculator.
- You are advised to use the attached DATA SHEETS.
- Show ALL formulae and substitutions in ALL calculations.
- 8. Round off your final numerical answers to a minimum of TWO decimal places.
- 9. Give brief motivations, discussions et cetera where required.
- 10. Write neatly and legibly.

(2)

(2)

### QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Four options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the correct answer and write only the letter (A–D) next to the question number (1.1–1.10) in the ANSWER BOOK, for example 1.11 E.

| 1.1T | he tendency | of an aton | n to attract | a bonded | pair of | electrons i | is known as . |  |
|------|-------------|------------|--------------|----------|---------|-------------|---------------|--|
|------|-------------|------------|--------------|----------|---------|-------------|---------------|--|

- A polarity.
- B electron affinity.
- C electronegativity.
- D ionisation energy.
- 1.2 Which ONE of the following atomic combinations will have the shortest bond length?
  - A C-C
  - B C-O
  - C C = C

$$D \quad C \equiv C \tag{2}$$

1.3 Which ONE of the following substances has a tetrahedral shape?

inmorephysics.com

- A CO<sub>2</sub>
- B SO<sub>2</sub>
- C HCN
- D NH<sub>4</sub><sup>+</sup> (2)

1.4 In which ONE of the following are ION-DIPOLE forces present?

- A NH<sub>3</sub> gas
- B CO<sub>2</sub> gas
- C LiCl aqueous solution
- D MgCl<sub>2</sub> solid c5.com

Copyright reserved Please turn over

| 1.5 | One mole of a gas, SEALED in a container of volume V, exerts pressure p. |  |  |  |  |  |
|-----|--|--|--|--|--|--|
|     | The volume of the container is decreased to 1/3V, while the temperature  |  |  |  |  |  |
|     | remains constant. What is the pressure now exerted by the gas?           |  |  |  |  |  |

A 1/3 p

B 3 p

C p

D 1,5 p (2)

- 1.6 Consider the following statements about ideal gases:
  - (i) The gas particles occupy no volume.
  - (ii) There are no intermolecular forces between the particles
  - (iii) The collisions between the particles are perfectly elastic

Which of the above statements are true?

- A (i) and (ii) only
- B (i) and (iii) only
- C (ii) and (iii) only

D (i), (ii) and (iii) (2)


- 1.7 Which ONE of the following solutions will have the greatest concentration of H<sup>+</sup> ions?
  - A 2 mol·dm<sup>-3</sup> NH<sub>3</sub>
  - B 1,5 mol·dm<sup>-3</sup> CH₃COOH
  - C 1,5 mol·dm<sup>-3</sup> H<sub>2</sub>SO<sub>4</sub>
  - D 1,5 mol·dm<sup>-3</sup> HCl (2)
- 1.8 Consider the following reaction:

$$HNO_3(aq) + X \rightarrow NO_3^-(aq) + HSO_4^-(aq)$$

In this reaction, X represents...

- A SO<sub>4</sub><sup>2</sup>-acting as a base
- B  $SO_4^{2-}$  acting as an acid.
- C H<sub>2</sub>SO<sub>4</sub> acting as a base.
- D  $H_2SO_4$  acting as an acid. (2)

1.9 The graph below shows the energy changes that take place during a chemical reaction.



Which ONE of the following expressions corresponds to the activation energy of the reverse reaction?

(2)

1.10 Consider the following redox reaction:

$$10C\ell^- + 2MnO_4^- + 16H^+ \rightleftharpoons 2Mn^{2+} + 8H_2O + 5C\ell_2$$

A product of the reduction half reaction is ...

(2) [2 x 10 = 20]

### QUESTION 2 (Start on a new page.)

Consider the following reaction of nitric acid (HNO<sub>3</sub>) with lithium oxide (Li<sub>2</sub>O).

 $2HNO_3$  (aq) + Li<sub>2</sub>O (s) $\rightarrow$  2LiNO<sub>3</sub> (aq) + H<sub>2</sub>O ( $\ell$ )  $\Delta H = -139 \text{ kJ} \cdot \text{mol}^{-1}$ 

- 2.1 Define the term ionic bond. (2)
- 2.2 Show that the bond between Li and O is ionic. (2)
- 2.3 Write down the name of the salt formed in this reaction. (1)
- 2.4 Consider the H<sub>2</sub>O molecule.
  - 2.4.1 (1) Write down the valency of the O atom.
  - 2.4.2 Draw the Lewis structure for the H<sub>2</sub>O molecule. (3)
  - 2.4.3 What is the shape of the H<sub>2</sub>O molecule? (1)
  - 2.4.4 NAME the ion that is formed when H2O forms a dative covalent bond. (1)
- 2.5 The above reaction takes place in a beaker. How will the temperature of the contents of the beaker be affected? Choose from INCREASES. DECREASES or REMAINS THE SAME. Give a reason for the answer. (2)
- (2)2.6 Define bond length.
- The diagram below shows the bonds that form between the atoms in the 2.7 HNO<sub>3</sub> molecule.

Consider the bonds between the following atoms: (a) H - Onmorephysics.com

- (b) N O
- (c) N = 0
- Is the bond length larger in (a) or in (b)? Explain the answer. (3)2.7.1
- Is the bond energy greater in (b) or in (c)? Give a reason for the 2.7.2 (2)answer. [20]

### QUESTION 3 (Start on a new page.)

The following table provides the bond energies between atoms.

| Bond | Energy (kJ·mol <sup>-1</sup> ) |
|------|--------------------------------|
| C≡C  | 839                            |
| C=O  | 804                            |
| 0=0  | 498                            |
| C-C  | 348                            |
| 0-0  | 145                            |
| H-O  | 463                            |
| C-O  | 358                            |
| H-C  | 413                            |


Consider the following reaction:

$$CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(g)$$

- 3.1 Define the term activation energy. (2)
- 3.2 Determine:
  - 3.2.1 The activation energy for the reaction (3)
  - 3.2.2 The heat of reaction in kJ·mol<sup>-1</sup> (4)
- 3.3 How will the value calculated in QUESTION 3.2.2 be affected if a suitable catalyst is used? Choose from INCREASES, DECREASES or REMAINS THE SAME. Give a reason for the answer. (2)
- 3.4 24 g of CH<sub>4</sub> reacts completely with excess O<sub>2</sub> at STP.
  - 3.4.1 Is there a NET ABSORPTION or a NET RELEASE of energy? (1)
  - 3.4.2 Calculate the energy referred to in QUESTION 3.4.1. (3)
  - 3.4.3 If 5,5 moles of gas is present when the reaction is complete, calculate the initial number of moles of O<sub>2</sub> (g) used in the reaction. (5) [201

### QUESTION 4 (Start on a new page.)

The graphs below show the relationship between vapour pressure and temperature for CHCl<sub>3</sub>, CCl<sub>4</sub> and H<sub>2</sub>O respectively. Atmospheric pressure is 101,3 kPa.




| 4.1 | Define the term vapour pressure. |  |            |  |  |  |  |
|-----|----------------------------------|--|------------|--|--|--|--|
| 4.2 | Write de                         | Write down the:  |            |  |  |  |  |
|     | 4.2.1                            | Vapour pressure of CC4 at 70°C   | (1)        |  |  |  |  |
|     | 4.2.2                            | Temperature at which the vapour pressure of CHCl3 is 100 kPa               | (1)        |  |  |  |  |
|     | 4.2.3                            | Phase of CHCl <sub>3</sub> at 65°C   | (1)        |  |  |  |  |
|     | 4.2.4                            | Boiling point of CC4   | (2)        |  |  |  |  |
| 4.3 | Fully ex                         | plain the difference in the boiling points of CC4 and H2O.                 | (4)        |  |  |  |  |
| 4.4 |                                  | soluble in H <sub>2</sub> O? Choose from YES or NO. reason for the answer. | (3)<br>[14 |  |  |  |  |

Copyright reserved Please turn over

### QUESTION 5 (Start on a new page.)

An experiment was conducted to investigate the relationship between pressure and volume of a fixed mass of gas at a temperature of 273 K.

The results obtained are shown in the graph below.




5.1 Named and state the law that was investigated. (3)

5.2 Calculate the volume of the gas when the pressure exerted by the gas is 101,3 kPa. (4)

5.3 Do real gases deviate from ideal gases at LOW pressures or HIGH pressures? Explain the answer. (3)

5.4 Draw a sketch graph of pressure (y-axis) vs volume (x-axis) for this experiment. No values or labels are required on the axes. (2)

The experiment is repeated using a greater mass of gas at 273 K. The results obtained are plotted on the same system of axes.



5.5 Which graph, A or B, will be obtained for the greater mass? Explain the answer.


[15]

(3)

### QUESTION 6 (Start on a new page.)

- 6.1 A 1,50 g sample of a compound containing carbon, hydrogen and oxygen was completely burned in air. The products formed were 1,74 g of carbon dioxide and 0,70 g of water.
  - 6.1.1 Define the term *empirical formula*. (2)
  - 6.1.2 Determine, by calculation, the empirical formula of the compound. (7)
  - 6.1.3 If the ratio of empirical formula to molecular formula is 1:3, determine the molecular formula of the compound (2)
- 6.2 An iron ore (impure sample) contains 30% Fe<sub>2</sub>O<sub>3</sub>.
  - 6.2.1 Write down the oxidation number of iron in Fe<sub>2</sub>O<sub>3</sub> (1)
  - 6.2.1 Calculate the mass of Fe that can be obtained from 150 g of this ore. (4)
- 6.3 In an experiment, a learner reacts 1,51 dm³ of sulphuric acid, H<sub>2</sub>SO<sub>4</sub>, of concentration 0,43 mol·dm⁻³ with solid lithium. The balanced equation for the reaction is:

The graph below shows how the mass of lithium changes with time during the reaction:



- 6.3.1 Define the term *limiting reagent*. (2)
- 6.3.2 Identify the limiting reagent in this reaction. Give a reason for the answer. (2)
- 6.3.3 Calculate the value of Y as shown in the graph. (7)

(5)

### QUESTION 7 (Start on a new page.)

7.1 Acetic acid (CH<sub>3</sub>COOH) ionises in water according to the following balanced equation:

$$CH_3COOH(\ell) + H_2O(\ell) \rightleftharpoons H_3O^+(aq) + CH_3COO^-(aq)$$

- 7.1.1 Define an acid according to the Arrhenius theory (2)
- 7.1.2 Write down the FORMULAE of TWO acids in this reaction. (2)
- 7.1.2 Is acetic acid a STRONG acid or a WEAK acid? Give a reason for the answer. (2)
- 7.2 A nitric acid solution, HNO<sub>3</sub>(aq), has a concentration of 0,15 mol·dm<sup>-3</sup>. In a reaction, 31,3 cm<sup>3</sup> of the HNO<sub>3</sub> solution neutralises 24,5 cm<sup>3</sup> of a calcium hydroxide solution, Ca(OH)<sub>2</sub>(aq), according to the following balanced equation:

$$2HNO_3(aq) + Ca(OH)_2(aq) \rightarrow Ca(NO_3)_2(aq) + 2H_2O(l)$$

7.2.1 Calculate the concentration of the Ca(OH)<sub>2</sub> solution.

250 cm<sup>3</sup> of the Nitric acid HNO<sub>3</sub> solution, with the same concentration of 0,15 mol·dm<sup>-3</sup> is now reacted with pure Zn.

$$Zn(s) + 2HNO3(aq) \rightarrow Zn(NO3)2(aq) + H2(g)$$

The pH of the final solution is 2,174.
Assume that the volume of the solution does not change.

- 7.2.2 Calculate the mass of Zn that reacted. (8)
- 7.2.3 Determine the volume that the hydrogen gas produced will occupy at 25°C.

  Take molar gas volume as 24000cm<sup>3</sup>·mol<sup>-1</sup> at 25°C

ake molar gas volume as 24000cm<sup>3</sup>·mol<sup>-1</sup> at 25°C (3) [22]

### QUESTION 8 (Start on a new page.)

8.1 A solution of iron (II) nitrate reacts with At (s) according to the following unbalanced ionic equation:

$$Al(s) + Fe^{2+}(aq) \rightleftharpoons Al^{3+}(aq) + Fe(s)$$

- 8.1.1 Define the term redox reaction. (2)
- 8.1.2 Identify the reducing agent in the above reaction. (1)
- 8.1.3 Give a reason why the nitrate ion, NO<sub>3</sub>-, is a spectator ion in the reaction. (1)
- 8.1.4 Give a reason why the reaction is unbalanced. (2)
- 8.2 The incomplete ionic reaction between dichromate ions ,Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup>, and tin (II) ions, Sn<sup>2+</sup>, in an acid medium is given below.

- 8.2.1 Write down the oxidation number of Cr in the Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> ion. (1)
- 8.2.2 Write down the reduction half-reaction. (2)
- 8.2.3 Using the Table of Standard Reduction Potentials, write down the balanced net ionic equation. (3)

  [12]

TOTAL:150

### **DATA FOR PHYSICAL SCIENCES GRADE 11** PAPER 2 (CHEMISTRY) GEGEWENS VIR FISIESÈ WETENSKAPPE GRAAD 11 **VRAESTEL 2 (CHEMIE)**

### TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESE KONSTANTES

| NAME/NAAM                                       | SYMBOL/SIMBOOL | VALUE/WAARDE                              |
|---|----------------|---|
| Standard pressure<br>Standaarddruk              | pθ             | 1,013 x 10 <sup>5</sup> Pa                |
| Molar gas volume at STP Molêre gasvolume by STD | Vm             | 22,4 dm <sup>3</sup> ·mol <sup>-1</sup>   |
| Standard temperature Standaardtemperatuur       | Тө             | 273 K                                     |
| Charge on electron<br>Lading op elektron        | е              | 1,6 x 10 <sup>-19</sup> C                 |
| Avogadro's constant  Avogadro-konstante         | NA             | 6,02 x 10 <sup>23</sup> mol <sup>-1</sup> |

#### TABLE 2: FORMULAE/TABEL 2: FORMULES

| $n=\frac{m}{M}$   | $n = \frac{N}{N_A}$                           |  |
|---|---|--|
| $c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$                    | $n = \frac{V}{V_m}$                           |  |
| $\frac{c_a v_a}{c_b v_b} = \frac{n_a}{n_b}$                   | pH = -log[H <sub>3</sub> O <sup>+</sup> ]     |  |
| P <sub>1</sub> V <sub>1</sub> = P <sub>2</sub> V <sub>2</sub> | PV = nRT                                      |  |
| K <sub>w</sub> = [H₃O <sup>+</sup> ][OH <sup>-</sup> ]        | ] = 1 x 10 <sup>-14</sup> at/ <i>by</i> 298 K |  |

|  |   | _        |
|--|---|----------|
|  | 3 | <u>~</u> |

NSC

Physical Sciences/P2

November 2025

|                   |             |  | -   |                    | -   |                   |                     | _ |
|-------------------|-------------|--|---|--------------------|---|-------------------|---------------------|---|
| Down              | n Dec       | 10<br>200<br>200<br>200<br>200<br>200<br>200<br>200<br>200<br>200<br>2 | 18<br>A   | W. Z. Z.           | R X E   | S SCOL            | E 3                 |   |
| 34                | 3           | 4.0<br>4.0<br>19   | તુ 17<br>લ CI<br>35,5   | 8 E &              | 25<br>127 - 53  | 25<br>A A         | 75<br>75            |   |
| <b>3</b> 2        |             | 8<br>မှ 0<br>က 16  | 16<br>เพื่ร<br>32   | 24 34<br>28 53     | 52<br>N Te<br>128   | 02<br>28 8        | 69<br>T             |   |
| 25                |             | 7<br>σ, N<br>14  | 15<br>N P   | 33<br>N As<br>75   | 8. 9. 5.<br>12. 12.   |                   | 88 P                |   |
| <b>4</b> §        |             |  |   |                    | 8.1<br>2.2.5<br>119   |                   | 67                  |   |
| 13                |             |  |   |                    | 49<br>115<br>115  |                   | 99 A                |   |
| 12                |             |  |   |                    | 17 Cd 112 Ld 48 L |                   | 8 E                 |   |
| 7                 |             | i  |   |                    | 1.9<br>108  | 79<br>197         | 64<br>64            |   |
| 10                |             | lood<br>bood   | mass  | - CONTRACTOR       | 2.2<br>106 P 46   | 78<br>195<br>195  | £ 5                 |   |
| 9<br>mber         | etal        | Simbool  | ative atomic mass<br>ewe atoommassa                               |                    | 2.2<br>103 E &  | 77<br>11<br>192   | 29 ES               |   |
| 8 9<br>omic numbe | Ato omgetal | 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2                               | elative   | AL                 | 22<br>5   | 76<br>OS<br>190   | P 4                 |   |
| 7<br>Atom         | ` _         | <b>†</b>   | Approximate relative atomic mass<br>Benaderde relatiewe atommassa | 25<br>1. Min<br>55 |   | 75<br>Re<br>186   | 9 P                 | - |
| 9                 | WE          | negativ  | Approx<br>Benad   | 4.6<br>7.5<br>52   |   | 74<br>W<br>184    | 88 2                |   |
| S                 | KEYISLEUTEI | Electronegativity –<br>Elektronegatiwiteit                             |   |                    | 41<br>Nb<br>92  | 73<br>Ta<br>181   | 88 8                |   |
| 4                 | ¥           |  |   | 2,r<br>2 ≒ &       | 40<br>7. Zr<br>91   | 72<br>© Hf<br>179 |                     |   |
| ဗ                 |             |  |   | ε,ι<br>2 % &       |   |                   | 89<br>. Ac          |   |
| 2 🗐               |             | 4 Be   | 12<br>24  | 20<br>Ca<br>40     | 8 S 8   |                   | 9 88<br>6 Ra<br>226 |   |
|                   |             | S'l  | Z'l   | 0°L                | 0'1   | 6'0               |                     |   |
| -€                | - = -       |  |   | 5 × 8              |   | 1000              | 87<br>F F           |   |
|                   | ı'z         | 0,1  | 6'0   | 8,0                | 8,0   | 7,0               | ۷'0                 |   |

| 7  | 3  | 175       | 103<br>L       |
|----|----|-----------|----------------|
| 2  | £  | 173       | 102<br>₹       |
| 69 | Ę  | 169       | 101<br>Md      |
| 8  | Ē  | 167       | 100<br>Fm      |
| 29 | 오  | 165       | 99<br>Es       |
| 99 | Dy | <b>18</b> | C 88           |
| 8  | 9  | 159       | 97<br>BK       |
| 4  | P9 | 157       | 96<br>Cm       |
| 63 | E  | 152       | 95<br>Am       |
| 62 | ES | 150       | \$ ₹           |
| 61 | Pa |           | 93<br>Np       |
| 99 | PN | 144       | 92<br>U<br>238 |
| 29 | Ь  | 141       | Pa<br>Pa       |
| 9  | e, | 40        | 97 28          |

# TABLE 4A: STANDARD REDUCTION POTENTIALS TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE



| Half-reactions/H  | Ε <sup>θ</sup> (V) |   |        |
|---|--------------------|---|--------|
| F <sub>2</sub> (g) + 2e <sup>-</sup>  |                    |   | + 2,87 |
| Co3+ + e-   | ==                 | Co <sup>2+</sup>                        | + 1,81 |
| H <sub>2</sub> O <sub>2</sub> + 2H <sup>+</sup> +2e <sup>-</sup>                  | ===                | 2H₂O                                    | +1,77  |
| MnO 4 + 8H+ + 5e-   | -                  | Mn <sup>2+</sup> + 4H <sub>2</sub> O    | + 1,51 |
| Cl <sub>2</sub> (g) + 2e <sup>-</sup>   |                    |   | + 1,36 |
| Cr <sub>2</sub> O <sup>2-</sup> <sub>7</sub> + 14H <sup>+</sup> + 6e <sup>-</sup> | =                  | 2Cr3+ + 7H₂O                            | + 1,33 |
| O <sub>2</sub> (g) + 4H <sup>+</sup> + 4e <sup>-</sup>                            |                    |   | + 1,23 |
| MnO <sub>2</sub> + 4H <sup>+</sup> + 2e <sup>-</sup>                              | =                  | Mn <sup>2+</sup> + 2H <sub>2</sub> O    | + 1,23 |
| Pt2+ + 2e-  |                    |   | + 1,20 |
| Br <sub>2</sub> (t) + 2e-   |                    |   | + 1,07 |
| NO 3 + 4H+ + 3e-  |                    |   | + 0,96 |
| Hg <sup>2+</sup> + 2e <sup>-</sup>  |                    |   | + 0,85 |
| Ag* + e-  |                    |   | + 0,80 |
| NO 3 + 2H+ + e-   |                    |   | + 0,80 |
| Fe <sup>3+</sup> + e <sup>-</sup>   |                    |   | + 0,77 |
| O <sub>2</sub> (g) + 2H <sup>+</sup> + 2e <sup>-</sup>                            |                    |   | + 0,68 |
| l <sub>2</sub> + 2e   |                    | 21-                                     | + 0,54 |
| Cu* + e-  |                    | 0                                       | + 0,52 |
| SO <sub>2</sub> + 4H <sup>+</sup> + 4e <sup>-</sup>                               | ==                 | 0 . 011 0                               | + 0,45 |
| 2H <sub>2</sub> O + O <sub>2</sub> + 4e <sup>-</sup>                              | -                  | 40H                                     | + 0,40 |
| Cu2+ + 2e-  | =                  | Cu                                      | + 0,34 |
| 3   |                    | SO <sub>2</sub> (g) + 2H <sub>2</sub> O | + 0,17 |
| Stanmore Cu2+++   | om                 | Cu*                                     | + 0,16 |
| Sn <sup>4+</sup> + 2e <sup>-</sup>  |                    | Sn <sup>2+</sup>                        | + 0,15 |
| S + 2H+ + 2e-   |                    | H₂S(g)                                  | + 0,14 |
| 2H+ + 2e-   |                    | H₂(g)                                   | 0,00   |
| Fe <sup>3+</sup> + 3e <sup>-</sup>  |                    |   | - 0,06 |
| Pb2+ + 2e-  | -                  | Pb                                      | -0,13  |
| Sn <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Sn                                      | - 0,14 |
| Ni <sup>2+</sup> + 2e <sup>-</sup>  | -                  | Ni                                      | - 0,27 |
| Co2+ + 2e-  | =                  | Co                                      | - 0,28 |
| Cd2+ + 2e-  | =                  | Cd                                      | - 0,40 |
| Cr3+ + e-   | =                  | Cr <sup>2+</sup>                        | - 0,41 |
| Fe <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Fe                                      | - 0,44 |
| Cr3+ + 3e-  | -                  | Cr                                      | - 0,74 |
| Zn2+ + 2e-  | =                  | Zn                                      | - 0,76 |
| 2H <sub>2</sub> O + 2e <sup>-</sup>   | **                 | H <sub>2</sub> (g) + 2OH                | - 0,83 |
| Cr2+ + 2e-  | =                  | Cr                                      | - 0,91 |
| Mn <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Mn                                      | - 1,18 |
| At3+ + 3e-  | =                  | Al                                      | - 1,66 |
| Mg <sup>2+</sup> + 2e <sup>-</sup>  | quit.              | Mg                                      | - 2,36 |
| Na+ + e-  | =                  | Na                                      | - 2,71 |
| Ca <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Ca                                      | - 2,87 |
| Sr <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Sr                                      | - 2,89 |
| Ba <sup>2+</sup> + 2e <sup>-</sup>  | =                  | Ва                                      | - 2,90 |
| Cs* + e*  | -                  | Cs                                      | - 2,92 |
| K+ + e-   | =                  | К                                       | - 2,93 |
|   |                    |   |        |

Increasing reducing ability/Toenemende reduserende vermoë

Increasing oxidising ability/Toenemende oksiderende vermoë

### **TABLE 4B: STANDARD REDUCTION POTENTIALS** TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE



Increasing oxidising ability/Toenemende oksiderende vermoë

| RD-REDUKSIEPOTE   | Eθ                   |   |        |
|---|----------------------|---|--------|
| Half-reactions/Half   | (v)                  |   |        |
| Li+ + e-  | =                    | Li                                      | - 3,05 |
| K⁺ + e⁻   | =                    | K                                       | - 2,93 |
| Cs+ + e-  | =                    | Cs                                      | - 2,92 |
| Ba <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Ва                                      | - 2,90 |
| Sr <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Sr                                      | - 2,89 |
| Ca <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Ca                                      | - 2,87 |
| Na+ + e-  |                      | Na                                      | - 2,71 |
| Mg <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Mg                                      | - 2,36 |
| Al <sup>3+</sup> + 3e <sup>-</sup>  | $\rightleftharpoons$ | Al                                      | - 1,66 |
| Mn <sup>2+</sup> + 2e <sup>-</sup>  |                      | Mn                                      | - 1,18 |
| Cr2+ + 2e-  | =                    | Cr                                      | - 0,91 |
| 2H <sub>2</sub> O + 2e <sup>-</sup>   |                      | H <sub>2</sub> (g) + 2OH <sup>-</sup>   | - 0,83 |
| Zn <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Zn                                      | - 0,76 |
| Cr3+ + 3e-  |                      | Cr                                      | - 0,74 |
| Fe <sup>2+</sup> + 2e <sup>-</sup>  |                      | Fe                                      | - 0,44 |
| Cr3+ + e-   | 27.10                | Cr <sup>2+</sup>                        | - 0,41 |
| Cd <sup>2+</sup> + 2e <sup>-</sup>  |                      | Cd                                      | - 0,40 |
| Co <sup>2+</sup> + 2e <sup>-</sup>  |                      | Co                                      | - 0,28 |
| Ni <sup>2+</sup> + 2e <sup>-</sup>  |                      | Ni                                      | - 0,27 |
| Sn <sup>2+</sup> + 2e <sup>-</sup>  |                      | Sn                                      | - 0,14 |
| Pb <sup>2+</sup> + 2e <sup>-</sup>  |                      | Pb                                      | - 0,13 |
| Fe <sup>3+</sup> + 3e <sup>-</sup>  |                      | Fe                                      | - 0,06 |
| 2H+ + 2e-   |                      | H <sub>2</sub> (g)                      | 0,00   |
| S + 2H1 + 2e  |                      |   | + 0,14 |
| Sn <sup>4+</sup> + 2e <sup>-</sup>  |                      | Sn <sup>2+</sup>                        | + 0,15 |
| Cu <sup>2+</sup> + e <sup>-</sup>   | =                    | Cu <sup>+</sup>                         | + 0,16 |
| SO <sub>4</sub> <sup>2-</sup> + 4H <sup>+</sup> + 2e <sup>-</sup>                 | =                    | SO <sub>2</sub> (g) + 2H <sub>2</sub> O | + 0,17 |
| Cu <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Cu                                      | + 0,34 |
| 2H <sub>2</sub> O + O <sub>2</sub> + 4e <sup>-</sup>                              |                      | 40H-                                    | + 0,40 |
| SO <sub>2</sub> + 4H <sup>+</sup> + 4e <sup>-</sup>                               | =                    |   | + 0,45 |
| Cu+ + e-  | =                    | Cu                                      | + 0,52 |
| l <sub>2</sub> + 2e <sup>-</sup>  |                      | 21-                                     | + 0,54 |
| O <sub>2</sub> (g) + 2H <sup>+</sup> + 2e <sup>-</sup>                            |                      |   | + 0,68 |
| Fe <sup>3+</sup> + e <sup>-</sup>   | =                    | Fe <sup>2+</sup>                        | + 0,77 |
| NO 3 + 2H+ + e-   |                      | $NO_2(g) + H_2O$                        | + 0,80 |
| Ag+ + e-  | $\rightleftharpoons$ | Ag                                      | + 0,80 |
| Hg <sup>2+</sup> + 2e <sup>-</sup>  | =                    | Hg(ℓ)                                   | + 0,85 |
| NO 3 + 4H+ + 3e-  | =                    | NO(g) + 2H <sub>2</sub> O               | + 0,96 |
| Br <sub>2</sub> (ℓ) + 2e <sup>-</sup>   | =                    | 2Br                                     | + 1,07 |
| Pt <sup>2+</sup> + 2 e <sup>-</sup>   |                      | Pt                                      | + 1,20 |
| MnO <sub>2</sub> + 4H <sup>+</sup> + 2e <sup>-</sup>                              | =                    | Mn <sup>2+</sup> + 2H <sub>2</sub> O    | + 1,23 |
| O <sub>2</sub> (g) + 4H <sup>+</sup> + 4e <sup>-</sup>                            | =                    | 2H <sub>2</sub> O                       | + 1,23 |
| Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup> + 14H <sup>+</sup> + 6e <sup>-</sup> | =                    | 2Cr3+ + 7H2O                            | + 1,33 |
| Cl <sub>2</sub> (g) + 2e <sup>-</sup>   | =                    | 2Ct                                     | + 1,36 |
| MnO - + 8H+ + 5e-   | =                    | Mn <sup>2+</sup> + 4H <sub>2</sub> O    | + 1,51 |
| H <sub>2</sub> O <sub>2</sub> + 2H <sup>+</sup> +2 e <sup>-</sup>                 | =                    | 2H₂O                                    | +1,77  |
| Co3+ + e-   |                      |   | + 1,81 |
|   | =                    | 2F-                                     | + 2,87 |

Increasing reducing ability/Toenemende reduserende vermoë



### NATIONAL SENIOR CERTIFICATE

PHYSICAL SCIENCES P2 (CHEMISTRY)

FINAL EXAMINATION

**NOVEMBER 2025** 

comprehistics com

tanmorephysics.com

**GRADE 11** 

These marking guidelines consist of 10 pages.

# Physical decided from Stanmore physics. COMWAZULU-NATAL/NOVEMBER 2025 Marking Guidelines

### **QUESTION 1**

| 1.1 | CVY            | (2) |
|-----|----------------|-----|
|     | All the second | \_/ |

1.3 
$$\mathsf{D}\checkmark\checkmark$$

$$1.4 \qquad C \checkmark \checkmark \tag{2}$$


tanmorephysics.com

(2)

### Physical desired from Stanmorephysics.com/AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

#### **QUESTION 2**

| 2.1.  | The force of attraction between oppositely charged ions, ✓ formed as a result of the transfer of electrons. ✓   | (2) |
|-------|---|-----|
| 2.2   | $\Delta$ EN = 3.5 $-$ 1<br>= 2.5<br>$\Delta$ EN > 2.1 $\checkmark$  | (2) |
| 2.3   | lithium nitrate ✓   | (1) |
| 2.4.1 | 2 ✓ (Accept: -2)  | (1) |
| 2.4.2 | Marking criteria/Nasienriglyne  ✓ Correct number of electrons surrounding each atom.  ✓ Single bond between O and each of the two H atoms.  ✓ Whole structure correct | (3) |



## Physical decided from Stanmore physics. com AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

### **QUESTION 3**

3.1 The minimum energy needed for a reaction to take place. 
$$\checkmark\checkmark$$
 (2)

3.2.1 E<sub>absorbed</sub> = 
$$(4 \times C - H) + (2 \times O = O)$$
  
=  $(4 \times 413) \checkmark + (2 \times 498) \checkmark$   
=  $2648 \text{ kJ} \cdot \text{mol}^{-1} \checkmark$ 

(3)

### 3.2.2 **POSITIVE MARKING FROM Q3.2.1**:

Ereleased = 
$$(2 \times C = O) + (4 \times O - H)$$
  
=  $(2 \times 804) \checkmark + (4 \times 463) \checkmark$   
=  $3460 \text{ kJ} \cdot \text{mol}^{-1}$ 

$$\Delta H = \sum E_{absorbed} - \sum E_{released}$$

$$= 2648 - 3460 \checkmark$$

$$= -812 \text{ kJ·mol}^{-1} \checkmark$$
(4)

3.3 REMAINS THE SAME <

A catalyst has no effect on energy of reactants or products 

OR A catalyst only affects activation energy.

(2)

(1)

### 3.4.2 POSITIVE MARKING FROM Q3.2.2

$$n = \frac{m}{M}$$

$$= \frac{24}{16} \checkmark$$

$$= 1,5 \text{ mol}$$
Energy released = 1,5 x 812 kJ  $\checkmark$  = 1218 kJ  $\checkmark$ 

(3)

#### 3.4.3 **POSITIVE MARKING FROM Q3.4.2**

no<sub>2</sub> that reacted = 2 x n<sub>CH4</sub> that reacted = 2 x 1,5 = 3 mol√

nco2 that formed = 1 x ncH4 that reacted = 1 x 1,5 = 1,5 mol√

n<sub>H2O</sub> that formed = 2 x n<sub>CH4</sub> that reacted = 2 x 1,5 = 3 mol√

 $n_{02}$  in excess = 5,5 - (1,5 + 3) = 1 mol $\checkmark$ 

Initial 
$$n_{02} = 3 + 1 = 4 \text{ mol}\sqrt{5}$$
 [50]

### Physical decided from Stanmore physics.com/AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

#### **QUESTION 4**

4.1 The temperature at which the vapour pressure of a liquid equals atmospheric pressure. ✓✓ (2)

4.2.1 80 kPa ✓ (1)

4.2.2 60°C ✓ [Accept answers between 60°C and 62°C] (1)

4.2.3 Gas ✓ (1)

4.2.4 75°C√√ (2)

4.3 H<sub>2</sub>O has a higher boiling point than CCl<sub>4</sub>√ Intermolecular forces between H<sub>2</sub>O molecules is hydrogen bonding, while intermolecular forces between CCl<sub>4</sub> molecules is London/Dispersion forces.√ Intermolecular forces in H<sub>2</sub>O are stronger / Intermolecular forces in CCl<sub>4</sub> are weaker. √

More energy is required to overcome the intermolecular forces in  $H_2O$ .  $\checkmark$  (4)

4.4 No. ✓

OPTION 1: CCl₄ molecules are polar ✓ while H₂O molecules are non-polar ✓

OPTION 2: the imtermolecular forces in H<sub>2</sub>O are stronger than in CCl<sub>4</sub> ✓ so they are not of comparable strength. ✓

(3) **[14]** 

(3)

#### **QUESTION 5**

5.1. Marking guidelines

If any of the underlined key words/phrases are omitted: minus 1 mark.

Boyle's Law ✓

<u>Pressure</u> of an enclosed gas is <u>inversely proportional to the volume</u> it occupies at <u>constant temperature</u>. ✓

5.2 **OPTION 1**:

$$V = \frac{1}{5.5 \times 10^{-3}}$$

$$p_1V_1 = p_2V_2 \checkmark$$

$$(115.5)(181.82) = (101.3)V_2 \checkmark$$

$$V_2 = 207.31 \text{ dm}^3 \checkmark$$

**OPTION 2:** 

$$\frac{\frac{\text{OPTION 2.}}{115 - 101,3\checkmark}}{5,5 - \frac{1}{\sqrt{}}\checkmark} = \frac{115}{5,5}\checkmark$$

$$V = 207,45 \text{ dm}^{3}\checkmark$$
(4)

# Physica are lossed from Stanmore physics.com AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

5.3 High√


The volume of the gas particles is no longer negligible  $\checkmark$  **OR** Intermolecular forces become significant.

Increasing the pressure does not decrease the volume as predicted by

Boyle's Law√

(3)

5.4



Marking criteria

Correct shape (hyperbola) ✓✓

5.5 A ✓

The gradient is proportional to the number of moles.

A has a larger gradient, ✓ and hence a larger number of moles of gas ✓.

(3) **[15]** 

(2)

#### **QUESTION 6**

6.1.1 The smallest whole number ratio of elements in a compound. ✓✓ (2 OR 0) (2)

6.1.2 Mass of C =  $\frac{12}{44}$  x 1,74  $\checkmark$  = 0,475 g

Mass of H =  $\frac{2}{18}$  x 0,70  $\checkmark$  = 0,078 g

Mass of O = 1.5 - (0.475 + 0.078)  $\checkmark$  = 0.947 g

C: H: 
$$n = \frac{m}{M}$$
  $n = \frac{m}{M}$   $= \frac{0,475}{12}$  porephysics  $= \frac{0,078}{1}$ 

O:  

$$n = \frac{m}{M}$$

$$= \frac{0,475}{12}$$
 
$$= \frac{0,947}{16}$$
 
$$= 0.04$$
 
$$= 0.078$$
 
$$= \frac{0,947}{16}$$
 
$$= 0.059$$

C:H:O

0,04:0,078:0.059

2:4:3

Molecular formula: C<sub>2</sub>H<sub>4</sub>O<sub>3</sub> ✓ (7)

6.1.3 (C<sub>2</sub>H<sub>4</sub>O<sub>3</sub>)<sub>3</sub>√

Molecular formula is C<sub>6</sub>H<sub>12</sub>O<sub>9</sub> ✓ (2)

# Physical decided from Stanmorephysics.com/AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

6.2.2 Mass of Fe<sub>2</sub>O<sub>3</sub> =  $\frac{30}{100}$  x 150  $\checkmark$  = 45g

Mass of Fe = 
$$\frac{112}{160} \checkmark x 45 \checkmark$$
  
= 31,5 g  $\checkmark$  (4)

- 6.3.1 Limiting reagent is the reactant that is used up completely in a reaction. 

  OR Limiting reagent determines the maximum amount of product formed in a reaction.
- 6.3.2 H<sub>2</sub>SO<sub>4</sub> ✓
  Lithium was left over after the reaction OR All the H<sub>2</sub>SO<sub>4</sub> was used up during the reaction. ✓
- 6.3.3 Marking criteria:
  - (a) Use of formula  $c = \frac{n}{V} \checkmark$
  - (b) Substitution into formula  $c = \frac{n}{V}$  to calculate no fo mol of H<sub>2</sub>SO<sub>4</sub> $\checkmark$
  - (c) Apply ratio (2:1) to calculate no of mol of Li used. ✓
  - (d) Substitution into formula  $n = \frac{m}{M}$  to calculate mass of Li used / initial mass of Li.  $\checkmark$
  - (e) Addition of mass of Li used to final mass of Li OR Addition of moles of Li used to final moles of Li.
  - (f) Final answer. ✓

$$c = \frac{n}{V} \checkmark^{(a)}$$

$$0,43 = \frac{n}{1,51} \checkmark^{(b)}$$

$$n_{H2SO4} = 0,65 \text{ mol } H_2SO_4$$

n<sub>Li</sub> used = 2 x n<sub>H2</sub>so<sub>4</sub> used = 2 x 0,65 = 1,3 mol

$$n = \frac{m}{M}$$

 $1,3 = \frac{m}{7} \sqrt{(d)}$ m = 9,1 g Li used  $n = \frac{1,24}{7}$ 

m = 0,18 mol Li used

Initial mass of Li = 9,1 + 1,24  $\checkmark$   $\checkmark$  (e) Y = 10,34 g  $\checkmark$  (f)

Initial moles of Li = 0,18 + 1,3  $\checkmark \checkmark$  (e) = 1,48 mol

$$n = \frac{m}{M}$$

 $n = \frac{m}{M}$ 

$$1,48 = \frac{m}{7} \sqrt{(d)}$$

 $Y = m = 10,33 \text{ g} \checkmark^{(f)}$ 

(7) **[27]** 

(2)

# Physical desired from Stanmore physics.com AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

### **QUESTION 7**

- An acid is a substance that produces hydrogen ions (H+)/hydronium 7.1.1 (2)ions (H<sub>3</sub>O<sup>+</sup>) when it dissolves in water. ✓✓
- 7.1.2 CH<sub>3</sub>COOH ✓ and H<sub>3</sub>O<sup>+</sup> ✓ (2)
- Weak acid. ✓ It ionises incompletely/partially in water to form a low 7.1.2 (2)concentration of H<sub>3</sub>O<sup>+</sup> ions. ✓

(5)

7.2.1 **OPTION 1** 

$$\frac{C_B V_B}{C_A V_A} = \frac{n_B}{n_A} \checkmark$$

$$\frac{C_B(24,5)\sqrt{}}{(0,15)(31,3)\sqrt{}} = \frac{1}{2}\sqrt{}$$

 $C_B = 0.1 \text{ mol} \cdot \text{dm}^{-3} \checkmark$ 

### **OPTION 2**

$$C = \frac{n}{V} \checkmark$$

$$0,15 = \frac{n}{0,0313}$$

$$n = 4.7 \times 10^{-3} \text{ mol HNO}_3$$

nmorephysics.com HNO<sub>3</sub>: Ca(OH)<sub>2</sub>

2:11

4,7 x 10<sup>-3</sup>: 2,35 x 10<sup>-3</sup> mol Ca(OH)<sub>2</sub>

$$C = \frac{n}{V}$$

$$C = \frac{2,35 \times 10^{-3}}{0,0313}$$

$$C = 0.1 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

tanmorephysics.com

### 7.2.2 Marking criteria:

- (a) Substitution into formula n = cV to calculate initial concentration of HNO<sub>3</sub>.
- (b) Substitution of pH value (2,174) in pH formula.
- (c) Substitution into formula n = cV to calculate final concentration of H₃O⁺.
- (d) Apply ratio n<sub>final</sub>(H<sub>3</sub>O<sup>+</sup>): n<sub>final</sub>(HNO<sub>3</sub>) =1:1 to calculate final moles of HNO<sub>3</sub>.
- (e) Subtraction: n<sub>reacted</sub> (HNO<sub>3</sub>) = n<sub>final</sub> (HNO<sub>3</sub>) n<sub>rnitial</sub> (HNO<sub>3</sub>).
- (f) Apply ratio n<sub>reacted</sub> (Zn) = ½ n<sub>reacted</sub> (HNO<sub>3</sub>) to calculate n<sub>reacted</sub> (Zn).
- (g) Substitute in formula m = nM to calculate mass of Zn that reactd.
- (h) Final answer.

$$\begin{split} n_{\text{initial}}(\text{HNO}_3) &= \text{cV} \\ &= \underline{0,15 \times 0,25} \,\, \checkmark^{(a)} \, = \, 0,0375 \,\, \text{mol} \\ \text{pH} &= -\text{log}[\text{H}_3\text{O}^+] \\ \underline{2,174} &= -\text{log}[\text{H}_3\text{O}^+] \,\, \checkmark^{(b)} \\ \text{[H}_3\text{O}^+] &= 6,7 \times 10^{-3} \,\, \text{mol} \cdot \text{dm}^{-3} \\ n_{\text{final}}(\text{H}_3\text{O}^+) &= \text{cV} \\ &= \underline{6,7 \times 10^{-3} \times 0,25} \,\, \checkmark^{(c)} \!\! = \, 1,675 \times 10^{-3} \,\, \text{mol} \\ n_{\text{final}}(\text{HNO}_3) &= \, 1,675 \times 10^{-3} \,\, \text{mol} \,\, \checkmark^{(d)} \end{split}$$

$$n_{\text{reacted}}(HNO_3) = 0.0375 - 1.675 \times 10^{-3}$$
  $\checkmark^{(e)}$ 

= 0,0358 mol

 $n_{\text{reacted}} (Zn) = \frac{1}{2} (0.0358) \sqrt{(f)}$ = 0.018 mol/s/cs.com

m = nM  
= 
$$0.018 \times 65 \checkmark^{(g)}$$
  
=  $1.16 \text{ g} \checkmark^{(h)}$ 

7.2.3

$$n_{produced}$$
 (H<sub>2</sub>) =  $\frac{1}{2}$   $n_{reacted}$  (HNO<sub>3</sub>) OR 1 x  $n_{reacted}$  (Zn)  
=  $\frac{1}{2}$  (0,0358)  $\checkmark$  OR 1(0,018)  
= 0,018 mol

$$V = nV_m$$

$$= 429.6 \text{ cm}^3 \checkmark (0.4296 \text{ dm}^3)$$

(3) **[22]** 

(8)

# Physical desired from Stanmore physics. com AZULU-NATAL/NOVEMBER 2025 Marking Guidelines

#### **QUESTION 8**

8.1.1 A reaction in which electrons are transferred. 
$$\checkmark\checkmark$$
 (2)

8.1.2 Al  $\checkmark$  (1)

8.1.3 NO<sub>3</sub> ions do not undergo oxidation nor reduction  $\checkmark$  OR
The oxidation numbers stay the same. (1)

8.1.4 The net charge on the left-hand side / for the reactants is not equal to the net charge on the right-hand side / for the products.  $\checkmark\checkmark$  (2)

8.2.1 +6  $\checkmark$  (1)

8.2.2  $Cr_2Or^{2-} + 14H^+ + 6e^- \checkmark \rightarrow 2Cr^{3+} + 7H_2O \checkmark$  (2)

8.2.3  $3Sn^{2+} \rightarrow 3Sn^{4+} + 6e^- \checkmark \rightarrow 2Cr^{3+} + 7H_2O \checkmark$  Balancing  $\checkmark$  (3)
[12]

Stansorephysics.com