



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

## SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS

**MATHEMATICS P1**

**MAY/JUNE 2026** stanmorephysics.com

stanmorephysics.com

**MARKS: 150**

**TIME: 3 hours**

**This question paper consists of 9 pages, 1 information sheet and  
an answer book of 21 pages.**



**INSTRUCTIONS AND INFORMATION.**

Read the following instructions and information carefully before answering the questions.

1. This question paper consists of 10 questions.
2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
3. Clearly show ALL calculations, diagrams, graphs, etc. that you have used in determining your answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round off answers correct to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. An information sheet with formulae is included at the end of the question paper.
9. Write neatly and legibly.



## QUESTION 1

1.1 Solve for  $x$ :

1.1.1  $x^2 - 9 = 0$  (2)

1.1.2  $2x^2 + 3x - 7 = 0$  (correct to TWO decimal places) (3)

1.1.3  $-x^2 + 3x > -4$  (4)

1.1.4  $\sqrt{7x+2} + 2x = 8$  (4)

1.2 Solve simultaneously for  $x$  and  $y$ :

$$\frac{2^x \cdot 2^y}{4} = 1 \quad \text{and} \quad -2x^2 + 3xy = y^2$$
 (6)

1.3 Given:  $(\sqrt{6p} + \sqrt[3]{128})(\sqrt{6p} - \sqrt[3]{128}) = 54 + a\sqrt{2}$ Calculate  $a - p$ .(4)  
[23]

## QUESTION 2

2.1 Given the arithmetic series:  $6 + 10 + 14 + \dots$ 

2.1.1 Which term in the series has the value of 602? (2)

2.1.2 Show that  $S_n = 2n^2 + 4n$  (2)2.1.3 Calculate the value of  $\sum_{n=100}^{150} (4n + 2)$  (2)2.2 The first four terms of a quadratic number pattern are:  $-1$  ;  $-4x$  ;  $11$  ;  $2x + 12$ 2.2.1 Calculate the value of  $x$ . (4)2.2.2 If  $x = -2$ , show that  $T_n = -3n^2 + 18n - 16$  (3)2.2.3 Consider the first  $n$  terms of the quadratic number pattern. Calculate the least value of  $n$  for which the sum of the first differences will be less than  $-1\,380$ .(4)  
[17]

**QUESTION 3**

Two clocks, A and B, were observed from 09:00 to 10:00.

The following observations were made:

- The time on both clocks was exactly 09:00.
- Clock A measured time correctly from 09:00 to 10:00.
- Clock B measured half the time that clock A measured in the first five minutes.
- Thereafter, for every five minutes, clock B measured 50% of the time it measured in the previous 5 minutes.



3.1 What was the time on clock B when clock A reached 10:00? (5)

3.2 How long will it take clock B to reach the time of 09:06?  
Justify your answer. (2)




[7]



**QUESTION 4**

Sketched below are the graphs of  $f(x) = -a^x + q$  and  $g(x) = -x^2 + 2x + 3$ .

B and C are the  $x$ -intercepts of  $g$ . D, the turning point of  $g$ , is also a point on the asymptote of  $f$ . T is a point on  $f$  such that DT is parallel to the  $y$ -axis.



- 4.1 Calculate the coordinates of D. (3)
- 4.2 Write down the values of  $x$  where  $g$  is increasing. (1)
- 4.3 Calculate the  $x$ -intercepts of  $g$ . (3)
- 4.4 Determine the values of  $k$  such that  $g(x) = k$  has TWO distinct positive roots. (2)
- 4.5 Write down the range of  $f$ . (1)
- 4.6 If  $DT = \frac{3}{2}$  units, calculate the values of  $a$  and  $q$ . (4)
- 4.7 Why is  $f$  a function? (1)
- 4.8 It is further given that  $f(x) = -\left(\frac{3}{2}\right)^x + 4$  and  $h(x) = -f(x) + 4$ . Determine the equation of  $h^{-1}$ , the inverse of  $h$ , in the form  $y = \dots$  (3)
- 4.9  $T'$  is the image of T on  $h$ . Sketch the graph of  $h^{-1}(x)$ . Indicate clearly on your graph the intercepts with the axes, the coordinates of  $T''$ , the image of  $T'$  on  $h^{-1}(x)$  and any asymptotes. (3)


[21]



**QUESTION 5**

The graphs of  $f(x) = \frac{-6}{x+2} - 1$  and  $g(x) = \frac{1}{2}x + 4$  are drawn below.

A and B are the x- and y-intercepts of  $f$  respectively. A is also a point of intersection of  $f$  and  $g$ .



- 5.1 Write down the domain of  $f$ . (1)
  - 5.2 Calculate the coordinates of:
    - 5.2.1 B (1)
    - 5.2.2 A (2)
  - 5.3  $h$  is the axis of symmetry of  $f$  that has a positive gradient. Determine the equation of  $h$  in the form  $h(x) = \dots$  (2)
  - 5.4 Calculate the coordinates of the point of intersection of  $g$  and  $h$ . (3)
  - 5.5 Use the graphs to determine the values of  $x$  for which  $f(x) + g(x) \geq 0$  (3)
  - 5.6 A right-angled triangle is created, having A as one of its vertices,  $D(k; 0)$  as another vertex and  $g$  as one of its sides. Calculate the values of  $k$  when the area of the right-angled triangle is 64 units<sup>2</sup>. (4)
- [16]**



## QUESTION 6

- 6.1 On 1 January 2025, the balance in Ben's savings account was R75 000. This savings account earns interest at  $r\%$  p.a., compounded half-yearly. Calculate the value of  $r$  if Ben will have R85 000 in the account on 1 January 2027. (3)
- 6.2 William wants to buy a house. The bank requires William to pay a deposit of 10% of the purchase price of the house. The bank will grant William a loan to cover the outstanding amount, subject to the following conditions:
- The loan is to be repaid over 20 years.
  - Interest on the loan is fixed at 11,5% p.a., compounded monthly.
  - An instalment of R10 665 is payable at the end of each month after the loan has been granted, starting one month after the loan has been granted.
- 6.2.1 Calculate the deposit required on the purchase price of the house. (5)
- 6.2.2 After negotiating with the seller, it was agreed that the purchase price of the house will be R1 100 000. William decides to pay a deposit of R120 000 and a monthly instalment of R11 000 at the end of each month, starting one month after the loan has been granted.
- (a) How long (in years) will it take for him to pay off the loan? (5)
- (b) How much will William have paid in total towards the purchase of the house at the end of 16 years after the loan was granted? (3)
- [16]**


## QUESTION 7

- 7.1 Determine  $f'(x)$  from first principles if it is given that  $f(x) = x^2 + 2x$  (5)
- 7.2 Determine:
- 7.2.1  $f'(x)$  if  $f(x) = x^2 - x - 6$  (2)
- 7.2.2  $\frac{d}{dx} \left( \sqrt[3]{27x^4} + \frac{2}{x^3} \right)$  (4)
- 7.3 The graph of  $k(x) = 2x^3 + bx^2 + cx + d$  has turning points at  $x = -2$  and  $x = 0$ . Determine the values of  $x$  for which:
- 7.3.1  $k'(x) = 0$  (1)
- 7.3.2  $k''(x) < 0$  (2)
- 7.3.3  $k'(x-3) > 0$  (2)
- [16]**



**QUESTION 8**

The graph of  $h(x) = -x^3 + px^2 + qx + r$  is drawn below.  $C(2; 0)$  and  $D(5; 0)$  are the  $x$ -intercepts and  $B$  is the  $y$ -intercept of  $h$ .  $A$  and  $C$  are the turning points of  $h$ .



- 8.1 Show that  $p = 9$ ,  $q = -24$  and  $r = 20$  (3)
- 8.2 Write down the length of  $OB$ . (1)
- 8.3 Calculate the coordinates of the point of inflection of  $h$ . (4)
- 8.4 Determine the equation of the tangent to  $h$  that has an angle of inclination of  $96,34^\circ$  and touches  $h$  between  $B$  and  $C$ . (5)

[13]

**QUESTION 9**

The oxygen content in the water,  $t$  days after organic waste was dumped into a lake, can be represented by  $C(t) = \frac{1}{1200}(t^3 - 15t^2 + 1200)$  for  $0 \leq t \leq 15$

- 9.1 What was the initial oxygen content in the lake at the time that organic waste was dumped? (2)
- 9.2 When will the oxygen content be at the lowest level? (3)
- 9.3 Calculate the value of  $t$  for which the rate of change of oxygen content is a maximum. (2)

[7]



**QUESTION 10**

10.1 Dr Calculus's wardrobe has:

TIES	TROUSERS	SHIRTS
6 blue	1 blue	3 red
4 red	1 black	2 pink
5 green	1 brown	4 white
	1 grey	3 blue
		4 green

10.1.1 An outfit comprises a tie, trousers and shirt. How many different outfits are possible from the items in the wardrobe? (1)

10.1.2 How many different outfits are possible if Dr Calculus does not wear red and pink at the same time? (2)

10.1.3 Calculate the probability that only blue and black items will be used in an outfit. (2)

10.2 A survey was conducted among 54 people to determine which of the three types of tea (A, B or C) they drank. The results are summarised below.

- Some people did not drink any tea at all.
- $P(\text{drinking types A and B and C}) = \frac{7}{54}$
- 9 people drank types B and C.
- 2 people drank types A and B but not type C.
- 18 people drank type B.
- No-one drank types A and C but not type B.
- Drinking type A is independent from drinking type B.
- $P(\text{not drinking type (A or B)}) = 2P(\text{not drinking types (A or B or C)})$ .

10.2.1 Calculate how many people drank type A tea. (3)

10.2.2 Use the given information to complete the Venn diagram provided in the ANSWER BOOK. (6)  
[14]

**TOTAL: 150**



**INFORMATION SHEET**

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d]$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; -1 < r < 1$$

$$F = \frac{x[(1 + i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1 + i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cdot \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2}ab \cdot \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2 \sin \alpha \cdot \cos \alpha$$

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$

