



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

## SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS

**PHYSICAL SCIENCES: CHEMISTRY (P2)**

**MAY/JUNE 2026**

Stanmorephysics.com

Stanmorephysics.com

**MARKS: 150**

**TIME: 3 hours**

**PHYSICAL SCIENCES: Paper 2**



10842E

**X10**



This question paper consists of 16 pages and 4 data sheets.



**INSTRUCTIONS AND INFORMATION.**

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. Show ALL formulae and substitutions in ALL calculations.
8. Round off your FINAL numerical answers to a minimum of TWO decimal places.
9. Give brief motivations, discussions, etc. where required.
10. You are advised to use the attached DATA SHEETS.
11. Write neatly and legibly.



**QUESTION 1: MULTIPLE-CHOICE QUESTIONS**

Various options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, e.g. 1.11 E.

1.1 Consider the following reaction.



This is an example of a/an ... reaction.

- A cracking
- B addition
- C hydrolysis
- D substitution

(2)

1.2 Four organic compounds are shown below.

<b>I</b>	$\begin{array}{c} \text{O} \quad \text{CH}_3 \\ \parallel \quad   \\ \text{CH}_2 - \text{C} - \text{CH} \\   \quad \quad   \\ \text{CH}_3 \quad \quad \text{CH}_3 \end{array}$	<b>II</b>	$\begin{array}{c} \text{CH}_3 \quad \text{O} \\   \quad \parallel \\ \text{CH}_3 - \text{C} - \text{C} - \text{CH}_3 \\   \\ \text{CH}_3 \end{array}$
<b>III</b>	$\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_3 - \text{CH} - \text{C} - \text{CH}_3 \\   \\ \text{CH}_3 - \text{CH}_2 \end{array}$	<b>IV</b>	$\begin{array}{c} \text{O} \quad \text{CH}_3 \\ \parallel \quad   \\ \text{CH}_3 - \text{CH} - \text{C} - \text{CH}_2 \\   \\ \text{CH}_3 \end{array}$


Which of the compounds above have the same IUPAC name?

- A I and II
- B II and III
- C I and IV
- D I and III

(2)



- 1.3 Consider the equation for the reaction below. **R** is the MAJOR PRODUCT.




Which ONE of the following is the CORRECT combination for the type of reaction and the formula of **R**?

	TYPE OF REACTION	FORMULA OF R
A	Addition	$\text{CH}_3\text{CH}_2\text{CHBrCH}(\text{CH}_3)\text{CH}_3$
B	Substitution	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CBr}(\text{CH}_3)\text{CH}_3$
C	Addition	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CBr}(\text{CH}_3)\text{CH}_3$
D	Substitution	$\text{CH}_3\text{CH}_2\text{CHBrCH}(\text{CH}_3)\text{CH}_3$

(2)

- 1.4 The Maxwell-Boltzmann distribution curve for a sample of a gas in a container is shown below. Stanmorephysics.com




Which ONE of the following will change the area under the curve?

- A The addition of a catalyst
- B A decrease in temperature
- C Gas particles escaping from the container
- D A decrease in concentration by increasing the volume of the container


(2)



1.5 The reaction below reaches equilibrium in a closed system.



A change is made to the system at time  $t_1$ . The reaction rate versus time graph is given below.



Which ONE of the following changes was made to the system at time  $t_1$ ?

- A A catalyst was added.
- B The pressure was decreased at constant temperature.
- C The temperature was increased.
- D The concentration of the  $\text{H}_2(\text{g})$  was increased. (2)

1.6 The equilibrium constant,  $K_c$ , for a hypothetical reaction is given by the following expression:

$$K_c = \frac{[\text{A}]^2}{[\text{C}]^3[\text{D}_2]}$$

The equation for this reaction is:

- A  $3\text{C}(\text{g}) + \text{D}_2(\text{g}) \rightleftharpoons 2\text{A}(\text{g}) + \text{B}(\text{s})$
- B  $2\text{A}(\text{g}) + \text{B}(\text{s}) \rightleftharpoons 3\text{C}(\text{g}) + \text{D}_2(\text{g})$
- C  $3\text{C}(\text{g}) + \text{D}_2(\text{g}) \rightleftharpoons 2\text{A}(\text{g}) + \text{B}(\text{g})$
- D  $2\text{A}(\text{l}) + \text{B}(\text{s}) \rightleftharpoons 3\text{C}(\text{g}) + \text{D}_2(\text{g})$  (2)



1.7 According to the Arrhenius theory, an acid is ...

- A a source of  $H^+$  ions in water.  
 B a source of  $OH^-$  ions in water.  
 C a substance that can accept a proton.  
 D a substance that can donate a proton.


(2)

1.8 A property of sodium nitrate solution,  $NaNO_3(aq)$ , is that it is...

- A acidic.  
 B basic.  
 C ampholytic.  
 D neutral.

(2)

1.9 An iron rod,  $Fe(s)$ , is placed in a beaker containing chromium(III) ions,  $Cr^{3+}(aq)$ .



Which ONE of the following is the CORRECT combination of products formed?

- A  $Fe^{2+}(aq)$  and  $Cr(s)$   
 B  $Fe^{2+}(aq)$  and  $Cr^{2+}(aq)$   
 C  $Fe^{3+}(aq)$  and  $Cr^{2+}(aq)$   
 D  $Fe^{3+}(aq)$  and  $Cr(aq)$

(2)



1.10 Which ONE of the following combinations CORRECTLY shows a product formed at the cathode and the change in pH during the electrolysis of a concentrated sodium chloride solution? Stanmorephysics.com

	PRODUCT FORMED AT CATHODE	CHANGE IN pH
A	Hydrogen	Decreases
B	Hydrogen	Increases
C	Chlorine	Decreases
D	Chlorine	Increases

(2)  
[20]

**QUESTION 2 (Start on a new page.)**

The letters **A** to **G** in the table below represent organic compounds.

<b>A</b>	$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{CH}_3 \\   \\ \text{CH}_3-\text{CH}-\text{C}\equiv\text{C}-\text{CH}_3 \end{array}$	<b>B</b>	
<b>C</b>	$\text{C}_5\text{H}_{10}\text{O}_2$	<b>D</b>	Hexanal
<b>E</b>	$\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	<b>F</b>	
<b>G</b>	$\text{C}_5\text{H}_{12}$		

- 2.1 Define the term *saturated compound*. Stanmorephysics.com (2)
- 2.2 Write down the letter(s) that represent(s):
- 2.2.1 A compound with a carbonyl group bonded to two carbon atoms (1)
- 2.2.2 The compound that is an isomer of compound **E** (1)
- 2.2.3 Two unsaturated hydrocarbons (1)
- 2.3 Write down the:
- 2.3.1 IUPAC name of compound **A** (3)
- 2.3.2 IUPAC name of compound **B** (3)



2.4 The molecular formula of compound **C**,  $C_5H_{10}O_2$ , represents two compounds that are functional isomers of each other.

2.4.1 Define the term *functional isomers*. (2)

Write down the:

2.4.2 STRUCTURAL FORMULA for the functional isomer with the stronger intermolecular forces (2)

2.4.3 CONDENSED FORMULA of the functional isomer with the lower boiling point (2)

2.5 Is compound **B** a PRIMARY, SECONDARY or TERTIARY haloalkane?

Give a reason for the answer. (2)


2.6 Using MOLECULAR FORMULAE, write down a balanced equation for the combustion of compound **G** in EXCESS oxygen. (3)

[22]



**QUESTION 3 (Start on a new page.)**

- 3.1 Consider the diagram of two methanol molecules shown below. I and II are forces.



Choose I OR II for EACH of the following:

- 3.1.1 The force, when broken or overcome, results in a chemical change (1)
- 3.1.2 The force that requires more energy to overcome (1)
- 3.1.3 The force responsible for the boiling point of methanol (1)
- 3.2 In an investigation, the boiling points of four compounds are compared. The boiling points of compounds A, B and C are given below. The boiling point of D is indicated as X.

	COMPOUNDS	BOILING POINT (°C)
A	Methanol	64,7
B	Ethanol	78,3
C	Butan-1-ol	117,7
D	Butanal	X

- 3.2.1 Define the term *boiling point*. (2)

The boiling points of compounds A, B and C are compared.

- 3.2.2 With reference to molecular structure, give the reason why this is a fair comparison. (1)
- 3.2.3 Name the intermolecular force responsible for the difference in these boiling points. (1)
- 3.2.4 Give a reason for the increase in boiling points in terms of the molecular structure of the compounds. (1)

The boiling points of compounds C and D are compared.

- 3.2.5 Will X, the boiling point of D, be GREATER THAN, LESS THAN or EQUAL TO 117,7 °C?


Fully explain the answer.

(4)  
[12]



**QUESTION 4 (Start on a new page.)**

Study the flow diagram below. Compounds **A**, **B**, **C**, **D** and **E** are organic compounds.



Reaction 6 is an addition reaction.

4.1 Write down the:

4.1.1 Type of reaction represented by reaction 1 (2)

4.1.2 Reaction condition for reaction 1 (1)

4.2 In reaction 3, compound **C** reacts with  $\text{HCOOH}$ , in the presence of concentrated  $\text{H}_2\text{SO}_4$ .

Write down the:

4.2.1 STRUCTURAL FORMULA of compound **C** (2)

4.2.2 Type of reaction represented by reaction 3 (1)

4.2.3 IUPAC name of compound **D** (2)

4.3 Write down TWO terms for the type of reaction represented by reaction 2. (2)

4.4 Compound **C** can be converted back to compound **B**.

Write down the FORMULA or NAME of the inorganic reagent needed for this conversion. Stanmorephysics.com (1)

4.5 Compounds **B** and **C** can be converted to compound **E**.

4.5.1 Write down ONE term for the type of organic reaction for BOTH reactions 4 and 5. (1)

Besides heat, write down a reaction condition for:

4.5.2 Reaction 4 (1)

4.5.3 Reaction 5 (1)

Consider reaction 4:


4.5.4 Using STRUCTURAL FORMULAE, write down the balanced equation. (4)

[18]




**QUESTION 5 (Start on a new page.)**

Pure powdered copper,  $\text{Cu(s)}$ , is oxidised IN EXCESS concentrated nitric acid,  $\text{HNO}_3(\text{conc.})$ , as shown by the balanced equation below.



- 5.1 Explain what is meant by 'IN EXCESS' in the statement above. (2)
- 5.2 The volume of nitrogen dioxide gas produced is shown in the graph below.



- 5.2.1 Use the collision theory to explain the decrease in the rate of the reaction between  $t_3$  and  $t_4$ . (3)
- 5.2.2 How does the rate of the reaction at  $t_2$  compare to that at  $t_1$ ? Choose from HIGHER THAN, LOWER THAN or REMAINS THE SAME. (1)
- 5.2.3 Give a reason for the answer to QUESTION 5.2.2 by referring to the graph. Stanmorephysics.com (1)
- 5.2.4 Calculate the value of time  $t_3$  if the average rate for the production of  $1,7 \text{ dm}^3$  of  $\text{NO}_2(\text{g})$  is  $0,02 \text{ dm}^3 \cdot \text{s}^{-1}$ . (2)
- 5.2.5 The initial concentration of the nitric acid is  $12 \text{ mol} \cdot \text{dm}^{-3}$ . If  $20 \text{ cm}^3$  of nitric acid is used, calculate the hydronium ion concentration after the reaction is completed. (Assume the volume of the acid does not change.) (6)
- Take the molar gas volume to be  $25 \text{ dm}^3 \cdot \text{mol}^{-1}$ .

A change is made to the reaction conditions and the reaction is repeated.

- 5.2.6 Which ONE of the following changes will result in more  $\text{NO}_2$  gas being produced when the reaction is completed? Choose from A, B or C. (Write down only the letter.)

A	A piece of pure copper with the same mass is used.
B	Impure copper with greater mass is used.
C	Higher concentration $\text{HNO}_3$ in excess is used.

(2)


- 5.3 Determine the oxidation number of nitrogen in  $\text{HNO}_3$ .

(2)


**[19]**

**QUESTION 6 (Start on a new page.)**

Iodine gas,  $I_2(g)$ , reacts with chlorine gas,  $Cl_2(g)$ , in a sealed container at temperature  $T$ . Equilibrium is reached according to the balanced equation below.



The graph below shows the masses of  $I_2(g)$ ,  $Cl_2(g)$  and  $ICl_3(g)$  versus time during the course of the reaction.



- 6.1 State Le Chatelier's principle. (2)
- 6.2 Why are the masses constant after  $t_2$ ? (1)
- 6.3 At  $t_1$ , the pressure is adjusted by changing the volume of the container at constant temperature. (3)
- Was the volume of the container INCREASED or DECREASED?
- Explain the answer. (3)
- 6.4 What other change to the equilibrium system will produce a SIMILAR change to the masses at  $t_1$ ? (2)
- 6.5 The volume of the container is  $2 \text{ dm}^3$ . At  $t_2$  the concentrations of the gases are:

$$\begin{aligned} [I_2] &= 0,15 \text{ mol}\cdot\text{dm}^{-3} \\ [Cl_2] &= 0,36 \text{ mol}\cdot\text{dm}^{-3} \\ [ICl_3] &= 0,05 \text{ mol}\cdot\text{dm}^{-3} \end{aligned}$$


More  $I_2$  is now added to the container. When a new equilibrium is reached, the concentration of  $ICl_3$  is  $0,09 \text{ mol}\cdot\text{dm}^{-3}$ . If the equilibrium constant,  $K_c$ , is  $0,357$ , calculate the number of moles of  $I_2$  added. (8)

[16]



**QUESTION 7 (Start on a new page.)**

7.1 Water ionises according to the balanced equation:



The equilibrium constant for the ionisation of water,  $K_w$ , varies with temperature, as shown in the table below.

TEMPERATURE (°C)	$K_w$
25	$1,00 \times 10^{-14}$
50	$5,48 \times 10^{-14}$

7.1.1 Is the ionisation of water EXOTHERMIC or ENDOTHERMIC?

Explain the answer.

(3)

7.1.2 Calculate the concentration of  $\text{H}_3\text{O}^+$  ions in water at 50 °C.

(2)

7.1.3 How will the pH of water be affected when the temperature increases? Choose from INCREASES, DECREASES or STAYS THE SAME. Give a reason for the answer by referring to the formula used to calculate pH.

(2)

7.2 Water is an ampholyte.

Write down the balanced equation for the reaction of water with  $\text{HCO}_3^-(\text{aq})$  where:

7.2.1 Water acts as an acid

(2)

7.2.2 Water acts as a base

(2)

7.3 The table below shows the equilibrium constants,  $K_a$ , of four acids, each having a concentration of  $0,1 \text{ mol}\cdot\text{dm}^{-3}$  at 25 °C. The acids are used as electrolytes.

ACID	$K_a$
$\text{H}_2\text{CO}_3$	$4,3 \times 10^{-7}$
$\text{CH}_3\text{COOH}$	$1,8 \times 10^{-5}$
$\text{HNO}_2$	$4,6 \times 10^{-4}$
$\text{H}_2\text{C}_2\text{O}_4$	$6,0 \times 10^{-2}$

7.3.1 Define the term *electrolyte*.



(1)

7.3.2 Which ONE of the above acids is the most effective electrolyte? Give a reason for the answer.


(2)



- 7.4 An antacid tablet contains a mixture of  $\text{Al}(\text{OH})_3$  and  $\text{Mg}(\text{OH})_2$  only. This tablet is reacted with  $50 \text{ cm}^3$  of a  $0,5 \text{ mol}\cdot\text{dm}^{-3}$   $\text{HCl}$  solution. The balanced equations for the two reactions that take place are:



The resulting solution is completely neutralised by  $26,5 \text{ cm}^3$  of a  $0,378 \text{ mol}\cdot\text{dm}^{-3}$   $\text{NaOH}$  solution.




- 7.4.1 Calculate the number of moles of hydroxide ions present in the tablet. (5)
- 7.4.2 If the tablet contains an equal number of moles of aluminium hydroxide and magnesium hydroxide, calculate the mass of  $\text{Al}(\text{OH})_3(\text{s})$  in the tablet. (5)

[24]



**QUESTION 8 (Start on a new page.)**

A group of learners set up a standard galvanic cell, as shown below.




- 8.1 Define the term *galvanic cell*. (2)
- 8.2 For this cell, write down the:
- 8.2.1 Cell notation (3)
- 8.2.2 Direction of the electron flow. Choose from **P TO Q** or **Q TO P**. (1)
- 8.2.3 Reduction half-reaction (2)
- 8.3 How will the reading on the voltmeter be affected when the concentration of the chloride ions is increased? Choose from **INCREASES**, **DECREASES** or **REMAINS THE SAME**. (2)
- Give a reason for the answer. [10]



**QUESTION 9 (Start on a new page.)**

An electrolytic cell is set up for the electrolysis of a concentrated copper(II) chloride solution,  $\text{CuCl}_2(\text{conc.})$ . X and Y are carbon electrodes.



- 9.1 Define the term *electrolysis*. (2)
- 9.2 Write down balanced equations for TWO reactions that will take place at the anode. (4)
- 9.3 Write down the NAME or FORMULA of the substance produced in higher amount at electrode Y. (1)
- 9.4 A solution of zinc iodide,  $\text{ZnI}_2$ , is added to the electrolyte of this cell. What product(s) will be formed at:
- 9.4.1 Electrode X (1)
- 9.4.2 Electrode Y (1)
- [9]

**TOTAL: 150**

DATA FOR PHYSICAL SCIENCES GRADE 12  
PAPER 2 (CHEMISTRY)

GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12  
VRAESTEL 2 (CHEMIE)

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure Standaarddruk	$p^\theta$	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP Molêre gasvolume by STD	$V_m$	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature Standaardtemperatuur	$T^\theta$	273 K
Charge on electron Lading op elektron	$e$	$1,6 \times 10^{-19} \text{ C}$
Avogadro's constant Avogadro-konstante	$N_A$	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} \text{ at/by } 298 \text{ K}$	
$E_{\text{cell}}^\theta = E_{\text{cathode}}^\theta - E_{\text{anode}}^\theta / E_{\text{sel}}^\theta = E_{\text{katode}}^\theta - E_{\text{anode}}^\theta$	
or/of $E_{\text{cell}}^\theta = E_{\text{reduction}}^\theta - E_{\text{oxidation}}^\theta / E_{\text{sel}}^\theta = E_{\text{reduksie}}^\theta - E_{\text{oksidasie}}^\theta$	
or/of $E_{\text{cell}}^\theta = E_{\text{oxidising agent}}^\theta - E_{\text{reducing agent}}^\theta / E_{\text{sel}}^\theta = E_{\text{oksideermiddel}}^\theta - E_{\text{reduseermiddel}}^\theta$	





Downloaded from [www.stannorephysics.com](http://www.stannorephysics.com)  
 TABLE 4A: STANDARD REDUCTION POTENTIALS  
 TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE



Increasing strength of oxidising agents/Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies	$E^{\theta}$ (V)
$F_2(g) + 2e^- = 2F^-$	+ 2,87
$Co^{3+} + e^- = Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- = 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^- = Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- = 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- = 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- = 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- = Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- = Pt$	+ 1,20
$Br_2(l) + 2e^- = 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- = NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- = Hg(l)$	+ 0,85
$Ag^+ + e^- = Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- = NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- = Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- = H_2O_2$	+ 0,68
$I_2 + 2e^- = 2I^-$	+ 0,54
$Cu^+ + e^- = Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- = S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- = 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- = Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- = SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- = Cu^+$	+ 0,16
$Sn^{4+} + 2e^- = Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- = H_2S(g)$	+ 0,14
$2H^+ + 2e^- = H_2(g)$	0,00
$Fe^{3+} + 3e^- = Fe$	- 0,06
$Pb^{2+} + 2e^- = Pb$	- 0,13
$Sn^{2+} + 2e^- = Sn$	- 0,14
$Ni^{2+} + 2e^- = Ni$	- 0,27
$Co^{2+} + 2e^- = Co$	- 0,28
$Cd^{2+} + 2e^- = Cd$	- 0,40
$Cr^{3+} + e^- = Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^- = Fe$	- 0,44
$Cr^{3+} + 3e^- = Cr$	- 0,74
$Zn^{2+} + 2e^- = Zn$	- 0,76
$2H_2O + 2e^- = H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^- = Cr$	- 0,91
$Mn^{2+} + 2e^- = Mn$	- 1,18
$At^{3+} + 3e^- = At$	- 1,66
$Mg^{2+} + 2e^- = Mg$	- 2,36
$Na^+ + e^- = Na$	- 2,71
$Ca^{2+} + 2e^- = Ca$	- 2,87
$Sr^{2+} + 2e^- = Sr$	- 2,89
$Ba^{2+} + 2e^- = Ba$	- 2,90
$Cs^+ + e^- = Cs$	- 2,92
$K^+ + e^- = K$	- 2,93
$Li^+ + e^- = Li$	- 3,05

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels



Downloaded from [Stanmorephysics.com](http://Stanmorephysics.com)

TABLE 4B: STANDARD REDUCTION POTENTIALS

TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE



Increasing strength of oxidising agents / Toenemende sterkte van oksideermiddels

Half-reactions/Halfreaksies	$E^{\ominus}$ (V)
$\text{Li}^+ + \text{e}^- \rightleftharpoons \text{Li}$	-3,05
$\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$	-2,93
$\text{Cs}^+ + \text{e}^- \rightleftharpoons \text{Cs}$	-2,92
$\text{Ba}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ba}$	-2,90
$\text{Sr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sr}$	-2,89
$\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$	-2,87
$\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$	-2,71
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2,36
$\text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al}$	-1,66
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	-1,18
$\text{Cr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cr}$	-0,91
$2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	-0,76
$\text{Cr}^{3+} + 3\text{e}^- \rightleftharpoons \text{Cr}$	-0,74
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	-0,44
$\text{Cr}^{3+} + \text{e}^- \rightleftharpoons \text{Cr}^{2+}$	-0,41
$\text{Cd}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cd}$	-0,40
$\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co}$	-0,28
$\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$	-0,27
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	-0,14
$\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$	-0,13
$\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$	-0,06
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+0,15
$\text{Cu}^{2+} + \text{e}^- \rightleftharpoons \text{Cu}^+$	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + \text{e}^- \rightleftharpoons \text{Cu}$	+0,52
$\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2$	+0,68
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+0,80
$\text{Hg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Hg}(\text{l})$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$	+1,07
$\text{Pt}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pt}$	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + \text{e}^- \rightleftharpoons \text{Co}^{2+}$	+1,81
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-$	+2,87

Increasing strength of reducing agents / Toenemende sterkte van reduceermiddels

